drug treatment with counselling, social work, or other psychological and social interventions. Good evidence from studies in psychiatric patients suggests that such psychosocial therapies combine well with drug treatment and are indicated for some psychological problems.⁸

E S PAYKEL

Department of Psychiatry,
University of Cambridge,
Addenbrooke's Hospital,
Cambridge CB2 2QQ

P FREELING

Division of General Practice and Primary Care,
St George's Hospital Medical School,
London SW17 0RE

- 1 Scott AIF, Freeman CPL. Edinburgh primary care depression study: treatment outcome, patient satisfaction, and cost after 16 weeks. *BMJ* 1992;304:883-7. (4 April.)
- 2 Paykel ES, Hollyman JA, Freeling P, Sedgwick P. Predictors of therapeutic benefit from amitriptyline in mild depression: a general practice placebo-controlled trial. *J Affective Disord* 1988;14:83-95.
- 3 Stewart JW, Quitkin FM, Liebowitz MR, McGrath PJ, Harrison WM, Klein DF. Efficacy of desipramine in depressed outpatients: response according to Research Diagnostic Criteria and severity of illness. *Arch Gen Psychiatry* 1983;40:202-7.
- 4 Blashki TG, Mowbray R, Davies B. Controlled trial of amitriptyline in general practice. *BMJ* 1971;1:133-8.
- 5 Thomson J, Rankin H, Ashcroft GW, Yates CM, McQueen JK, Cummings SW. The treatment of depression in general practice: a comparison of L-tryptophan, amitriptyline, and a combination of L-tryptophan and amitriptyline, with placebo. *Psychol Med* 1982;12:741-51.
- 6 Thompson C, Thompson CM. The prescribing of antidepressants in general practice. II. A placebo-controlled trial of low-dose dothiepin. *Human Psychopharmacology* 1989;4:191-204.
- 7 Porter AMW. Depressive illness in a general practice. A demographic study and a controlled trial of imipramine. *BMJ* 1970;1:773-8.
- 8 Paykel ES. Treatment of depression: the relevance of research for clinical practice. *Br J Psychiatry* 1989;155:754-63.

Measuring temperatures

SIR,—In his paper on using thermometers in general practice Steven Clarke shows the wide variation in the practice of temperature taking and identifies the problems associated with using the axillary temperature.¹ He and Duncan Keeley, in his editorial,² concentrate on the problem of detecting fever in children; another group in which measuring temperature may be difficult is elderly people.

We and colleagues have shown that the temperature of an afebrile elderly person varies according to the site of measurement.³ Simultaneously measured rectal and axillary temperatures differed by a mean of 0.91°C (95% confidence interval 0.87 to 0.95°C). Whereas a rectal temperature of 37.6°C was outside our normal range and signified fever, the equivalent figure for an axillary temperature was 37.1°C. The picture is further confused by our finding that the difference in the temperature measured in the auditory canal and sublingually was increased if patients with confusion, previous stroke, parkinsonism, and micrognathia or patients who were not wearing their dentures were included in the normal afebrile population. Mouth breathing, difficulties in maintaining the position of the thermometer, and the ambient temperature may influence the measurement.

The environment in a patient's home is different from that in a warm hospital ward. In unselected elderly patients the mean change in rectal temperature in the 24 hours after admission to hospital was 0.4°C.⁴ In our study 61% of patients who had a low or normal body temperature on admission had a raised temperature at one or more sites the next day.

The rectal temperature and temperature in the proximal auditory canal will detect fever in roughly 86% of febrile elderly patients, the sublingual temperature in 66%, and the axillary temperature in 32%.⁵ If important clinical decisions are to be based on a patient's temperature the limitations of the various methods of measurement must be understood. Keeley's advice to forget the axilla is

as important for elderly people as it is for the young.

JULIUS WEINBERG

Department of Public Health Medicine,
East Berkshire Health Authority,
Windsor,
Berkshire SL4 3AW

ADAM DAROWSKI

Department of Medicine for the Elderly,
Upton Hospital,
Slough,
Berkshire SL1 2BL

- 1 Clarke S. Use of thermometers in general practice. *BMJ* 1992;304:961-3. (11 April.)
- 2 Keeley D. Taking infants' temperatures. *BMJ* 1992;304:931-2. (11 April.)
- 3 Darowski A, Najim Z, Weinberg JR, Guz A. Normal rectal, auditory canal, sublingual and axillary temperatures in elderly afebrile patients in a warm environment. *Age Ageing* 1991;21:113-9.
- 4 Darowski A, Najim Z, Weinberg JR, Guz A. The increase in body temperature of elderly patients in the first twenty-four hours following admission to hospital. *Age Ageing* 1991;21:107-12.
- 5 Darowski A, Najim Z, Weinberg JR, Guz A. The febrile response to mild infections in elderly hospital inpatients. *Age Ageing* 1991;21:193-8.

Ciprofloxacin in bacterial diarrhoea

SIR,—Y J Drabu and colleagues report on a patient infected with *Salmonella virchow* who displayed reduced susceptibility to ciprofloxacin after repeat dosing over 15 months.¹ The authors state, "Ciprofloxacin is not licensed for the treatment of bacterial diarrhoea." This is incorrect, as can be seen in the current datasheet, which indicates that ciprofloxacin may be used to treat enteric fever and infective diarrhoea. The datasheet also states that such organisms as *Salmonella* spp, *Campylobacter coli*, *C jejuni*, and *Escherichia coli* are fully sensitive to ciprofloxacin.

B J O'KEEFFE

Bayer,
Newbury,
Berkshire RG13 1JA

- 1 Drabu YJ, Sharma AK, Yardumian A, Blakemore PH. Antibiotic sensitivity testing in enteric salmonella. *BMJ* 1992;304:686. (14 March.)

Childhood mortality after a high dose of vitamin A

SIR,—Mike Lavender and Chris Vickery question whether vitamin A supplementation could be readily integrated into community health programmes, as colleagues and I suggested in our paper.^{1,2} I acknowledge differences between our programme and the basic health services found elsewhere in Nepal, which were necessitated by the requirements of rigorous documentation. I wish, however, to highlight elements in our programme's design that enhance the possibility of replicating it.

In the more than two years since our study vitamin A supplementation in Jumla has been carried out every four months by intermittent mass coverage of all children at central points in their communities—similar to patterns established by the existing expanded programme on immunisation. Coverage has remained high. This does not rely on the two weekly household visiting regimen of our pneumonia case management programme, which has been running concurrently but separately. Each worker does about 100 children a day and could easily do more in more densely populated areas. Such routine periodic contacts are certainly operationally feasible and contrast with the difficulties cited by Lavender and Vickery in sustaining routine house visiting programmes.

The fact that no other nutritional services were available to our population is an argument for the programme being able to be replicated. Many

primary health care programmes include elaborate components on nutrition education and growth monitoring, but, as Lavender and Vickery imply when they comment on the low levels of routine household contact with health personnel, such components are generally poorly executed. The programme we described has strength because it does not rely on this; we strongly disagree with Lavender and Vickery's claim that such an intervention should not be considered in the absence of a more comprehensive nutrition education programme, which adds great operational complexity. This is an appropriate long term goal, but first things first.

Our vital events monitoring system was entirely separate from the service delivery system, precisely to ensure that the intensity of monitoring would have no bearing on the delivery of services. The total cost of the programme per vitamin A dose delivered was under \$0.20; in most places such a cost is affordable.

The reduction in mortality of 26% in our study is modest compared with reductions found in other, less extreme settings. In Sarlahi, Nepal, the protective effect of periodic vitamin A supplementation was 30%, and in southern India weekly supplementation resulted in a 54% decline in mortality.^{3,4}

Our argument that this approach could be readily replicated does not imply that effort would not be required; our longstanding work with health services throughout Nepal and the proved effect on reducing deaths of children lead us to conclude, however, that this effort is feasible and worth while.

NILS M P DAULAIRE

INTERCEPT,
PO Box 168,
Hanover,
New Hampshire 03755,
USA

- 1 Lavender M, Vickery C. Childhood mortality after a high dose of vitamin A. *BMJ* 1992;304:640. (7 March.)
- 2 Daulaire NMP, Starbuck ES, Houston RM, Church MS, Stuckel TA, Pandey MR. Childhood mortality after a high dose of vitamin A in a high risk population. *BMJ* 1992;304:207-10. (25 January.)
- 3 West KP, Pokhrel RP, Katz J, LeClerq SC, Khatri SK, Shrestha SR, et al. Efficacy of vitamin A in reducing preschool child mortality in Nepal. *Lancet* 1991;338:67-71.
- 4 Rahmathullah L, Underwood BA, Thulasiraj RD, Milton RC, Ramaswamy K, Rahmathullah R, et al. Reduced mortality among children in southern India receiving a small weekly dose of vitamin A. *N Engl J Med* 1990;323:929-35.

Asthma in general practice

SIR,—Cedrick R Martys expresses disappointment at the inability of the Darley Dale asthma clinic¹ to show appreciable changes in morbidity compared with the Aylsham nurse run asthma clinic.^{2,3} Several fundamental differences between the two clinics are worth highlighting.

Our clinic was for patients who were already receiving prophylactic treatment. They were judged to have more severe disease and hence most to gain. The Darley Dale patients were selected on the basis of having asthma and would have included patients with fairly mild asthma and perhaps little to gain from an intensive asthma programme.

Our study was based on those patients who actually attended the clinic whereas the Darley Dale clinic was judged on the total population with asthma regardless of whether they attended the clinic. It seems unfair to judge a clinic by patients who did not attend, just as it is unrealistic to expect general practitioners to use the clinic protocol (which may take 45 minutes) in their 5-10 minute consultations.

Hilton et al and Jenkinson et al have shown that providing patients with knowledge does little to alter morbidity.^{4,5} Our behavioural approach, incorporating written self management plans, moves the control away from the doctor towards

the patients, giving them permission to change treatment and, in particular, to increase inhaled steroids as their lung function falls.

Martys misquotes our study in saying that the peak flow meter is not crucial to managing asthma. We considered that both peak flow meters and symptoms play an important part in self management by teaching the patients the importance of their symptoms and what action they should take.⁶ This is best achieved by written self management plans linked to diary cards and reinforced by colour coded labels on the peak flow meter.⁷

The Darley Dale clinic saw 78 patients in one year with six monthly review. Our clinic ran for nine hours a week and monitored 115 patients in a year with a median of four consultations with the nurse and one consultation with a doctor. The overall consultation rate did not change, but the redistribution in work resulted in a halving of emergency steroid courses, nebulisations, and absenteeism.

Martys recognises that gathering information about a patient's condition does not necessarily change the management of asthma. Unless we can develop techniques that result in steroids being taken in the right doses and at the right time then little is likely to change.

IAN CHARLTON
GILLIAN CHARLTON

Kincumber 2251,
New South Wales,
Australia

- 1 Martys C. Asthma care in Darley Dale: general practitioner audit. *BMJ* 1992;304:758-60. (21 March.)
- 2 Charlton IC, Charlton G, Broomfield J, Mullee MA. Audit of the effect of a nurse run asthma clinic on workload and patient morbidity in a general practice. *Br J Gen Pract* 1991;41:227-31.
- 3 Charlton I, Charlton G, Broomfield J, Campbell M. An evaluation of a nurse-run asthma clinic in general practice using an attitudes and morbidity questionnaire. *Fam Pract* (in press).
- 4 Hilton S, Sibbald B, Anderson HR, Freeling P. Controlled evaluation of the effects of patient education on asthma morbidity in general practice. *Lancet* 1986;i:26-9.
- 5 Jenkinson D, Davison J, Jones S, Hawtin P. Comparison of effects of a self management booklet and audiocassette for patients with asthma. *BMJ* 1988;297:267-70.
- 6 Charlton I, Charlton G, Broomfield J, Mullee M. Evaluation of peak flow and symptoms only self-management plans for control of asthma in general practice. *BMJ* 1990;301:1355-9.
- 7 Charlton I, Charlton G. New perspectives in asthma care. *Practitioner* 1990;234:30-2.

Videotaped interviews with children suspected of being sexually abused

SIR,—M R Wiseman and colleagues report that agreement between professional groups and a consensus rating was good when raters were asked to assess the likelihood of child sexual abuse on the basis of videorecorded interviews with children.¹ There are several reasons, however, for thinking that their study overestimates the level of agreement that will generally be found during such assessments.

Firstly, the design of the research tends to maximise the chances of agreement since, as they point out, the cases were not randomly selected and the professionals rating the videos were highly experienced.

Secondly, their study of videotaped interviews is mainly a test of the reliability of the criteria used to define the probability of sexual abuse. In clinical practice there will be additional variability arising from differences in the amount and quality of other information that raters have (such as direct interviews and statements). Such information was available only to the consensus panel in the authors' study.

Thirdly, the statistic that the authors use to assess agreement, the raw percentage agreement, does not correct for the fact that 50% agreement

would occur by chance. If the data in their first table are reanalysed with a statistic that corrects for chance agreement, the κ statistic, then the level of agreement for all groups is 0.71, less than the 0.85 raw agreement calculated from the same data.

Thus the level of agreement in routine clinical practice is most unlikely to be any higher than the authors report, and for the reasons given above it will probably be much lower. Even if the level of agreement in practice is just 0.2 lower, at a κ coefficient of 0.51, there would be disagreement on a considerable number of cases. For instance, two professionals rating 100 videotapes at this level of agreement (with a "true" rate of high likelihood cases the same as in the reported study) would disagree on 24 cases. Thus more work seems to be needed on standardising the assessment of children suspected of being sexually abused.

RICHARD HARRINGTON

University Department of Psychiatry,
Queen Elizabeth Psychiatric Hospital,
Birmingham B15 2QZ

- 1 Wiseman MR, Vizard E, Bentovim A, Leventhal J. Reliability of videotaped interviews with children suspected of being sexually abused. *BMJ* 1992;304:1089-91. (25 April.)

Being a house officer

SIR,—Most of us prefer to forget about our house jobs as soon as we reach the relative sanctuary of a senior house officer post. We each spent our preregistration year working in different institutions in different parts of Britain, yet we found that John Lear's comments described accurately the misery and frustration of our own experiences as junior house officers¹; we would be surprised if many other readers did not have the same response. The comments of Lear's consultants, with their implication that he was out of the ordinary in having problems, show their perception that the fault lies with the people rather than with the system within which they work. Junior doctors are still not listened to when they are honest and brave enough to express their feelings of inadequacy.

If we are not to train another generation of doctors who become disillusioned so quickly with their first experience of work, two changes are necessary: firstly, medical undergraduate training must prepare new doctors for the job they have to do when they qualify instead of trying to produce brilliant diagnosticians; and, secondly, senior doctors need to take their juniors' complaints seriously and lobby on their behalf for more humane conditions of work.

JEREMY GRIMSHAW

Department of General Practice,
University of Aberdeen,
Medical School,
Aberdeen AB9 2ZD

BRENDA WILSON
DANNY RUTA

Department of Public Health,
University of Aberdeen

- 1 Lear J. The beginning of the end. *BMJ* 1992;304:1122. (25 April.)

SIR,—I am intrigued that at the end of John Lear's personal view the *BMJ* decided to publish a response from his consultants.¹ The response seems to cast doubt on the veracity of Lear's eloquent description of the difficulties he faced as a house officer. The suggestion that he has been economical with the truth tends to diminish the impact of the message he is conveying. In the short response from the anonymous consultants I am reminded obliquely of the importance of a referee and a referee. Unfortunately, this may have the effect of discouraging other disaffected preregistration house officers from speaking out.

I think it should concern all in the medical profession that the consultants wrote, "The patients were as well 'protected' during this time as we could devise." The crucial issue must be the protection of house officers. Recognition that the preregistration year is, and must remain, primarily a training experience would ensure that house officers no longer had to "survive" a traumatic first year in their chosen career. Awareness has been raised; change has surely got to occur.

O JUNAID

Department of Health Care of the Elderly,
University Hospital,
Queen's Medical Centre,
Nottingham NG7 2UH

- 1 Lear J. The beginning of the end. *BMJ* 1992;304:1122. (25 April.)

SIR,—John Lear speaks for many in his personal view on being a house officer.¹ The open acknowledgment of the difficulties of this challenging year allows for change.

Stress in students and junior doctors may affect their physical, psychological, and social health as well as impair their performance as doctors.^{2,3} We are looking at the experience of house officers with a view to preparing students for this extraordinary year. Key topics that we have identified as being appropriate for training are clinical skills,⁴ personal support,⁵ and management skills at both the personal and the professional level.⁶

This summer we plan to run the first of a series of workshops for our clinical students, entitled "Being a doctor: are you ready?" We would be interested to hear from other groups running similar or alternative projects.

C VAUGHAN
S PEZESHGI

Department of General Practice and Primary Care,
King's College School of Medicine and Dentistry,
London SE5 9PJ

- 1 Lear J. The beginning of the end. *BMJ* 1992;304:1122. (25 April.)
- 2 Dowling S, Barrett S. *Doctors in the making. The experience of the pre-registration year.* Bristol: Bristol University, 1991.
- 3 Dyer C. Manslaughter convictions for making mistakes. *BMJ* 1991;303:1218.
- 4 Moss F, McManus IC. The anxieties of new clinical students. *Med Educ* 1992;26:17-20.
- 5 Firth-Cozens J. Emotional distress in junior house officers. *BMJ* 1987;295:533-6.
- 6 Calman KC, Donaldson M. The pre-registration house officer year: a critical incident study. *Med Educ* 1991;25:51-9.

Maternity services

SIR,—Correspondents are critical of the report of the Commons select committee aiming at empowering women to have more choice and control over the place and style of birth.¹ Indeed, it would be surprising if the medical profession did not oppose recommendations to "demedicalise" a service that doctors have tried to monopolise for generations. But the select committee's findings reflect accurately both the needs of women and the views of enlightened health professionals.

Paul Sackin misses the point in claiming that the report is unrealistic in its expectation that all general practices can offer a home birth service.¹ Though it is true that many general practitioners do not wish to provide intrapartum care themselves, they can and should offer every woman a genuine choice (which includes home birth for women considered to be at low risk), referring her to colleagues if necessary.

Adam Balen and John McGarry are shroud waving.¹ A growing body of evidence shows that care by a general practitioner and community midwife is no less safe than consultant care for women already selected to be at low risk,² and there is no evidence to support the claim that the safest policy is for all women to give birth in hospital.³