of proteins. Assays for total protein show poor performance in quality assurance schemes because of the diversity of methods and standards.² At all levels of protein excretion caused by glomerular disorders albumin is the main contributor. Albumin is a discrete protein, and specific methods for measuring it are now widely available.

Shihabi *et al* suggested that both urine total protein and albumin should be routinely estimated.³ In our experience, in a laboratory that receives more than 2500 requests for measurement of urine protein each year, there is a linear relation between urine total protein and albumin concentrations ($r=0.924$; $p<0.001$). We consider that urine albumin concentration should replace total protein concentration for the assessment of glomerular proteinuria.

FIONA C BALLANTYNE
DENIS ST J O'REILLY

Special Techniques Unit,
Institute of Biochemistry,
Royal Infirmary,
Glasgow G4 0SF

- Winocour PH. Microalbuminuria. *BMJ* 1992;304:1196-7. (9 May.)
- Chambers RE, Bullock DG, Whicher JT. External quality assessment of total urinary protein estimation in the United Kingdom. *Ann Clin Biochem* 1991;28:467-73.
- Shihabi ZK, Konecny JC, O'Connor ML. Albuminuria vs urinary total protein for detecting chronic renal disorders. *Clin Chem* 1991;37:621-4.

NCEPOD: more training needed

EDITOR,—S J Nixon highlights the fact that the national confidential enquiry into perioperative deaths for 1990 has to report the same deficiencies as the confidential enquiry into perioperative deaths for 1985-6.¹ In surgery for the complications of peptic ulcer, deaths in 120 patients are analysed; three quarters of the patients were elderly and seriously ill, but one third were managed by an unsupervised registrar. The risks for these patients were real, while the trainees were doubly handicapped—faced with undue responsibility and denied help from a senior with whom to discuss the choice and technique of treatment. The overall perioperative mortality is rather less than 1%, but there must be concern over some of the reported deaths and other aspects of current surgical training.

Firstly, reviews of registrars' experience provide further evidence of lack of supervision in theatre.^{2,3}

Secondly, a survey by the Association of Surgeons showed that training had suffered through bed closures in seven English regions. It has become difficult to learn the technique of hernia repair, with few patients being admitted and most day surgery being performed by consultants.⁴

Thirdly, theatres are now open for more limited periods and there is pressure to operate on more patients, so defter, trained surgeons do most of the work. Deprived trainees then operate on emergencies out of hours.

Fourthly, modern trends have resulted in an appreciable contraction of training opportunities in biliary and peptic ulcer surgery.

The royal surgical colleges share the vital task of inspecting and approving hospitals as suitable for training, but it is difficult to ascertain the true extent of the experience and supervision being given to trainees. The introduction of log books is a step in the right direction. The Royal Australasian College of Surgeons has taken the logical further step of requiring each hospital to submit a summary of each log book for central scrutiny, the outcome of which may be that the college requires corrective action to be taken if recognition for training is to continue.

Now is the time for the colleges to show unmistakably that service and training are distinct

activities. Service currently receives nearly all the attention. Training must be accorded specific status, with a planned programme prosecuted in protected time. Managements will require facts, not opinions, about training arrangements before they will entertain change: the training committees of the colleges will have the authority to require such changes only when they can present hospitals with accurate numerical information on the quality of the experience being offered.

PETER F JONES

Cults,
Aberdeen AB11 9HR

- Nixon SJ. NCEPOD: revisiting perioperative mortality. *BMJ* 1992;304:1128-9. (2 May.)
- Diggory RT. Training of surgical registrars. *BMJ* 1988;297:455-6.
- Steele RJC, Logie JRC, Munro A. Technical training in surgery: the trainees' view. *Br J Surg* 1989;76:1291-3.
- Association of Surgeons of Great Britain and Ireland. The effect of bed/theatre closures on junior staff training in general surgery in 1990. In: *British Journal of Surgery travelling fellow's report 1990*. London: Association of Surgeons of Great Britain and Ireland, 1991:6-9.

Improving preregistration training

EDITOR,—Improvements in preregistration training have to start with a reduction in hours devoted to service if disabling tiredness is not to destroy educational opportunity (and endanger quality of care).¹ Hours of work can effectively be shortened while service is maintained but only if there are more people to share the work²: hence my proposal to double up on preregistration house officers, which would solve the problem of hours for them without the need to train more doctors. Reducing "inappropriate tasks,"³ important though that is, cannot alone achieve this. Shift working might do it but inevitably disrupts continuity of care and educational programmes more.

Closer supervision is indeed needed at all levels of seniority^{4,5} and should start at once. Much that new house staff have to learn, however, is good technique in practical procedures, basic organisation of self and others, and simple decision making (which is not as simple in practice as in theory); for all of these the best tutor is often a slightly more senior fellow apprentice who has in turn been well supervised. To introduce an apprentice partner who would not only teach but learn by teaching basic practical competence, even if for only part of the working day, would be one (but only one) of several necessary steps forward in improving supervision. To denigrate such an apprentice partner as a "proxy consultant"⁶ is both to misunderstand the need and to overlook the fact that consultants are often not the most suitable people to meet it.

Well supervised training during limited hours in the front line of service needs to be complemented by withdrawal to an uninterrupted, reflective educational programme. There is much to be said for making this formal programme "common to all specialties" and for considering it to be "the first step on the road to a more integrated view of continuing education in medicine" as much as the last step in general medical education. The preregistration period should become the universal joint of medical education and training.

Certainly the preregistration period, lengthened or not, needs to be seen in the context of the preceding undergraduate course, shortened or not. There could not be a better moment. The General Medical Council is at war with "pot filling,"^{7,8} and most schools are now searching for a curriculum that will minimise information and maximise the ability to correlate and to solve problems.

Service hours must be shortened; supervision at all levels of skill and function must be strengthened; and an appropriate protected educational programme must be devised and implemented. It

will cost money (which any responsible employer would accept as a necessary investment in its expert workforce) and effort. But radical improvement will be achieved only if continuous constructive pressure comes from the trenches: if recent graduates rise above their own justified frustration, disappointment, and even anger and use their experience and vision to help build a better future for those who follow them.

Might the Council of Deans now take the initiative and convene a small representative national group to set out a range of options for radical reform?

PETER RICHARDS

St Mary's Hospital Medical School,
Imperial College,
London W2 1PG

- Richards P. Educational improvement of the preregistration period of general clinical training. *BMJ* 1992;304:625-7. (7 March.)
- Andrews J. Improving preregistration training. *BMJ* 1992;304:981. (11 April.)
- Rees G. Improving preregistration training. *BMJ* 1992;304:981. (11 April.)
- Mitchell TR, Sherwood T. Improving preregistration training. *BMJ* 1992;304:981. (11 April.)
- Bahrami J. Improving preregistration training. *BMJ* 1992;304:981. (11 April.)
- Dowling S. Improving preregistration training. *BMJ* 1992;304:980. (11 April.)
- Meakin RP. Improving preregistration training. *BMJ* 1992;304:980. (11 April.)
- General Medical Council. *Undergraduate medical education—consultation document sent to deans of medical schools May 1991*. London: GMC, 1991.

Draconian sentence imposed on Vietnamese doctor

EDITOR,—Dr Nguyen Dan Que graduated with distinction from Saigon University in 1966 at the age of 24. He studied in Brussels, London, and Paris from 1967 to 1974 and was made director of Cho-Ray Hospital in Ho Chi Minh City in 1975.

He specialised in thyroid disease and diabetes and wrote 13 papers from 1975 to 1978. Gradually, however, he became critical of the low standard of health care and treatment in his country and expressed his criticism openly. Arrested in February 1978 and accused of trying to overthrow the government, he was kept in solitary confinement for several months and not given enough to eat. For two months he was kept in shackles in a cell, less than 2×2 m, without sanitary facilities. He was not put on trial but was released in 1988 after 10 years' imprisonment.

After his release the police kept his diplomas, papers, albums, diaries, and address book, and he was not allowed to practise. Nevertheless, he wrote a paper about the treatment of 17 patients with insufficient insulin. After 1988 more insulin was sent to Vietnam from abroad.

Soon after his release Dr Nguyen helped to found a non-violent political movement, whose aims included "respect for the human, civil, and property rights of the people, . . . a pluralist political system, and free elections." He was arrested again in June 1990 and held incommunicado for 18 months. He was tortured soon after his arrest. He has a gastric ulcer, which has bled twice, and he has lost much weight. His family and the public were excluded from the court that sentenced him, and he was not allowed to speak in his own defence. On 29 November last year he was sentenced to 20 years' imprisonment followed by five years' house arrest. The BBC Far Eastern Service described this as "draconian . . . by any standard."

Dr Nguyen's health is deteriorating, and he is kept under a harsh regime. If he survives until the age of 75 he will then be free. The medical group of British Amnesty has appealed to the Vietnamese government for his release and return to medical practice; it has undertaken to provide him with

insulin regularly if this happens. I hope that doctors all over the world will help to persuade the Vietnamese government to free a heroic colleague before it is too late.

HAROLD HILLMAN

University of Surrey,
Guildford GU2 5XH

Keeping babies in prison

EDITOR,—Luisa Dillner's editorial on keeping babies in prison is a welcome reminder of one of the greatest inhumanities towards children in the United Kingdom today.¹ The Children Act emphasises the rights of children and the importance of promoting their upbringing within their families,² yet our judicial system seems to take the opposite view.

We are not surprised that infants brought up in prison did not show significant developmental delay on Griffiths developmental testing as this scale was not designed to measure the "harmful" effects of an inadequate social and emotional environment. An assessment of play and social interaction in these very young children would be much more informative.

The few children who are allowed to spend their early months in prison with their mothers are at least spared sudden enforced separation, often with fostering and always with totally inadequate visiting arrangements. Dillner did not mention the thousands of children whose fathers are in prison, many of whom are rendered in need by social and economic consequences. How does a child maintain a positive relationship with a parent in prison? Why do we continue to punish children in this way? More consideration should be given to non-custodial sentencing for non-violent offenders who do not pose a threat to children (as in the Nordic countries and increasingly in other European countries such as Italy) and to developing the provision of quality child care both in and out of prison.

A humane society is one that cares for its children. The British Association for Community Child Health is a subspecialty group of the British Paediatric Association; we believe that the decision to separate children from their parents is rarely justified and that the rights of the children should always be paramount.

SIMON LENTON
MARION CROUCHMAN

British Association for Community Child Health
(British Paediatric Association),
London NW1 4LB

1 Dillner L. Keeping babies in prison. *BMJ* 1992;304:932-3. (11 April.)

2 *Children act*. London: HMSO, 1991.

Predicting psychiatric admission rates

EDITOR,—Brian Jarman and colleagues report high correlations between psychiatric admission rates and indices of local socioeconomic deprivation.¹ We recently analysed data on the use of psychiatric services obtained from case registers in two areas in northern Italy, urban south Verona and rural Portogruaro, for 1983-9. In south Verona we, like Jarman and colleagues in their study, found high correlations between the all diagnoses psychiatric admission rates and the following census variables: living alone, unemployment, percentage of the total population who are dependants, and percentage who are divorced, separated, or widowed. These correlations were higher for schizophrenic patients than for all admissions (r^2 for best stepwise multiple regression model 0.94 and 0.79 respectively).

We suspect, however, that such a model may have important limitations. In rural Portogruaro, for example, there were no consistent associations between census variables and use of services. Furthermore, when we analysed only those admissions with a diagnosis of neurosis (ICD ninth revision) no significant correlations were found. Further research should perhaps look specifically at whether such correlations are found only in urban areas and only for psychotic, especially schizophrenic, patients.

GRAHAM THORNICROFT

PRISM (Psychiatric Research in Service Measurement),
Institute of Psychiatry,
London SE5 8AF

MICHELE TANSELLA

Servizio di Psicologia Medica,
Istituto di Psichiatria,
Verona,
Italy

1 Jarman B, Hirsch S, White P, Driscoll R. Predicting psychiatric admission rates. *BMJ* 1992;304:1146-51. (2 May.)

Decision analysis in medicine

EDITOR,—Although many doctors consider formal decision theory¹ to have little practical application, decision making is so central to the practice of medicine that all help is welcome. As part of an examination of a recent technical innovation (thrombolytic treatment for myocardial infarction) and the development of appropriate policy options we sought help from decision analysis. We constructed a decision tree, using a combination of national, international, and local data, to estimate the probabilities of the various outcomes. We applied a range of utilities reflecting possible preferences of patients, and the various options available were substantially clarified. Three points, however, arose from our studies, two prompting caution and one enthusiasm.

Firstly, the quality of the available data needed to complete these models is generally poor. We need much more fairly simple local information about the processes and outcomes of everyday activities such as, for example, what actually happens when a patient arrives at the front door of an accident and emergency department. One of us has tried, recently, to provide an example.² Secondly, gazing at decision trees has a bemusing effect: they can be taken to mean much more than their essentially crude weightings can tell us. Finally, doctors have talked a lot about the ways in which they can achieve true participation with patients in their health care. Here at last is a good place to start. Utilities can describe patients' hopes, fears, attitudes, understanding, and imperatives: doctors should begin to use them.

ALISON ROUND
ANDREW MARSHALL
MAURICE BACKETT

Plymouth Health Authority,
Plymouth PL6 5QZ

1 Thornton JG, Lilford RJ, Johnson N. Decision analysis in medicine. *BMJ* 1992;304:1099-103. (25 April.)

2 Hendra TJ, Marshall AM. Increased prescription of thrombolytic treatment to elderly patients with suspected acute myocardial infarction associated with audit. *BMJ* 1992;304:423-6. (15 February.)

EDITOR,—Two aspects of J G Thornton and colleagues' article on decision analysis in undergraduate medical training require comment.¹ The first is the decision analytical model adopted. Expected utility derives from econometric models of rational decision making in riskless situations using complete information. This does not reflect what happens in real life, when many complex medical decisions are made in risky situations with incomplete information. Subjective expected utility is an axiomatic utility analysis that can be

used in risky situations where complete information is not available.^{2,3} It has been used extensively to model human choice and decision making and is more appropriate in medical decision making.

My second comment concerns the acceptability of axiomatic decision making by patients. This cannot be justified by stating that "most people do agree that this is how they wish to make decisions." Rather, the test should be whether or not people make decisions that are consistent with the axioms of subjective expected utility. The research evidence is clear: they do not.³

Eraker and Politser reviewed the evidence on how decisions were reached between physicians and patients in the context of decision analysis.⁴ They concluded that decision analysis had the advantage of being "explicit, quantitative and prescriptive" but there was "a large body of empirically based research in behavioural decision theory [which] indicates that there are many potential biases and distortions that could affect any recommendation based exclusively on decision analysis."

Those working in clinical medicine can cite instances of the gulf that can exist between what people say they want and how they behave. It is salutary to recall such instances. Otherwise we may be becoming more "explicit, quantitative and prescriptive" on the basis of a false assumption: that people are rational about health care decisions.

PHIL C MACKIE

Directorate of Public Health Medicine,
Darlington Health Authority,
Memorial Hospital,
Darlington, DL3 6HX

1 Thornton JG, Lilford RJ, Johnson N. Decision analysis in medicine. *BMJ* 1992;304:1099-103. (25 April.)

2 Edwards W. The theory of decision making. *Psychol Bull* 1954;51:380-417.

3 Wright G. *Behavioural decision theory: an introduction*. Harmondsworth: Penguin, 1984.

4 Eraker SA, Politser P. How decisions are reached: physician and patient. *Ann Intern Med* 1982;97:262-8.

EDITOR,—The type of classic decision analysis described by J C Thornton and colleagues¹ has been an important development in the history of decision support systems. We believe, however, that there are compelling reasons why these systems have not been extensively used in a clinical setting.

Classic decision analysis may be very useful in resource allocation exercises on precisely defined problems, such as amniocentesis for prenatal diagnosis, where all outcomes are known at the beginning of the exercise. Probabilities can be determined accurately from published work and the value of each outcome can be estimated with some precision. Real clinical problems are seldom so simple and the variety of outcomes of the action is often not apparent at the start of the analysis. It is often necessary to introduce new options as the case develops and new information is acquired. There are major problems in calculating accurately what each outcome is worth to the patient; without this information the system cannot function. The authors allude to this problem in their paper.

The alternative approach is to use logical and qualitative techniques to represent and to reason with medical knowledge. These techniques offer a more flexible approach than simple numerical methods. Effective decision making is not dependent on the availability of precise numbers; achieving a proper understanding of the logical structure of the decision is more important.² The Oxford System of Medicine is a computer program which provides this type of more general reasoning and is also able to reason with numbers when appropriate. The system generates arguments for and against different decision options by logical inference from facts in a knowledge base. The arguments are considered and the logically correct decision options are presented. It is possible to examine the facts that were used to make the decision