a conserved breast, although the definition of local recurrence does depend on the extent and meticulousness of histopathological examination. Wide excision to ensure clear margins certainly compromises cosmetic results, particularly in small breasts, and the relative importance of cosmesis and local recurrence needs to be determined for each patient. Adequate radiotherapy minimises the risk of local recurrence, and if there is local recurrence survival does not seem to be affected adversely. The effect of a booster dose of radiation in relation to completeness of excision is currently the subject of a clinical trial by the European Organisation for Research and Treatment of Cancer.

R D RUBENS

Division of Oncology,
United Medical and Dental Schools of
Guy's and St Thomas's Hospitals,
London SE1 9RT

Babies' sleeping position

EDITOR,—Like Sarah Stewart-Brown, I am interested in the changes in infants' sleeping position over time.¹ I have looked at baby care manuals published in Britain since 1700 and have visited the Museum of Childhood in London. I have also looked at pottery babies in cradles in the Fitzwilliam Museum in Cambridge.

I have found no evidence of babies sleeping other than on their backs from the fifteenth century to early this century, when they began to be placed on their sides to avoid the stated danger of inhaling vomit. I have not yet found the evidence on which this advice was based. They remained on their sides at least until the late 1940s and early 1950s. The earliest reference to babies being put on their fronts to reduce the risk of inhaling vomit that I have found in a British book on baby care dates from 1961.²

The first British edition of Spock's *Baby and Child Care* (1955) states, "It's considered a little safer for a baby to sleep on his back in the first six months. . . ."³ I could not find why Spock thought that sleeping on the back was a little safer, but he thought that sleeping on the front was the alternative. He makes no mention of sleeping on the side. This could be the first suggestion in Britain that sleeping on the front was at least possible if not recommended. Stewart-Brown may be right that the doctors in special care baby units made a ghastly mistake, but the groundwork for such a shift in practice may have been made years earlier.

The idea may have crossed the Atlantic by other routes. *Recent Advances in Paediatrics* (1954) contains a chapter on sudden infant death by the Cambridge pathologist A M Barrett. He quoted an American paper from 1944 and suggested that the higher incidence of death in the prone position was probably due to the higher prevalence of that sleeping position in the United States.⁴

The stated advantages of babies sleeping on their fronts had all the benefits of being reasonable, logical, and easy to understand that make a health message acceptable. Whatever the origins of the advice, a terrible mistake was made. The recent return to the recommendation that babies should sleep on their backs may be the first decision since the professionalisation of baby care to be based on a reasonably scientifically established principle rather than mere assertion.

CHRISTINE HILEY

Department of Paediatrics,
Cambridge University School of Clinical Medicine,
Addenbrooke's Hospital,
Cambridge CB2 2QQ

1 Stewart-Brown S. Cot death and sleeping position. *BMJ* 1992; 304:1508. (6 June.)

2 Dally A. *The A-Z of babies*. London: Parrish, 1961.

3 Spock B. *Baby and child care*. London: Bodley Head, 1955.

4 Barrett AM. Sudden death in infancy. In: Gairdner D, ed. *Recent advances in paediatrics*. London: J and A Churchill, 1954: 301-20.

Percutaneous endoscopic gastrostomy feeding

EDITOR,—R H R Park and colleagues have highlighted important advantages of percutaneous endoscopic gastrostomy feeding over nasogastric tube feeding.¹ Neurogenic dysphagia is the predominant indication for inserting percutaneous endoscopic gastrostomy tubes,² and we have inserted 15 such tubes for neurogenic dysphagia out of a total of 22 tubes inserted over the past 18 months. We report the usefulness of percutaneous endoscopic gastrostomy in a particular subset of patients with neurogenic dysphagia: those who have difficulty in swallowing liquids but are able to swallow some solids.

A 62 year old man sustained three episodes of cerebral infarction in 1984, 1988, and 1990. He regained his mobility but had disabling pseudo-bulbar palsy; he was able to swallow solids slowly but aspirated liquids. Despite spending seven hours a day over his meals he ingested only about 2.9 MJ a day and lost 5 kg in weight. A nasogastric tube was inserted with difficulty, but he found it uncomfortable and a social handicap. It also blocked repeatedly, and he suffered great distress when it was reinserted. Assessment of swallowing by videofluoroscopy showed impaired oral and pharyngeal phases of swallowing. A percutaneous endoscopic gastrostomy tube was inserted in March 1991, and the patient gained 12 kg in weight over the next eight months. He could also enjoy some of the pleasures of normal eating as he could swallow some solids. His energy intake had to be reduced as he was becoming overweight, and he was able to go away on holiday, finding this method of feeding a great social success.

A 65 year old woman had a partially resolved right hemiparesis but virtually complete dysphagia after a brainstem stroke in December 1991. Her speech was intact, and videofluoroscopy showed persistent cricopharyngeal dysfunction. A fine bore nasogastric tube could not be passed despite repeated attempts and intervention by specialist ear, nose, and throat staff. A percutaneous endoscopic gastrostomy tube was inserted, and the patient was successfully rehabilitated at home; she considered the unobtrusiveness of the tube to be a great social benefit. After four months her swallowing started to improve: she was able to swallow some solids but remained unable to swallow liquid.

In patients with neurogenic dysphagia affecting their ability to swallow liquid the preferred method of giving commercial liquid feeds may be through a percutaneous endoscopic gastrostomy tube; the energy intake should be adjusted according to volitional intake of solids. Patients retain the ability to enjoy the taste of some solids, and in mobile patients this method of feeding offers considerable social and cosmetic benefits.

S GHOSH

M A EASTWOOD

Gastrointestinal Unit,
Western General Hospital,
Edinburgh EH4 2XU

1 Park RHR, Allison MC, Lang J, Spence E, Morris AJ, Danesh BJZ, et al. Randomised comparison of percutaneous endoscopic gastrostomy and nasogastric tube feeding in patients with persisting neurological dysphagia. *BMJ* 1992;304:1406-9. (30 May.)

2 Larson DE, Burton DD, Shroeder KW, DiMagno EP. Percutaneous endoscopic gastrostomy. Indications, success, complications and mortality in 314 consecutive patients. *Gastroenterology* 1987;93:48-52.

EDITOR,—R H R Park and colleagues report superior short term results in patients with persisting neurological dysphagia managed with percutaneous endoscopic gastrostomy compared with those managed with nasogastric tube feeding.¹ We agree that percutaneous endoscopic gastrostomy is the most appropriate management in such

patients. We have audited the long term outcome in 49 patients managed by percutaneous endoscopic gastrostomy, using an integrated team approach, in our unit over five years.

All patients undergoing percutaneous endoscopic gastrostomy in this hospital are managed by a multidisciplinary home enteral nutrition team, consisting of a gastroenterologist, nutritionist, dietitian, and nurse. We have instituted a seven day training programme for patients and families before discharge and a telephone hotline for patients to the team to help to deal with problems that arise and to provide advice.

Forty nine consecutive patients had percutaneous endoscopic gastrostomy. The main diagnoses were cerebrovascular accident (17 patients), motor neurone disease (13), malignancy (nine; mainly oropharyngeal), and others (10). The median duration of feeding was 175 days, and the total experience was 24 patient years. Thirty three patients were able to return home. Four patients died within 30 days, one from aspiration pneumonia two days after insertion of the gastrostomy tube. Of the 45 patients who survived more than 30 days, 22 experienced 41 complications. Twenty one of these complications were resolved without the need for a hospital visit, by a telephone discussion with the home enteral nutrition team and, in some cases, a home visit by the dietitian. The 41 complications observed were mechanical or blockage of the tube (21 cases), gastrointestinal symptoms (12), and local sepsis (eight), none of which was serious. Fifteen late deaths occurred, one due to gastric perforation related to the gastrostomy tube.

Our results show that the long term results of percutaneous endoscopic gastrostomy are excellent, that the long term rate of complications is lower than that previously described, and that most complications are minor and do not require a hospital visit. We suggest that patients are best cared for by an integrated team of nutritionists and gastroenterologists.

F E MURRAY J RAWLINGS

M A HULL C J HAWKEY

S P ALLISON

Departments of Nutrition and Therapeutics,
University Hospital,
Nottingham NG7 2UH

1 Park RHR, Allison MC, Lang J, Spence E, Morris AJ, Danesh BJZ, et al. Randomised comparison of percutaneous endoscopic gastrostomy and nasogastric tube feeding in patients with persisting neurological dysphagia. *BMJ* 1992;304:1406-9. (30 May.)

EDITOR,—We question the ethics of R H R Park and colleagues' study comparing fine bore nasogastric tube feeding with percutaneous endoscopic gastrostomy feeding and the issue of informed consent.¹ Despite Ian Forgacs and colleagues' reservations concerning overenthusiasm in adopting a technological advance,² percutaneous endoscopic gastrostomy feeding is well established, has few complications, and has been recommended for any patient in whom informed clinical judgment suggests that enteral tube feeding will be required for at least a month.³

To randomise patients at four weeks (when they are presumably still being fed by nasogastric tube) and to inform them that the "potential advantages of gastrostomy tube feeding were untested against the standard method of tube feeding" is not, in our opinion, appropriate. One of us recently reported personal experience confirming the long term safety and efficacy of percutaneous endoscopic gastrostomy feeding and suggesting that, in patients with neurological disorders of swallowing, recovery seems to be enhanced when a nasogastric tube is replaced by a percutaneous endoscopic gastrostomy tube⁴; the presence of a nasopharyngeal tube is unlikely to help recovery of speech and swallowing. The complications of nasogastric feed-