Teaching vaginal examination

EDITOR,—Having just spent a week examining medical students in the final MB, I am convinced of the need for an improved standard of training in all aspects of clinical examination, especially vaginal examination. As a teacher I could take no sense of pride in watching fumbling medical students desperately trying to impress on an examiner that they knew what they were doing when clearly they had rarely, if ever, inserted a vaginal speculum during their training.

If the operating theatre is not an appropriate place to learn vaginal examination then where is? Surely not the outpatient clinic, where throughput is so rapid that there is no time for the student and patient to develop rapport, or the ward, which is always bustling with people in these enlightened days of open visiting.

By taking such a negative attitude towards vaginal examination of anaesthetised patients and differentiating between the vaginal and other organs Susan Bewley is instilling into us the views of an articulate and vociferous minority who will, if allowed to continue, undermine the education of future generations of doctors.¹ It is hard enough to encourage medical students to come to the operating theatre without making them feel that if they examine patients they may be committing a criminal offence.

Some years ago, in response to demands from the local community health council, we introduced signed consent to clinical examination by medical students as part of the form giving consent to operation. Only a handful of women have ever refused to sign this part of the form. But surely by introducing different "rules" for medical students we are making the learning process more difficult and differentiating the medical student-patient relationship from the doctor-patient relationship.

Personally, I would prefer to see a new generation of well trained doctors who are able to relate appropriately to women who require gynaecological examination rather than a nation of women whose vaginas are protected from battery by medical students.

LINDA CARDOZO

King's College Hospital,
London SE5 9RS

1 Bewley S. The law, medical students, and assault. *BMJ* 1992;304:1551-3. (13 June.)

EDITOR,—After reading Susan Bewley's article I was left with one question: do medical students really need to learn how to perform a vaginal examination?¹ I think not. In 11 years of practising general internal medicine, both in hospital and in the community, I have yet to do a vaginal examination. I passed my postgraduate examinations without having to know about it. Even if I found or suspected something abnormal on vaginal examination a more experienced finger than mine would be needed to confirm and interpret the finding. I can thus spare my female patients (and myself) the additional embarrassing examination by referring them to the appropriate expert.

Vaginal examination could easily be taught to first time senior house officers in obstetrics and gynaecology (either general practice trainees or prospective gynaecologists) without anybody suffering. The trainee would then be a qualified

Advice to authors

Priority will be given to letters that are less than 400 words long and are typed with double spacing. All authors should sign the letter. Please enclose a stamped addressed envelope for acknowledgment.

doctor, a member of the team with a direct interest in the patient's welfare. Under such circumstances consent would not be an ethical or legal problem. All other doctors apart from those practising obstetrics and gynaecology need know only that vaginal examination, like psychoanalysis, is best left to those trained in it.

ANTHONY PAPAGIANNIS

Department of Respiratory Medicine,
Llandough Hospital,
Penarth,
South Glamorgan CF6 1XX

1 Bewley S. The law, medical students, and assault. *BMJ* 1992;304:1551-3. (13 June.)

EDITOR,—Susan Bewley's article opens a new debate on an old subject—namely, the need for respect for patients and how their consent is obtained by doctors and others who care for them.¹ The charge of battery for examination without consent has not been tested in court, and the risk of such a charge being raised is minimal. Bewley emotively uses vaginal examinations by medical students to illustrate the question of respect and consent. After 30 years of teaching medical students to perform vaginal examinations I believe that such teaching could and should be completely abandoned.

All trainee general practitioners and gynaecologists are taught the value and limitations of vaginal examination as postgraduates. Doctors working in other specialties do not perform vaginal examinations, so only those who need the skill should acquire it, and they should do so as doctors, not as medical students. Gynaecologists could then concentrate on teaching medical students about women's needs for medical care related to their reproductive organs, and the sensitivities and human values be illustrated by giving such care.

The present approach whereby all students are taught vaginal examination is unjustified as well as educationally harmful to the students. I am sure that if the teaching of vaginal examination to medical students was stopped for 12 months as an experiment it would never be considered necessary to restart it.

JAMES MCGARRY

Division of Obstetrics and Gynaecology,
Southern General Hospital,
Glasgow G51 4TF

1 Bewley S. The law, medical students, and assault. *BMJ* 1992;304:1551-3. (13 June.)

Management of breast cancer

EDITOR,—In responding to J Michael Dixon's review of the management of elderly patients with breast cancer¹ I H Kunkler argues for a palliative approach aimed at improving quality rather than duration of life, particularly for the large proportion

of women with locally advanced disease.² In Edinburgh Roberts *et al* showed that even during a screening trial more than a third of symptomatic women present with advanced disease.³ The proportion is highest in women living alone—and elderly people commonly live alone. It would, however, be too easy to assume that palliation alone is required. Of 100 elderly or frail patients (median age 76.3 years) treated in this centre with tamoxifen as primary treatment for non-metastatic breast cancer, 47 had T4 tumours and 14 had T3 tumours.⁴ Initially 68 patients responded, but 53 of these required subsequent treatment. Despite failure of local control overall in 42% of patients the five year survival was almost 60%.

In two reviews of radiotherapy in this department over the past 16 years we have found that the technique and dose are important. Langlands *et al* showed that low dose palliative techniques given in a few fractions for convenience were more commonly used in elderly people but produced poorer control of disease (table).⁵ More recently, Price *et al* showed that even high dose but palliative techniques produce poorer locoregional control of disease.⁶

Relation between radiation dose, mean age, and control of local disease at death

Radiation dose (Gy)	No of patients	Mean age (years)	Control of local disease at death (%)
30	23	72.3	19
40	10	65.4	
45	132	59.2	41

Clearly, elderly patients can survive many years after breast cancer is diagnosed. Treatment should be as radical as the stage requires and the patient can tolerate. Admittedly, radiation therapy can take several weeks to deliver. Arrangements for travel and so on should be such as to ensure minimal stress. The alternative of short palliative techniques will be poor control of disease, resulting in the increased stress precipitated by advancing local disease.

A RODGER

Radiation Oncology Unit,
Department of Clinical Oncology,
Western General Hospital,
Edinburgh EH4 2XU

- Dixon JM. Treatment of elderly patients with breast cancer. *BMJ* 1992;304:996-7. (18 April.)
- Kunkler IH. Treating elderly patients with breast cancer. *BMJ* 1992;304:1377. (23 May.)
- Roberts MM, Alexander FE, Elton RA, Rodger A. Breast cancer stage, social class and impact of screening. *Eur J Surg Oncol* 1990;16:18-21.
- Akhtar SS, Allan SG, Rodger A, Chetty U, Smyth JF, Leonard RCF. A ten year experience of tamoxifen as primary treatment of breast cancer in 100 elderly and frail patients. *Eur J Surg Oncol* 1991;17:30-5.
- Langlands AO, Kerr GR, Shaw S. The management of locally advanced breast cancer by x ray therapy. *Clin Oncol* 1976;2: 365-71.
- Price A, Kerr GR, Rodger A. Primary radiotherapy for T4 breast cancer. *Clin Oncol* (in press).

EDITOR,—Neither R D Rubens nor J Michael Dixon considered an important element of the management of breast cancer.^{1,2} A King's Fund consensus conference recognised the high rates of psychiatric morbidity in patients with primary breast cancer and recommended that a psychiatrist should be attached to each district team.³ A recent

paper by Jennifer Barraclough and colleagues underlines this; in their sample 45 out of 204 patients with breast cancer had a major depressive episode lasting at least three months after a surgical treatment.⁴ Similarly, Maguire *et al* showed that one in four women had significant psychological symptoms, including depression, anxiety, and sexual dysfunction, 12 months after mastectomy.⁵

Studies have concentrated on patients treated surgically, but all patients with breast cancer are at risk of psychological sequelae.⁶ Detecting these is important as psychological problems in patients with cancer respond well to several treatments.⁷

Though a third of all breast cancer occurs in those aged over 70, elderly people have been excluded from most studies assessing psychological morbidity. This is regrettable as breast cancer in elderly people tends to be treated more conservatively; operable breast cancer is less common, and, for many, quality of life is more important than quantity. Psychological health is therefore important. The studies that have not excluded elderly patients have not presented the results separately, so little is known about the psychological problems of elderly patients with breast cancer.

An unsupportive social network and debility are associated with a higher risk of psychological problems in studies of patients with cancer under 70.⁸ These factors may be more prominent in elderly patients. Moreover, depression is common in elderly people and, at least in the past, has often gone untreated.⁹

There may be considerable unrecognised and untreated psychological morbidity in elderly patients with breast cancer. Specific research is needed to clarify this. Awareness of this and detection and treatment of psychological distress by medical and nursing staff should form an important part of any management plan.

KWAME MCKENZIE

Maudsley Hospital,
London SE5 8AZ

SIMON LOVESTONE

Institute of Psychiatry,
London SE5 8AZ

- 1 Rubens RD. Management of early breast cancer. *BMJ* 1992;304:1361-4. (23 May.)
- 2 Dixon MJ. Treatment of elderly patients with breast cancer. *BMJ* 1992;304:996-7. (18 April.)
- 3 Consensus development conference: treatment of primary breast cancer. *BMJ* 1986;293:946-7.
- 4 Barraclough J, Pinder P, Cruddas M, Osmond C, Taylor I, Perry M. Life events and breast cancer prognosis. *BMJ* 1992;304:1078-81. (25 April.)
- 5 Maguire G, Lee E, Bevington D, Kuchemann C, Crabtree R, Cornell C. Psychiatric problems in the first year after mastectomy. *BMJ* 1978;i:963-5.
- 6 Fallowfield LJ, Baum M. Psychological welfare of patients with breast cancer. *J R Soc Med* 1989;82:4-5.
- 7 Lovestone L, Fahy T. Psychological factors in breast cancer. *BMJ* 1991;302:1219-20.
- 8 Dean C. Psychiatric morbidity following mastectomy: pre-operative predictors of type of illness. *J Psychosom Res* 1987;35:385-92.
- 9 MacDonald AJD. Do general practitioners "miss" depression in elderly patients? *BMJ* 1986;292:1365-7.

EDITOR,—R D Rubens has overestimated the absolute difference in 10 year mortality that can be produced by the widely tested forms of adjuvant treatment for high risk patients with early breast cancer but may have underestimated the proportion who gain some shorter term benefit.¹

As he says, adjuvant treatments such as ovarian ablation for premenopausal women, or at least a few years of tamoxifen for older women, reduce the annual death rate by about one quarter among both low risk and high risk women. But, although this will reduce 10 year mortality by about a quarter for low risk women (changing, for example, a 20% risk of death into about a 15% risk of death), it will reduce 10 year mortality by somewhat less than a quarter in high risk women. Among high risk women the absolute reduction in 10 year mortality is about 10 fewer deaths for every 100 women

Effects of persistent proportional reduction of 30% in annual mortality: eventual absolute benefit is about 12 per 100 (for example, 62 v 50 or 35 v 23)

treated, irrespective of whether there would have been 80, 60, or 40 deaths in the absence of treatment.

Turning from probabilities of death to probabilities of survival, 10 fewer deaths would mean that a 10 year survival of 20% is increased to about 30%, of 40% is increased to about 50%, and of 60% is increased to about 70%. But if the 10 year survival is increased from 40% and becomes 50% this does not necessarily mean that 10% are protected and 90% are not; it could be that in many of those who died during the first 10 years recurrence of death was delayed, or that many of those who have not yet died at 10 years but will do so later may likewise gain at least some delay. All we know for sure is the overall survival pattern in the absence and presence of treatment. This is illustrated in the figure for the particular case of a 30% reduction in annual mortality (which might well be achievable² and eventually produces an absolute difference of about 12 per 100 for high risk women). But there is at present no way of telling whether the figure shows a large benefit for a small proportion of women or a smaller benefit for a larger proportion.

RICHARD PETO

Clinical Trial Service Unit and
ICRF Cancer Studies Unit,
Radcliffe Infirmary,
Oxford OX2 6HE

- 1 Rubens RD. Management of early breast cancer. *BMJ* 1992;304:1361-4. (23 May.)
- 2 Early Breast Cancer Trialists' Collaborative Group. Systemic treatment of early breast cancer by hormonal, cytotoxic, or immune therapy: 133 randomised trials involving 31,000 recurrences and 24,000 deaths among 75,000 women. *Lancet* 1992;339:1-15,71-85.

EDITOR,—I have two comments to make about R D Rubens's article on the management of early breast cancer.¹

Firstly, in the discussion of mastectomy there is no mention of the use of immediate reconstruction of the breast. This is now routinely offered in many units. In the Edinburgh breast unit during 1990, 144 patients underwent mastectomy and 112 were offered immediate reconstruction, of whom 68 accepted. As Rubens points out, "the drawback with mastectomy is loss of the breast"; immediate reconstruction in some small part compensates for this by increasing patients' freedom of dress and self confidence. Although with increased use of primary systemic treatment mastectomy is likely to become less common, some women will still require it, and for these women immediate reconstruction should be available.

Secondly, in relation to breast conservation Rubens suggests that recent work at Guy's Hospital indicates that excision of the macroscopic tumour is all that is required if adequate doses of radiotherapy are given. Of 12 centres that have reported studies comparing the histology of excision margins and local recurrence after conservation treatment (excision and radiotherapy), five (Paris; Amsterdam; Leuven; Marseilles-Basle; Boston) have

shown significantly increased rates of recurrence of breast cancer if margins were involved.^{2,3} Four of the 12 centres (Nottingham, Los Angeles, Milan, Villejuif) have reported a non-significant increase in local failure rates⁴⁻⁶ despite higher doses of radiotherapy having been given in two studies if margins were involved. Two centres (Pennsylvania, Richmond) have given higher doses of radiotherapy to patients with carcinoma at the margins and reported similar local control rates in all groups irrespective of involvement of the margins,^{1,7} although between a third and a half of the patients in both centres underwent a re-excision and only patients with focal involvement of the margin proceeded with breast conservation.

This leaves one study in which histologically involved margins were neither related to recurrence nor used to select for radiotherapy dose.⁸ This study did, however, show that patients who had simple excision of the macroscopic tumour had more than twice the local recurrence rate of patients undergoing wide local excision. In contrast to the views of Rubens, these data suggest that the dual aims of local disease control and optimal cosmesis are most likely to be obtained by an excision that achieves clear histological margins by removing a rim of surrounding normal breast tissue followed by the minimum effective dose of radiotherapy.⁹

J MICHAEL DIXON

Department of Surgery,
Edinburgh University,
Royal Infirmary,
Edinburgh EH3 9YW

- 1 Rubens RD. Management of early breast cancer. *BMJ* 1992;304:1361-4. (23 May.)
- 2 Solin LJ, Fowble BL, Schultz DJ, Goodman RL. The significance of the pathology margins of the tumor excision on the outcome of patients treated with definitive irradiation for early stage breast cancer. *Int J Radiat Oncol Biol Phys* 1991;21:279-89.
- 3 Schnitt SJ, Connolly JL, Harris JR, Hellman S, Cohen RB. Pathologic predictors of early local recurrence in stage I and stage II breast cancer treated by primary radiation therapy. *Cancer* 1984;53:1049-57.
- 4 Van Limbergen E, van den Bogaert W, van der Schueren E, Rijnders A. Tumor excision and radiotherapy as primary treatment for breast cancer. Analysis of patient and treatment parameters and local control. *Radiother Oncol* 1987;8:1-9.
- 5 Locker AP, Ellis IO, Morgan DAL, Elston CW, Mitchell A, Blamey RW. Factors influencing local recurrence after excision and radiotherapy for primary breast cancer. *Br J Surg* 1989;76:890-4.
- 6 Veronesi U, Volterrani F, Luini A. Quadrantectomy versus lumpectomy for small size breast cancer. *Eur J Cancer* 1990;26:671-3.
- 7 Schmidt-Ullrich R, Wazer DE, Tercilla O. Tumor margin assessment as a guide to optimal conservation surgery and irradiation in early stage breast carcinoma. *Int J Radiat Oncol Biol Phys* 1989;17:733-8.
- 8 Ghossein NA, Alpert S, Barba AJ, Pressman P, Stacey P, Lorenz P, *et al*. Importance of adequate surgical excision prior to radiotherapy in the local control of breast cancer in patients treated conservatively. *Arch Surg* 1992;127:411-5.
- 9 Wazer DE, Di Petrillo T, Schmidt-Ullrich R, Safai H, Marchant DJ, Smith TJ, *et al*. Factors influencing cosmetic outcome and complication risk after conservative surgery and radiotherapy for early-stage breast carcinoma. *J Clin Oncol* 1992;10:356-63.

AUTHOR'S REPLY,—Richard Peto's comments are based on table II of my article. This table shows the relative efficacies, in different prognostic categories, of hypothetical treatments that absolutely reduce five year mortality by 25% or 50% respectively. The aim was to show how the effectiveness of treatment can vary in different groups of patients at an arbitrary time and so help in the selection of treatment for individual patients. The table was not meant to be a direct extrapolation from the reduction in the annual odds of death as identified from the recent meta-analysis. Peto's remarks are otherwise most helpful in showing the limits of interpretation permissible from current overview analyses.

J Michael Dixon's comment about the availability of immediate reconstruction for patients in whom mastectomy is unavoidable is worth noting. He also rightly emphasises the relation between unclear margins of excision and local recurrence in