Outcome of excision of natal cleft sinus and primary closure compared with that of excision and secondary healing. Figures are medians (ranges)

	Primary closure		Secondary healing (n=23)
	Success (n=17)	Failure (n=6)	
Days in hospital	1 (0)* (day case)	1 (1-2)*	3 (2-3)
Days to healing	14 (0)* (removal of sutures)	37.5 (26-48)†	41 (29-51)
Days off work	17 (15-21)*	38.5 (28-49)†	42 (28-56)
Outpatient visits	2 (0)*	5.5 (4-6)†	6 (4-7)
Visits by district nurse	0*	3 (2-5)†	4 (4-6)

*p<0.001, †NS compared with secondary healing group.

sinus. Wounds that failed to heal after primary closure were then allowed to heal by secondary intention, and the outcome of these patients' treatment was not different from that of the patients in the secondary healing group except that their stay in hospital was shorter (table).

We advocate day case excision and primary closure of uncomplicated natal cleft sinuses. The treatment results in less discomfort for patients, earlier return to work, and better use of health care resources in 74% of cases. The remaining 26% of patients will have a shorter stay in hospital and will otherwise not be different from patients given "standard" treatment of excision and healing by secondary intention.

H T KHAWAJA
P C WEAVER

Department of Surgery,
St Mary's Hospital,
Portsmouth

Health Economics Research Group,
Brunel University,
Uxbridge,
Middlesex

S BRYAN

- 1 Bissett IP, Isbister WH. The management of patients with pilonidal disease—a comparative study. *Aust NZ J Surg* 1987;57:939-42.
2 Allen-Mersh TG. Pilonidal sinus: finding the right track for treatment. *Br J Surg* 1990;77:123-32.

Home accidents in elderly people

EDITOR,—We agree with the conclusions put forward in both the paper¹ and the editorial² on home accidents in elderly people but believe that they should have gone further. As falls in this age group are often multifactorial and many go unreported¹ all medical professionals having contact with the older population have a duty to reduce potential risk factors. As with many of the new reforms of the health service, this will ultimately fall on the primary care team and general practitioners in particular.

Medical risk factors are already assessed as part of normal working practice. Home visits presently made by general practitioners could provide an opportunity for screening environmental risk factors. We believe that the yearly review of all patients aged 75 and over should also be performed in the home, rather than in the practice premises, as part of such a screening programme.

In our department occupational therapists making home visits use these opportunities to assess and alter potential environmental risk factors. This often entails removing loose mats, nailing down loose carpet edges, and adding hand-rails. One other simple yet effective alteration in view of the large proportion of falls associated with poor lighting¹ is to change the light bulbs used to a higher wattage—for example, to 100 W.

Should we advocate adding a hammer, some nails, and some 100 W light bulbs to the medical

bags of all general practitioners and hospital doctors making home visits? The relatively small expense might save only a few admissions to hospital, but when one day in a hospital bed costs at least £150 it may be a sound investment.

The increased workload resulting from screening the home environment of all patients aged 75 and over may be unacceptable to general practitioners. Could this screening therefore be performed by community occupational therapists or health visitors, who already have experience in such matters? This requires further research to determine practicality and cost effectiveness.

KEVIN SHOTLIFF
DEBESH MUKHERJEE
ABHIRAM PRASAD

Department of Geriatric Medicine,
St George's Hospital,
London SW17 0RE

- 1 Graham HJ, Firth J. Home accidents in older people: role of primary health care team. *BMJ* 1992;305:30-2. (4 July.)
2 Livesley B. Reducing home accidents in elderly people. *BMJ* 1992;305:2-3. (4 July.)

EDITOR,—Brian Livesley and Helen J Graham and Julia Firth make the point that more effort by the primary care services might reduce falls and fractures in elderly people living at home.^{1,2} An important theme in our recent randomised trial on preventing fractures by working with the whole community was that falls in elderly people may be a consequence of preventive measures.³ Attempting to create a safer environment for non-disabled people aged 70 and over seemed to increase the number of reported falls. Elderly people want independence, and to achieve this they often seem to demand of themselves a greater degree of mobility than they actually have. A limiting factor to what they will attempt is an unacceptable risk of falling. Thus reducing the risk of falling (either in reality or perceptually) seems to encourage an increase in mobility, either to the desired level or until the risk of falling becomes unacceptable.

It is obviously right to try to make the environment safer, but a relevant outcome for this policy is the achievement of independence and thus an improved quality of life rather than a reduction in falls. We do not attempt to overrestrict the activities of children because they might fall. We may have to rethink the extent to which we accept such risks for elderly people despite the grave consequences.

NORMAN J VETTER
PETER A LEWIS

University of Wales College of Medicine,
Cardiff CF4 4XN

- 1 Livesley B. Reducing home accidents in elderly people. *BMJ* 1992;305:2-3. (4 July.)
2 Graham HJ, Firth J. Home accidents in older people: role of primary health care team. *BMJ* 1992;305:30-2. (4 July.)
3 Vetter NJ, Lewis PA, Ford D. Can health visitors prevent fractures in elderly people? *BMJ* 1992;304:888-90. (4 April.)

Use of Lucozade and glucagon by ambulance staff in hypoglycaemia

EDITOR,—In 1987 the Department of Health and Social Security circulated a hazard warning notice to all health authorities regarding blood glucose tests done outside the laboratory.¹ We are surprised that J M Steel and colleagues did not mention this in their short report.² In the Merthyr and Cynon Valley Unit of Mid Glamorgan a formal policy was set up to meet the criteria suggested by the department. The policy covered training users, selecting and maintaining equipment, and quality control of meters and sticks. All staff who perform

blood glucose tests were trained, and all tests were done with Reflux S meters.

In 1989 Mid Glamorgan paramedical staff were included in the scheme. They use 50% glucose, glucagon, or oral carbohydrates. Quality control is even more important in the ambulance service in view of the effects of temperature and humidity on stored test sticks. These difficulties and the clinical conditions which can give misleading results are emphasised in training. Monthly quality control checks are carried out. For example, in one quality control exercise 162 ambulance staff were tested on their ability to diagnose hypoglycaemia on an unknown test sample. Altogether 159 staff obtained results within two standard deviations of the mean; the remaining three staff still correctly diagnosed hypoglycaemia.

All schemes must have adequate training and quality control.

K JONES

Prince Charles Hospital,
Merthyr Tydfil,
Mid Glamorgan CF47 9DT

H STEVENS

Mid Glamorgan Ambulance Service,
East Glamorgan Hospital,
Pontypridd,
Mid Glamorgan CF38 1AB

D MOODY-JONES

East Glamorgan General Hospital,
Pontypridd,
Mid Glamorgan CF38 1AB

- 1 Department of Health and Social Security. *Health services management. Blood glucose measurements: reliability of results produced in extra-laboratory areas.* London: DHSS, 1987. (HN JAM (87)13.)
2 Steel JM, Allwinkle J, Moffat R, Carrington DJ. Use of Lucozade as glucagon by ambulance staff for treating hypoglycaemia. *BMJ* 1992;304:1283-4. (16 May.)

Monitoring Creutzfeldt-Jakob disease

EDITOR,—Systematic surveillance of Creutzfeldt-Jakob disease in the wake of the epidemic of bovine spongiform encephalopathy, as reported by R G Will and colleagues, is to be welcomed.¹ We recently identified the disease in a 78 year old woman and were grateful to the National Creutzfeldt-Jakob Disease Surveillance Unit for help in postmortem investigation.

We have some concerns, however, about the identification of cases and collection of data. To our knowledge, ours is the only case so far reported by a psychiatric department. Neurology departments have recently been targeted for this surveillance exercise. Will *et al* pointed out, however, that diagnosis of the disease in elderly people has apparently decreased, which could be explained by poor ascertainment of cases.² Someone in his or her fifth or sixth decade who develops an organic brain syndrome of subacute onset is likely to be referred for neurological assessment whereas an older person is much less likely to be so intensively investigated and, if referred anywhere, is likely to be referred to the old age psychiatry service.

Creutzfeldt-Jakob disease in elderly people is rare compared with other dementias, but the importance of making the diagnosis cannot be underestimated. We urge our colleagues working in the psychiatry of old age to be aware of the surveillance project and to contact Will and colleagues if they strongly suspect the diagnosis.

DONALD LYONS
CAROLYN JOHNSTONE

Department of Old Age Psychiatry,
Leverdale Hospital,
Glasgow G53 7TU

- 1 Will RG, Ironside JW, Bell JE. Bovine spongiform encephalopathy and risk to health. *BMJ* 1992;305:53. (4 July.)
2 Will RG, Matthews WB, Smith PG, Hudson C. A retrospective study of Creutzfeldt-Jakob disease in England and Wales 1970-1979. II. Epidemiology. *J Neurol Neurosurg Psychiatry* 1986;49:749-55.