of people damaged as a result of taking medically prescribed drugs, there has not been one documented case of death due to nutrients and only a relatively small handful of cases reporting serious side effects. The cost to human health of excess salt consumption must be far greater.

The most reactionary aspect of the proposed legislation is that American and British journals are publishing a large and growing number of refereed articles reporting the value for human health of vitamin and mineral intake, not only in food form but also in nutrient form. Will the NHS of the future be happy to have nutrients given only through prescribed and costly "drugs"?

WALTER BARKER

Early Childhood Development Unit,
University of Bristol,
Bristol BS8 1HP

1 Richards T. Vitamins face control worldwide. *BMJ* 1992;305:491. (29 August.)

Oestrogen replacement after oophorectomy

EDITOR,—We are puzzled as to why C Harriet M Anderson and colleagues question whether oestradiol provided by transdermal patches will be as effective as that provided by implants in conferring long term benefits of oestrogen replacement therapy.¹ They raise this question solely on the basis of a small difference between the two routes of administration in suppressing gonadotrophin. Although this would suggest that the oestradiol dose delivered by the patches was lower than that delivered by the implant, there was actually no significant difference between the two treatments in concentrations of circulating oestradiol. It would therefore seem that delivered doses were actually similar.

The authors imply that a difference in gonadotrophin suppression will lead to a different effect on the short term and long term sequelae of the menopause. This extrapolation is based on a study which purported to show a dose-response effect (but actually used arbitrary lines) on urinary excretion of calcium and creatinine, a crude and indirect assessment of bone turnover.² It has already been shown that oestrogen sensitive tissues do not always respond in a dose dependent manner to increasing dosage; higher doses may reduce sensitivity to oestrogen, possibly because of down regulation of the receptor mechanism.³ In the study of Anderson *et al* the two methods of giving oestradiol had equivalent biological activity in terms of suppressing vasomotor symptoms. With regard to long term effects, the 0.05 mg transdermal dose of oestradiol is effective in preventing postmenopausal bone loss.⁴ It is also effective in producing beneficial changes in lipid and lipoproteins⁵ and in arterial pulsatility index,⁶ which suggest that it would be effective in reducing the risk of cardiovascular disease.

Thus there is no good evidence that the long term benefits of transdermal oestradiol 0.05 mg will be any less than those of oestradiol implants 50 mg. Both routes have the advantage of avoiding the hepatic first pass effect of steroids given orally, and the patches are easier to apply and discontinue.

JOHN C STEVENSON TIM C HILLARD
SOVRA I J WHITCROFT MALCOLM I WHITEHEAD

Wynn Institute for Metabolic Research,
London NW8 9SQ

- 1 Anderson CHM, Raju SK, Forsling ML, Wheeler MJ. Oestrogen replacement after oophorectomy: comparison of patches and implants. *BMJ* 1992;305:90-1. (11 July.)
- 2 Selby PL, Peacock M. Dose dependent response of symptoms, pituitary, and bone to transdermal oestrogen in postmenopausal women. *BMJ* 1986;293:1337-9.
- 3 Dyer GI, Young O, Townsend PT, Collins WP, Whitehead MI, Jelowitz J. Dose-related changes in vaginal cytology after topical conjugated equine oestrogens. *BMJ* 1982;284:789.
- 4 Stevenson JC, Cust MP, Ganger KF, Hillard TC, Lees B,

Whitehead MI. Effects of transdermal versus oral hormone replacement therapy on bone density in spine and proximal femur in postmenopausal women. *Lancet* 1990;335:265-9.

- 5 Crook D, Cust MP, Ganger KF, Worthington M, Hillard TC, Stevenson JC, *et al*. Comparison of transdermal and oral estrogen/progestin hormone replacement therapy: effects on serum lipids and lipoproteins. *Am J Obstet Gynecol* 1992;166:950-5.
- 6 Ganger KF, Vyas S, Whitehead M, Crook D, Meire H, Campbell S. Pulsatility index in internal carotid artery in relation to transdermal oestradiol and time since menopause. *Lancet* 1991;338:839-42.

EDITOR,—We agree with Harriet Anderson and colleagues that hormone concentrations are not routinely measured in patients receiving oestrogen replacement.¹

Although their study showed significantly lower gonadotrophin concentrations in the implant group, there was no statistically significant difference in the oestradiol levels in the two groups and both were equally effective in preventing hot flushes. As the beneficial effect of hormone replacement therapy correlates with oestradiol concentrations and not gonadotrophin concentrations, it should not cause any panic.

A serum oestradiol concentration of > 120 pmol/l provides no further additional effect and may be related to side effects like endometrial hyperplasia²; in the study of Anderson *et al* oestradiol concentrations were much higher in both groups. Oestrogen probably acts through intermediate products on bone metabolism; these could be prostaglandin E₂, calcitonin, or changes in ratio of insulin-like growth factor and growth hormone.³ Although calcitriol seems to be a safe and effective treatment for osteoporosis,⁴ oestrogen replacement therapy is of value in preventing further bone loss and fractures and may benefit the cardiovascular system as well.

As regards the cardiovascular system, while oral oestrogen results in a significant increase in plasma triglycerides and in all major lipoprotein factors, percutaneous administration exerts a weak effect on liver metabolism but substantially increases concentrations of high density lipoprotein cholesterol.⁵ Steingold *et al* confirmed that transdermal route may have less adverse hepatic effects.⁷ Chetkowski *et al* also showed efficacy of transdermal patches for biological effects.⁸

As implants and patches are equally good in maintaining oestradiol levels and suppressing menopausal symptoms and their beneficial effects on cardiovascular and bone metabolism, we see no reason in forcing implants on oophorectomised premenopausal or postmenopausal patients if they are keen to have patches.

B W ELIOT

J B SHARMA

L J MASCARENHAS

Acute Services Unit,
Northampton General Hospital,
Northampton NN1 5BD

- 1 Anderson CHM, Raju SK, Forsling ML, Wheeler MJ. Oestrogen replacement after oophorectomy: comparison of patches and implants. *BMJ* 1992;305:90-1. (11 July.)
- 2 Sarlet N, Reginster JY, Gaspard U. Determination of mineral serum oestradiol for prevention of postmenopausal bone loss. *Calcif Tissue Int* 1989;44(suppl):S69.
- 3 Duursma SA, Raymakers JA, Boereboom FTJ, Scheven BAA. Estrogen and bone metabolism. *Obstet Gynaecol Surv* 1992;47:38-44.
- 4 Tilyard MW, Spears GFS, Thomson J, Dovey S. Treatment of postmenopausal osteoporosis with calcitriol or calcium. *N Engl J Med* 1992;326:357-62.
- 5 Lufkin EG, Hodgson SF, Kotowicz MA, O'Fallon WM, Wnekner HW, Riggs BL, *et al*. The use of transdermal estrogen treatment in osteoporosis. In: Christiansen C, Overgaard K, eds. *Osteoporosis 1990*. Copenhagen: Osteopress, 1990:1995-8.
- 6 Tepper R, Goldberger S, May JY, Luz JJ, Beyth Y. Hormonal replacement therapy in postmenopausal women and cardiovascular disease: an overview. *Obstet Gynaecol Surv* 1992;47:426-31.
- 7 Steingold KA, Matt DW, Diezleiger D. Comparison of transdermal to oral estradiol administration on hormonal and hepatic parameters in women with premature ovarian failure. *J Clin Endocrinol Metab* 1991;73:275-81.
- 8 Chetkowski RJ, Meldrum DR, Steingold KA, Randle D, Lu, JK, Eggena P, *et al*. Biologic effects of transdermal estradiol. *N Engl J Med* 1986;314:1615-20.

EDITOR,—Harriet Anderson and colleagues state that premenopausal women undergoing prophylactic oophorectomy are given oestrogen replacement therapy to prevent menopausal symptoms.¹ It is unfortunate that they did not emphasise the real issues of long term skeletal and cardiovascular protection.² It is well recognised that premature ovarian failure appreciably increases the risk of death from myocardial causes.³ We question the ethics of delaying oestrogen replacement for seven days, thereby subjecting premenopausal women to miserable vasomotor symptoms. Was this really necessary?

In their final sentence Anderson and colleagues cast doubts as to whether transdermal therapy "will be as effective as the implant in conferring the many long-term benefits of oestrogen replacement." This is inappropriate for two reasons. Firstly, there are no long term epidemiological data regarding skeletal and cardiovascular protection for implants. Most of the published epidemiological data relate to oral conjugated oestrogens.^{4,5} Data do exist, however, showing comparable lipid changes and skeletal preservation of oral and transdermal therapy.^{6,7} Secondly, the dosages studied are not equivalent—a fact which is acknowledged in one of the studies quoted by Anderson and colleagues: Stanczyk *et al* stated that comparable oestradiol concentrations are obtained with 0.05 mg oestrogen patches and 25 mg implants.⁸

The authors state that the 0.05 mg patch can suppress gonadotrophin release after spontaneous menopause but not after oophorectomy. The papers cited do not confirm this: both studies found similar concentrations of follicle stimulating hormone and luteinising hormone after four weeks of oestrogen therapy.^{9,10} Indeed Chetkowski *et al*, referring to spontaneously postmenopausal women treated with various doses of oral and transdermal oestrogens, state: "even the largest doses of these medications did not lower the gonadotrophins to the levels observed in the premenopausal subjects." Anderson *et al* explain the recorded difference in gonadotrophin levels by "the higher sustained oestradiol concentrations in the implant group" yet their results state that the mean oestradiol concentrations did not differ significantly.

There is no evidence that concentrations of follicle stimulating hormone and luteinising hormone in postmenopausal women taking hormone replacement therapy are of any value in determining efficacy of treatment. It seems inappropriate on the basis of this study to cast doubt on the efficacy of transdermal administration of oestrogen. Evidence already exists indicating a similar biological effect to comparable doses of conjugated oral oestrogens, for which there is an abundance of epidemiological evidence.

BRIAN A REID

KEVIN F GANGER

Department of Obstetrics and Gynaecology,
St Mary's Hospital Medical School,
London W2 1PG

- 1 Anderson CHM, Raju SK, Forsling ML, Wheeler MJ. Oestrogen replacement after oophorectomy: comparison of patches and implants. *BMJ* 1992;305:90-1. (11 July.)
- 2 Stevenson JC. Osteoporosis and cardiovascular disease in women: converging paths? *Lancet* 1990;336:1121-2.
- 3 Colditz GA, Willett WC, Stampfer MJ, Rosner B, Speizer FE, Hennekens CH. Menopause and the risk of coronary heart disease in women. *N Engl J Med* 1987;316:1105-10.
- 4 Stampfer MJ, Colditz GA, Willett WC, Manson JE, Rosner B, Speizer FE, *et al*. Postmenopausal estrogen therapy and cardiovascular disease. Ten year follow up from the Nurses' Health Study. *N Engl J Med* 1991;325:756-62.
- 5 Paganini-Hill A, Ross RK, Gerkens VR, Henderson BE, Arthur M, Mack TM. Menopausal estrogen therapy and hip fractures. *Ann Intern Med* 1981;95:28-31.
- 6 Stevenson JC, Cust MP, Ganger KF, Hillard TC, Lees B, Whitehead MI. Effects of transdermal versus oral hormone replacement therapy on bone density in spine and proximal femur in postmenopausal women. *Lancet* 1990;336:265-9.
- 7 Crook D, Cust MP, Ganger KF, Worthington M, Hillard TC, Stevenson JC, *et al*. Comparison of transdermal and oral estrogen-progestin replacement therapy: effects on serum lipids and lipoproteins. *Am J Obstet Gynecol* 1992;166:950-5.
- 8 Stanczyk FZ, Shoupe D, Nunez V, Macias-Gonzales P. A randomised comparison of nonoral estradiol delivery in

- postmenopausal women. *Am J Obstet Gynecol* 1988;159:1540-6.
- 9 Chetkowski RJ, Meldrum DR, Steingold KA, Randle D, Lu JK, Eggena P, *et al.* Biologic effects of transdermal estradiol. *N Engl J Med* 1986;314:1615-2.
- 10 Kamel EM, Maurer SA, Hochler MG, Hoffman DI, Reber RW. Gonadotrophin dynamics in women receiving immediate or delayed transdermal estradiol after oophorectomy. *Obstet Gynecol* 1991;78:98-102.

AUTHORS' REPLY.—Our study showed significantly higher concentrations of gonadotrophin in oophorectomised women treated with the 0.05 mg oestradiol patch than in those given a 50 mg oestradiol implant. As stated in our paper, we think that the lower gonadotrophin concentrations in the implant group were due to the higher oestradiol concentrations observed. Unpublished data from this group of patients at 12 months show significantly different oestradiol concentrations: a mean (SE) in the patch group of 224 (43) pmol/l and in the implant group 544 (42) pmol/l (95% confidence interval for difference between means -443 to -197, $p < 0.0001$). As stated by Stevenson *et al.*, patches have been shown to prevent postmenopausal bone loss, as have implants,¹ but it has also been observed that percentage increase in bone density correlates with plasma concentrations of oestradiol,² and so treatments resulting in higher oestradiol concentrations may be more effective in preventing this long term consequence of the menopause.

This suggests that Eliot *et al* may not be correct in their assertion that serum oestradiol levels greater than 120 pmol/l provide no additional effect. We agree that implants should not be forced on postmenopausal women who are keen to have patches and vice versa.

Reid and Ganger question the ethics of our study. Delaying treatment ensured that all women had equivalent baseline hormone profiles before starting oestrogen replacement; given that one of the aims was to compare hormone profiles, this was a necessary part of the study design. They also question the doses used in this study: we did not set out to compare equivalent doses, but rather, the recommended starting doses. Careful reading of Chetkowski *et al*'s findings reveals that transdermal oestradiol significantly decreased gonadotrophin levels in a dose dependent manner.³ Kamel's paper referred to administration of the 0.2 mg oestradiol patch immediately after oophorectomy; suppression of gonadotrophins was not maintained.⁴ Our study showed that the 0.05 mg patch did not suppress gonadotrophin release after oophorectomy (mean concentration of follicle stimulating hormone after oophorectomy, 37.3 IU/l; after four months' treatment, 53.4 IU/l). Although there is no evidence from studies on the long term benefits of the 0.05 mg patch compared with the 50 mg implant, our study shows that there are differences in gonadotrophin concentrations which we suggest are due to differences in oestradiol concentrations and which may be reflected in the long term benefits of oestrogen replacement therapy.

C HARRIET M ANDERSON
K SHANTI RAJU

Department of Gynaecology,
St Thomas's Hospital,
London SE1 7EH

- 1 Savvas M, Studd JWW, Fogelman I, Dooley M, Montgomery J, Murby B. Skeletal effects of oral oestrogen compared with subcutaneous oestrogen and testosterone in postmenopausal women. *BMJ* 1988;297:331-3.
- 2 Studd JWW, Savvas M, Watson N, Garnett T, Fogelman I, Cooper D. The relationship between plasma oestradiol and the increase in bone density in postmenopausal women after treatment with subcutaneous hormone implants. *Am J Obstet Gynecol* 1991;163:1474-9.
- 3 Chetkowski RJ, Meldrum DR, Steingold KA, Randle D, Lu JK, Eggena P, *et al.* Biologic effects of transdermal estradiol. *N Engl J Med* 1986;314:1615-20.
- 4 Kamel EM, Maurer SA, Hochler MG, Hoffman DI, Reber RW. Gonadotrophin dynamics in women receiving immediate or delayed transdermal oestradiol after oophorectomy. *Obstet Gynecol* 1991;78:98-102.

Midwifery and body fluid contamination

EDITOR.—Josaphat J Kabukoba and Pam Young contend that current practices to prevent the skin of staff being contaminated by mothers' body fluids during delivery are inadequate and that staff may be at risk of contracting viral diseases while practising obstetrics in the United Kingdom.¹ This has enormous implications for medical and paramedical staff practising in developing countries, especially in sub-Saharan Africa, where one in 40 adult men and women are estimated to be infected with HIV.²

Recently, we have seen three nurses with HIV infection probably acquired occupationally while they were working in obstetric and accident and emergency departments in Africa. The first patient was a white missionary nurse who had never had sexual intercourse but had worked for many years in general nursing and midwifery. The second was an African nurse from Zambia whose only sexual contact was her husband, who was HIV negative. She had worked in an accident and emergency department, although she may have received a blood transfusion. The third patient was a white nurse who had worked for two years in Zambia but denied having had sexual intercourse during that time. She had also worked as a midwife but had had vaccinations and minor ophthalmic surgery while in Zambia.

Kabukoba and Young found that the incidence of broken skin among doctors and midwives examined immediately after they performed obstetric procedures was 23%. Furthermore, 34% of the assistants, of whom 35% did not wear gloves, were shown to be contaminated with body fluids immediately after vaginal deliveries. This has important implications for those working in developing countries, as highlighted by the above cases. Staff doing high risk work should be encouraged at the very least to wear gloves to prevent or minimise the risk of transmission of HIV and hepatitis B virus.

RUFUS FERNANDO
PATRICK TERRY
FRED WILLMOTT

Department of Genitourinary Medicine,
Royal South Hants Hospital,
Southampton SO9 4PE

- 1 Kabukoba JJ, Young P. Midwifery and body fluid contamination. *BMJ* 1992;305:226. (25 July.)
- 2 Gibb DM, Newell ML. HIV infection in children: epidemiological and diagnostic aspects. *International Journal of Sexually Transmitted Diseases and AIDS* 1992;3:235.

EDITOR.—A R Smyth and colleagues¹ may have misunderstood our paper on midwifery and body fluid contamination.² We did not discuss HIV infection across intact skin. This issue followed on after the discussion of the finding that 23% of staff had broken skin on their hands and arms. These skin breaks may be portals of entry for the virus, which may then be picked by Langerhans cells.

The studies quoted by Smyth *et al* report follow ups of single exposure incidents of health workers to infected body fluids.^{3,4} Obstetric staff have a cumulative risk during their long careers. With the rising prevalence of HIV it is not difficult to see that midwives and doctors face different risks from people having a single exposure. Any similarity to smoking and lung cancer?

We stated that the purpose of the study was to assess current methods of protection as issued by the royal colleges. We gave details of the methods of protection in use for the procedures mentioned and the levels of protection provided by these methods. Contamination even among surgeons doing caesarean sections who were "maximally" protected was 30%. Therefore the conclusion that the methods in use are inadequate is sound.

We will soon be publishing details of a new type of protection. A waterproof armsleeve with a watertight seal between it and the glove has been shown to reduce contamination to less than 5%. This is the answer to not only manual removal⁵ but also to all other procedures on the labour ward and theatre when spillage of blood is expected.

The risk of HIV infection may be small, but we do not know how small. Also it is constantly changing as prevalence changes. Authorities and staff must ensure that risks are kept to the minimum by adopting appropriate practices and methods, or we may, as suggested by Fernando *et al*,⁶ be too late.

JJ KABUKOBA

St George's Hospital Medical School,
London SW17 0RE

- 1 Smyth AR, Symth RL, Hart CA. Midwifery and body fluid contamination. *BMJ* 1992;305:474. (22 August.)
- 2 Kabukoba JJ, Young P. Midwifery and body fluid contamination. *BMJ* 1992;305:226. (25 July.)
- 3 Hart CA. Aids and the anaesthetist. In: Stoutenbeek CP, Van Saene HKF, eds. *Infection and the anaesthetist*. London: Bailliere Tindall, 1991:243-60. (Bailliere's clinical anaesthesiology.)
- 4 Centre for Disease Control. Human immunodeficiency virus infections in health care workers exposed to blood of infected patients. *MMWR* 1987;36:285-9.
- 5 Pearson MJ. Midwifery and body fluid contamination. *BMJ* 1992;305:474. (22 August.)
- 6 Fernando R, Terry P, Willmott F. Midwifery and body fluid contamination. *BMJ* 1992;305:713.

Day surgery for cataracts

EDITOR.—Hugh F Thomas and Roger Humphry confuse the issue of the advantages of day surgery when they fall into the trap of assuming the advantages of local anaesthesia over general anaesthesia.¹ With current anaesthetic agents it is possible to provide general anaesthesia while still offering early ambulation and feeding, with a low incidence of nausea and vomiting, and of coronary and embolic complications and less need for postoperative nursing care.

COLIN DRYDEN

Department of Anaesthesia,
Western Infirmary,
Glasgow G11 6NT

- 1 Thomas HF, Humphry R. Day surgery for cataracts. *BMJ* 1992;305:536-7. (5 September.)

Cardiorespiratory distress after sumatriptan given by injection

EDITOR.—Sumatriptan is a serotonin-1 (5HT-1) agonist that treats migraine by inducing cerebral vasospasm and is used to treat migraine and cluster headaches.^{1,2} A recent report suggested that coronary vasospasm may be induced in susceptible patients.³ We report two cases which show that serious ventricular arrhythmias may also be induced by this drug.

In the first case, a 42 year old woman who had suffered from migraine for many years and who had no previous cardiac history apart from mild hypertension (treated with Moduretic) collapsed within three minutes of receiving a first subcutaneous injection of sumatriptan. She was found to be in coarse ventricular fibrillation when the ambulance staff arrived seven minutes later. Sinus rhythm was regained by a single 200 KJ DC shock, and no further arrhythmias occurred. Subsequent investigations showed a normal 24 hour ambulatory electrocardiogram, an equivocal exercise stress test, and a 40% stenosis of the left anterior descending artery on cardiac catheterisation.

In the second case, a 67 year old man who was a life long migraine sufferer and who had had