ception are high among adolescents but that they lack the sense to assess personal risk.^{8,10}

These studies suggest that cognitive development plays an important part in adolescents' sexual behaviour. Hein mentioned four features of adolescent sexual behaviour: feelings of invulnerability, concrete versus abstract thinking, peer conformity, and denial.¹¹ Adventure and gratification (short term and immediate) could also be included in this list.

We need to teach adolescents skills and values in making decisions about their sexuality, including delaying sexual intercourse. These interpersonal behavioural skills are more likely to be acquired in a supportive social and economical environment and a society that is comfortable when talking about both the positive and negative consequences of sexual relationships.

JAYSHREE PILLAYE

Public Health Division,
Health Education Authority,
London WC1H 9TX

- 1 Smith R. Promoting sexual health. *BMJ* 1992;305:70-1. (11 July.)
- 2 Secretary of State for Health. *The health of the nation: a strategy for health in England*. London: HMSO, 1992. (Cm 1986.)
- 3 Office of Population Censuses and Surveys. *1985 Birth statistics*. London: HMSO, 1986. (Series FM1 No 12.)
- 4 Office of Population Censuses and Surveys. *1990 Birth statistics*. London: HMSO, 1992. (Series FM1 No 19.)
- 5 Catchpole MA. Sexually transmitted diseases in England and Wales: 1981-1990. *CDR Review* 1992;No 1.
- 6 Cancer Research Campaign. *Cancer of the cervix uteri*. London: CRC, 1990. (Factsheet 12.2.)
- 7 Health Education Authority. *Young adults' health and lifestyle: sexual behaviour*. London: Health Education Authority, 1990.
- 8 Balassone ML. Risk of contraceptive discontinuation among adolescents. *J Adol Health Care* 1989;10:527-33.
- 9 Rickert VI, Jay MS, Gottlieb A, Bridges C. Adolescents and AIDS. Females' attitudes and behaviours toward condom purchase and use. *J Adol Health Care* 1989;10:313-6.
- 10 Durant RH, Sanders JM. Sexual behaviour and contraceptive risk taking among sexually active adolescent females. *J Adol Health Care* 1989;10:1-9.
- 11 Hein K. AIDS in adolescence. *J Adol Health Care* 1989;10:305-55.

EDITOR,—In his editorial Richard Smith writes of a meeting of social engineers of indeterminate status at Leeds Castle.¹ This faceless group's purpose is to change people's attitudes in some unspecified way that is supposed to promote sexual health—a term used without definition. The intended propaganda offensive seems to be but an intensified and more explicit version of the sort of thing in place now, replete with a jauntier but equally meretricious alternative slogan to "safe sex." The wish is to encourage "a more erotic, positive, and diverse kind of sexual behaviour" as a contribution to the struggle against HIV.

In the front line as a general practitioner and a parent, I thought this approach wrong 30 years ago and feel sufficiently vindicated to prophesy that it will continue to be counterproductive: it has been accompanied by increasing promiscuity, abortions, venereal disease, and broken marriages.

Ominously, Smith says that those at this meeting not only condemned opposition to their views from moralists but, with breathtaking hubris, determined that it was their "duty not to bend to these people." No doubt there are those in the group who have the power to manipulate opinion in the media, including the *BMJ*, to suppress argument uncongenial to "amoralists" or "immoralists" (whatever the suitable word is for the opposite of people labelled moralist). I would hope that professional integrity would induce them to turn from such wickedness, but I am not the least bit confident of this now that they have proclaimed a higher duty.

JOHN FINDLATER

Carnforth,
Lancashire LA5 0TT

- 1 Smith R. Promoting sexual health. *BMJ* 1992;305:70-1. (11 July.)

EDITOR,—In discussing the best way to tackle HIV infection Richard Smith mentions that people should be able to exercise choice in sexual relationships.¹ To make an informed choice, however, people need to know certain facts about use of condoms. A discussion of attempts at developing a condom made of plastic stated, "While a more expensive material than latex, plastic is more uniform with fewer pinholes or variations in thickness."² In the same issue of *Population Reports* a recent analysis for the World Health Organisation of data from nine published studies was quoted. It found that people using condoms faced a risk about two thirds that of non-users of developing gonorrhoea, trichomoniasis, or chlamydia infection; when data on transmission of HIV from 10 studies were combined the analysis found a relative risk of 0.4 for people using condoms (M Rosenberg *et al*, strategies for viricide research meeting, World Health Organisation, March 1990).

A survey of people experienced in using condoms from a family planning clinic in Manchester reported that "52% of respondents had experienced condoms bursting or slipping off in the previous 3 months."³ The authors went on, "We were surprised to find how commonly our clients had experienced 'accidents,' even those who had apparently relied on condoms as their method of choice for contraception." The senior author of the above report also criticised the statement that "Most condom failure is due to incorrect usage" and added "We do the public no favours by promoting the idea among medical staff that it is all the fault of the user when things go wrong."⁴

JOHN KELLY

Birmingham B15 2PH

- 1 Smith R. Promoting sexual health. *BMJ* 1992;305:70-1. (11 July.)
- 2 New condoms in the 1990s. *Popul Rep [H]* 1990;8:28-31.
- 3 Kirkman RJE, Morris J, Webb AMC. User experience: Mates* v. Nuforms*. *British Journal of Family Planning* 1990;15:107-11.
- 4 Kirkman RJE. Condom use and failure. *Lancet* 1990;336:1009.

EDITOR,—Richard Smith's editorial on promoting sexual health addresses an important and sensitive public health issue.¹ His statement that "Roughly half of 16 year olds in Britain have had sexual intercourse" requires close examination, particularly because his article, rather than original data, is liable to be used by the mass media.

National statistics are not available, but recent surveys have shown that most teenagers are not sexually active before their 16th birthday.^{2,3} Although half of teenagers are probably not virgins by their 17th birthday, so that Smith's statement is technically correct, we believe that his statement may mislead and be interpreted, particularly by the media, as suggesting that the rate of sexual activity is higher than it actually is for teenagers below the age of consent. For example, a recent headline in the *Independent* read "Nearly half of young people lose virginity before 16,"⁴ even though the original article quoted did not give this information.⁵ An analysis of the given data (presuming that the age cohorts are equal) suggests a figure of roughly 34%—that is, a third rather than half, which is entirely consistent with other data.²

Society's view of what is normal has an important influence on individual behaviour.⁶ "Everyone's doing it" is a common myth among teenagers. Those attempting to reduce risk taking by teenagers in other aspects of their lives have found it useful to expose similar myths.⁶ If teenagers are led, by both the medical and the general press, to believe that most of their peers are sexually active this will increase pressure on them in their relationships still further. Those under 16 who are virgins may be labelled as a minority group, with an implication of abnormality.

Should we perhaps be more circumspect in our reporting of such figures? There is enough heat

within teenage relationships without the medical profession stoking the fire.

A R MELLANBY

F A PHELPS

J H TRIPP

Department of Child Health,
Postgraduate Medical School,
Exeter EX2 5SQ

- 1 Smith R. Promoting sexual health. *BMJ* 1992;305:70-1. (11 July.)
- 2 Curtis H, Lawrence C, Tripp J. Teenage sexuality: implications for controlling AIDS. *Arch Dis Child* 1989;64:1240-5.
- 3 Ford N. The sexual and contraceptive lifestyles of young people: part 1. *British Journal of Family Planning* 1992;18:52-5.
- 4 Hall C. Nearly half of young people lose virginity before 16. *Independent* 1992 July 21:2.
- 5 Baric L. Social expectations vs personal preferences—two ways of influencing health behaviour. *Journal of the Institute of Health Education* 1977;15:23-8.
- 6 Rundall TG, Bruvold WH. A meta-analysis of school-based smoking and alcohol use prevention programs. *Health Education Quarterly* 1988;15:317-34.

EDITOR,—Richard Smith highlights the problems encountered at a recent conference at Leeds Castle in finding a way of changing sexual behaviour to reduce the chances of HIV infection and unwanted teenage pregnancy.¹ The present emphasis is on the need to find a clear and readily understandable message. Given the gravity of the risks, would it be wrong to suggest that this should be "No sex outside marriage"? Such an approach might ease the burden on some teenagers, who may perceive themselves to be under pressure from their peers and the establishment to embark on sexual activities for which they are not really ready.

J W MYLES

Peterborough PE3 9PG

- 1 Smith R. Promoting sexual health. *BMJ* 1992;305:70-1. (11 July.)

EDITOR,—We endorse Richard Smith's view that the struggle against HIV needs reinvigorating and agree that the clarity of messages regarding health is important.¹ We are concerned that messages from some professional bodies are not only unclear but also damaging. For example, a statement from the Royal College of Surgeons of England earlier this year includes the recommendation: "If a patient is in a high risk category, he or she should be offered an HIV test, provided that facilities for counselling are available both before and after the test."² This is with a view to an operating surgeon deciding what extra precautions should be taken to minimise transmission of HIV during treatment or surgery.

No guidance is given regarding who should be included in such a category (it does not include known or suspected cases of HIV infection, for which there is a separate recommendation). Any assessment of high risk would require a detailed history, not just about injecting drug use and homosexual practice but also about heterosexual activity since the proportion of adults with AIDS reported to be heterosexual is 20%.³ We doubt whether such a history is part of the routine preoperative assessment undertaken by surgeons. In our view, the term "high risk category" is inappropriate and stigmatising and, in the context of control of infection, leads to false reassurance with regard to patients who are not perceived as being at high risk. Indeed, the recommendation is contrary to that given by the World Health Organisation, which advocates universal precautions regarding blood and body fluid (that is, that all blood and certain body fluids are assumed to be infectious for hepatitis B virus and HIV and other bloodborne pathogens) as the cornerstone of preventing transmission of bloodborne infections.⁴

Terminology and messages are important: the continued use of the term high risk category suggests that there is still a widely held belief that HIV infection occurs only in specific well defined