Royal College of General Practitioners, they are in a better position than us to validate or reject such a conjecture. Our information is that no such effect is evident in the figures. As for the third hypothesis, which they implicitly attribute to us, our version was that "open cast coal mining increases episodes of asthma." To distinguish between open cast coal mining and coal dust is crucial for further work.

Finally, though it would be foolish to ignore Tudor Hart's authoritative paper (and others he quotes) on the underdiagnosis of asthma, we must reiterate that our study concerned episodes and not prevalence. His comments are therefore inappropriate.

J M F TEMPLE

Glynneath,
West Glamorgan SA11 5AL

A M SYKES

University College,
Swansea

1 Correspondence. Asthma and open cast mining. *BMJ* 1992;305:644-5. (12 September.)

Asthma deaths in New Zealand

EDITOR,—Julian Crane and colleagues present time trend data correlating the rise and fall of fenoterol's market share in New Zealand between 1976 and 1989 with a coinciding epidemic of deaths due to asthma.¹ Three considerations are valuable in judging the merit of evidence from this form of data.

Firstly, the authors' argument of an effect from fenoterol would have been more convincing had the death rate followed the rise, plateau, and fall of fenoterol's market share. Of the three phases, however, only the rise during 1976-9 fits such reasoning. Moreover, that such a steep rise can be ascribed only to fenoterol is debatable. Applying the relative risk of 2.0 found in New Zealand for fenoterol during 1977-81,² we find that the "epidemic" should have increased the pre-1976 rate of 1.5/100 000 to no more than 2.0/100 000, lower than the peak of 4.1/100 000 observed in 1979.

Secondly, other equally plausible explanations emerge from time trend analyses. For example, the initial rise could have coincided with the onset of overuse of all inhaled β agonists, as a class. This would be consistent with the findings of our

Observed rate (■—■) of asthma death per 100 000 people aged 5-34 in New Zealand, 1974-89. Predicted rate (---) 1977-89, from linear Poisson regression, along with 95% prediction limits

Saskatchewan study of at least a tenfold increase in the risk of death from asthma in people using yearly more than one metered dose inhaler of any β agonist per month,³ not only fenoterol. For the decrease in the death rate from asthma after 1977 an alternative explanation could be the more recent worldwide trend favouring the increasing use of inhaled steroids.

Lastly, Crane and colleagues submit that the drop in the death rate from asthma from 2.2/100 000 in the first half of 1989 to 1.1/100 000 in the second half is significant. Instead, the appropriate scientific question should be whether the second rate for 1989 is significantly different from the rate expected taking the pre-existing 12 year decline and the yearly fluctuations into account. We fitted a linear Poisson regression model⁴ for the period 1977-89 and obtained 95% prediction bands (figure). It is clear from this analysis that the rate of 1.1/100 000 from the second half of 1989 falls within the range expected solely from statistical variability. Accordingly, the "drop" in the second half of 1989 is not significant, contrary to the assertion of the authors.

In summary, although the fenoterol hypothesis advanced by Crane and colleagues is possible, inconsistencies in the logic, the existence of plausible alternative explanations, and inaccurate statistical analyses render it unlikely. The pitfalls inherent in time trend data analyses should be considered when assessing evidence on the risk and safety of this and other drugs.

SAMY SUISSA

PIERRE ERNST

WALTER O SPITZER

Department of Epidemiology and Biostatistics,
McGill University,
Department of Medicine, Montreal General Hospital,
Montreal, Quebec, Canada H3A 1A2

- 1 Crane J, Pearce N, Burgess C, Beasley R, Jackson R. Asthma deaths in New Zealand. *BMJ* 1992;304:1307. (16 May.)
- 2 Pearce N, Grainger J, Atkinson M, Crane J, Burgess C, Culling C, et al. Case-control study of prescribed fenoterol and death from asthma in New Zealand: 1977-1981. *Thorax* 1990;45:170-5.
- 3 Spitzer WO, Suissa S, Ernst P, Horwitz R, Habbick B, Cockcroft D, et al. The use of β -agonists and the risk of death and near-death from asthma. *N Engl J Med* 1992;326:501-6.
- 4 McCullagh P, Nelder JA. *Generalized linear models*. 2nd ed. London: Chapman and Hall, 1989.

EDITOR,—Julian Crane and colleagues misrepresent the recent trends in mortality from asthma to defend their hypothesis that restriction on the use of fenoterol in New Zealand in 1989 led to a reduction in this mortality.¹ The recent figures fall well within the 95% confidence intervals predicted by trends in mortality before 1989 in both the 5-34 and 35-55 year old populations (figure).

The research group in Wellington has always held that there was no substantive evidence that fenoterol was prescribed to patients with more severe asthma.² "Channelling" occurs when patients who have either suffered adverse effects or not benefited from established treatments are prescribed a new drug. Petri *et al* have described channelling of inhaled β agonists to patients differing in the severity of their asthma in the Netherlands.³ They found that 2.0-2.5 times more treatment with systemic steroids and categories of other drugs for asthma was prescribed concomitantly to patients receiving fenoterol compared with those receiving salbutamol (with terbutaline intermediate).⁴ We have made a similar observation in New Zealand.⁵

The research group in Wellington has argued that the association between fenoterol and death due to asthma was not due to confounding by severity since the estimated relative risk for fenoterol did not decrease when the group examined the subgroups of patients with the most severe disease. This argument is inconsistent with data from the group's third study.⁶ Patients with

Mortality from asthma in New Zealand, 1980-90 (broken lines indicate 95% confidence intervals)

the highest baseline risk of death are those with a recent hospital admission and prescriptions for at least three classes of drugs for asthma when patients using fenoterol are excluded (relative risk=4.3). Among these high risk patients the overall relative risk for fenoterol decreases from 2.66 to 2.21. Therefore, confounding by severity of asthma cannot be excluded as a reason for the association between fenoterol and mortality from asthma. If baseline severity of asthma did increase as a result of continuous use of inhaled β agonists or use of high doses of these drugs⁶ then indices of chronic severity would be intermediate in the causal pathway and would be expected to reduce the relative risk. Alternatively, the severity of asthma increased in New Zealand as a result of other factors and newer medicines for asthma, including fenoterol, were prescribed to patients with more severe asthma—that is, channelling occurred.

Crane and colleagues' letter was given wide media coverage in New Zealand before there was any opportunity of peer review. As I have shown, changes in mortality have actually been modest since use of fenoterol was restricted in New Zealand.

JEFF GARRETT

Department of Respiratory Medicine,
Green Lane Hospital,
Auckland,
New Zealand

- 1 Crane J, Burgess C, Pearce N, Beasley R. Asthma deaths in New Zealand. *BMJ* 1992;304:1307. (16 May.)
- 2 Pearce NE, Beasley R, Burgess C, Grainger J, Crane J. Confounding by severity does not explain the association between fenoterol and asthma death. *Aust NZ J Med* 1990;20(suppl 1):515.
- 3 Petri H, Urquhart J, Herings R, Bakker A. Characteristics of patients prescribed three different inhalational beta-2 agonists: an example of the channelling phenomenon. *Post-Marketing Surveillance* 1991;5:57-66.
- 4 Garrett J, Turner P. The severity of asthma in relation to beta agonist prescribing. *NZ Med J* 1991;104:39-40.
- 5 Grainger J, Woodman K, Pearce N, Crane J, Burgess C, Keane A, et al. Prescribed fenoterol and death from asthma in New Zealand, 1981-7: a further case-control study. *Thorax* 1991;46:105-11.
- 6 Sears MR, Taylor DR, Print CG, Lake DC, Li Q, Flannery EM, et al. Regular inhaled beta agonist treatment in bronchial asthma. *Lancet* 1990;336:1391-6.