higher in the treated group²; for these two reasons the use of this particular drug has waned. The two other trials each showed a significant benefit in reduction of coronary risk with two very different drugs.^{3,4} This is strong evidence that use of carefully selected drugs to lower cholesterol concentration has an important clinical benefit. I do not believe that the meta-analysis alters this view.

JOHN ANDERSON

Medical Professorial Unit,
St Bartholomew's Hospital,
London EC1A 7BE

- 1 Ravnskov U. Cholesterol lowering trials in coronary heart disease: frequency of citation and outcome. *BMJ* 1992;305:15-9. (4 July.)
- 2 Committee of Principal Investigators. A cooperative trial in the primary prevention of ischaemic heart disease using clofibrate. *Br Heart J* 1978;40:1069-103.
- 3 Lipid Research Clinics Program. The Lipid Research Clinics coronary primary prevention trial results. I. Reduction in incidence of coronary heart disease. *JAMA* 1984;251:351-64.
- 4 Frick MH, Elo O, Haapa K, Heinonen OP, Heinsalmi P, Helo P, et al. Helsinki heart study: primary prevention trial with gemfibrozil in middle-aged men with dyslipidemia. *N Engl J Med* 1987;317:1237-45.

EDITOR,—U Ravnskov's paper is an impressive demonstration of the citation bias that favours references to supportive rather than unsupportive trials.¹ Ravnskov unintentionally illustrates this thesis by not mentioning the cholesterol lowering arm of our diet and reinfarction trial, the results of which were unsupportive or inconclusive, depending on the expectations, given a net cholesterol reduction of 3.6%.²

Similar analyses of trials in other topics of research would be illuminating. Publication bias is well recognised; citation bias must also be considered in the evaluation of medical evidence of all kinds.

M L BURR
A M FEHILY
P M SWEETNAM
P C ELWOOD

MRC Epidemiology Unit (South Wales),
Llandough Hospital,
Penarth,
South Glamorgan CF6 1XX

- 1 Ravnskov U. Cholesterol lowering trials in coronary heart disease: frequency of citation and outcome. *BMJ* 1992;305:15-9. (4 July.)
- 2 Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PL, et al. Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989;ii:757-61.

EDITOR,—In most human endeavours winners get cited more often than losers because people are most interested in winners. Yet this citation bias does not negate the fact that winners usually win because they are better. So also with prevention trials: the successful ones get cited more often, but only if they are perceived to be better. For example, in U Ravnskov's table I, strikingly positive trials such as that of Dorr *et al* and the Stockholm secondary prevention study (odds ratio for fatal coronary heart disease 0.40 (men) and 0.56 respectively) each accumulated only eight citations a year, presumably because their designs were flawed.¹ In contrast, the less positive but larger and better designed Lipid Research Clinics trial and Helsinki heart study (odds ratio for fatal coronary heart disease 0.78 and 0.73 respectively) collected 153 and 164 citations a year respectively.

Ravnskov should realise that preferential citation reflects the perceived quality of the evidence rather than the outcome.

GILBERT R THOMPSON

Medical Research Council Lipoprotein Team,
Hammersmith Hospital,
London W12 0HS

- 1 Ravnskov U. Cholesterol lowering trials in coronary heart disease: frequency of citation and outcome. *BMJ* 1992;305:15-9. (4 July.)

AUTHOR'S REPLY,—P N Durrington and colleagues still think that lowering cholesterol is useful as my unsupportive results are due to bias. According to them, I should have excluded multifactorial trials, non-randomised trials, and two trials that allegedly did not use the intention to treat principle. They have obviously not made any calculations because if these trials had been excluded the overall results would have been even more unfavourable. They find support in Holme's meta-analysis, although his review included four of the trials that I should have excluded; and also in a seven year old abstract by Peto *et al*. Let us discuss the matter again when Peto *et al* have published their results in detail.

Trials of oestrogens and thyroxine should also have been excluded; these drugs "do not simply reduce cholesterol," as F L Game and R H Neary put it. No drug simply reduces cholesterol concentration; most of them have several diverse side effects, but trial directors make things easy for themselves if they claim their drug to be cardioprotective when the number of coronary events is lowered but cardiotoxic when the number is increased.

J F Goodwin, and also Game and Neary, think that I have biased my results by excluding the trials of regression of atheroma. These trials, however, used angiographic changes as the primary end point. They did record cardiac morbidity and mortality also, but, owing to the few events that occurred, their inclusion should not have changed the overall results. Furthermore, regression of atheroma was not proved by these studies. What they recorded is a trivial, generalised widening of the coronary vessels and this is also what is seen in the rare published angiographs.¹ But atheromatosis is a localised disease, and its rate of progression is independent of the cholesterol concentration; this has been shown by at least six angiographic studies that have never been cited.²

Game and Neary suggest that the expected correlation between outcome and degree of cholesterol lowering was neutralised by a greater lowering in those at greater risk. Their idea cannot be tested from published data, but it seems unlikely because a high serum cholesterol concentration is not a risk factor after one or more myocardial infarctions,³ and nine of the 13 trials that had analysed the relation were secondary preventive trials.

I excluded the unsupportive trial of Burr *et al* because some of the controls were advised to eat more fish.⁴ But it must be unintentional that my critics, by citing mainly favourable reports, have illustrated the point of my paper.

UFFE RAVNSKOV

Råbygatan 2,
S-223 61 Lund,
Sweden

- 1 Brown G, Albers JJ, Fisher LD, Schaeser SM, Lin J-T, Kaplan C, et al. Regression of coronary artery disease as a result of intensive lipid-lowering therapy in men with high levels of apolipoprotein B. *N Engl J Med* 1990;323:1289-98.
- 2 Ravnskov U. An elevated serum cholesterol level is secondary, not causal, in coronary heart disease. *Med Hypotheses* 1991;36:238-41.
- 3 Shanoff HM, Little JA, Csima A. Studies of male survivors of myocardial infarction. XII. Relation of serum lipids and lipoproteins to survival over a 10-year period. *Can Med Assoc J* 1970;103:927-31.
- 4 Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PL, et al. Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989;ii:757-61.

Adenosine and cardiac arrhythmias

EDITOR,—Having seen the common reluctance of doctors to diagnose ventricular tachycardia, we agree with Clifford J Garratt and colleagues that "both accident and emergency departments and cardiac care units would be safer places for patients

with arrhythmias if the first instinct of the doctor on the spot was to reach for adenosine rather than verapamil."¹ We wish, however, to sound a note of caution.

Torsades de pointes developed in a 24 year old woman who had narrow complex tachycardia of 160 beats/min with a prolonged QT interval when 6 mg adenosine was given intravenously to establish the diagnosis.² This led the authors to suggest that adenosine should not be used in the presence of the prolonged QT syndrome. Another woman with a narrow complex tachycardia and a systolic blood pressure of 62 mmHg became briefly unresponsive with fixed dilated pupils after being given 12 mg adenosine; sinus rhythm resumed after some minutes.² A woman with supraventricular tachycardia of 176 beats/min went into asystole for 17 seconds and became unconscious, but the rhythm ultimately reverted to normal sinus rhythm.³

As with many other drugs initially declared safe, problems like those described above may occur; more importantly, adenosine may be used inappropriately, as in the woman with a systolic blood pressure of 62 mmHg, in whom cardioversion should have been used.

S S TACHAKRA

S ROBINSON

Central Middlesex Hospital NHS Trust,
London NW10 7NS

- 1 Garratt CJ, Malcolm AD, Camm AJ. Adenosine and cardiac arrhythmias. *BMJ* 1992;305:3-4. (4 July.)
- 2 Wesley RC, Turnquest P. Torsades de pointes after intravenous adenosine in the presence of QT syndrome. *Am Heart J* 1992;123:794-6.
- 3 Reed R, Falk JL, O'Brien J. Untoward reaction to adenosine for supraventricular tachycardia. *Am J Emerg Med* 1991;9:566-70.

Cardiopulmonary resuscitation in British hospitals

EDITOR,—Hugh Tunstall-Pedoe and colleagues' survey of cardiopulmonary resuscitation in British hospitals provides a baseline against which other results can be measured.¹ We have been auditing the outcome of cardiopulmonary resuscitation at the City Hospital, Nottingham, since September 1991 with the aim of determining whether the in house resuscitation training programme is adequate and well directed. The hospital had 1208 admissions to the coronary care unit during 1991.

Data on cardiac arrests were recorded on an audit form by the team leader. Completion of data collection was measured by cross checking with "arrest calls" put out from the hospital switchboard and showed that over 95% of calls were audited. Our results to April this year are in good agreement with those of Tunstall-Pedoe and colleagues' study. One hundred and forty three forms were collected in seven months (about 245 a year). Sixty (43%) out of 141 patients (two arrests were repeat arrests and excluded) survived resuscitation, of whom 42 (70%) were alive at 24 hours. At a median follow up of six weeks 29 patients (69% of survivors at 24 hours) were still alive. Survival data for one year are not yet available. Similarly, our figures for the type of arrest (cardiac 49% (70/143), respiratory 17% (25/143), and mixed cardiac and respiratory 24% (35/143)) and immediate survival related to age (age <50, 68% (13/19); 50-70, 60% (24/40); and >70, 38% (25/65); data on age incomplete) agree with those of Tunstall-Pedoe and colleagues' survey.

The authors comment that the paper is an initial publication and will be followed by subgroup analyses including analysis of specific rhythms. Unfortunately, this means that they have not presented data on the outcomes related to the initial cardiac rhythm observed by the team leader, which we have found is of major importance. We observed that immediate survival was 71%

(20/28) after an arrest in which the initial rhythm was ventricular fibrillation but only 20% (6/30) after an asystolic arrest and 8% (2/23) after electromechanical dissociation. These differences probably reflect not only the reversibility of the disturbance of the cardiac rhythm but also the underlying disease processes leading to the arrest. No patient who initially survived an asystolic arrest or electromechanical dissociation either survived to six weeks or left hospital (though numbers are small).

The information presented by Tunstall-Pedoe and colleagues serves as an excellent benchmark by which others can measure their success or failure rates. The addition of information on the initial rhythm will surely, as our data suggest, show great differences in survival and will need to be included before the data can truly act as a standard.

Our audit has already changed practice. Funds are being found for a resuscitation officer, and certain areas of the hospital are improving the telephone links to the switchboard to reduce the delay in calling the arrest team. The resuscitation working group has made changes to equipment to facilitate rapid resuscitation, and further changes have been proposed. The poor outcome after asystolic arrests and electromechanical dissociation has led to changes in emphasis in training in resuscitation, and the good outcome with ventricular fibrillation encourages good training and practice.

In the light of the Tunstall-Pedoe and colleagues' and our results there seems little doubt that all hospitals should have a measure of quality control for cardiopulmonary resuscitation and that feedback from this will have major effects on local management and training. Both studies give cause for optimism as immediate survival is higher than anecdote suggests and a considerable number of patients have prolonged survival.

ALISTAIR MCINTYRE

Department of Medicine,
Division of Therapeutics,
University Hospital,
Queen's Medical Centre,
Nottingham NG7 2UH

1 Tunstall-Pedoe H, Bailey L, Chamberlain DA, Marsden AK, Ward ME, Zideman DA. Survey of 3765 cardiopulmonary resuscitations in British hospitals (the BRESUS study): methods and overall results. *BMJ* 1992;304:1347-51. (23 May.)

EDITOR,—Hugh Tunstall-Pedoe and colleagues' survey of cardiopulmonary resuscitation in British hospitals has produced a considerable amount of information of value to those interested in auditing this subject.¹ We take issue with their conclusions on two points.

Firstly, they suggest that as 71% of the mortality after cardiopulmonary resuscitation occurs at the time of the initial arrest then the greatest potential for improving the survival rate is to concentrate on improving the speed of reaction, training, and competence in cardiopulmonary resuscitation. No one could disagree with the need for improved training, but the authors have not presented any evidence to support their belief that it will improve survival.

The outcome of cardiopulmonary resuscitation depends heavily on the patients selected for it. Selecting only patients in whom the technique is thought to have a high chance of success will increase the overall survival rate whereas including large numbers with a poor prognosis will decrease it. We are not told what, if any, criteria were used to exclude patients from cardiopulmonary resuscitation in the study hospitals or if these criteria were validated between hospitals. Indeed, we are not aware that any agreed policy exists in the United Kingdom for making such "do not resuscitate" decisions. We suspect that policy varies among hospitals and even among wards.

In our own hospital we recently found that 61 of 350 inpatients were classified as "do not resuscitate" but that there was considerable variation between wards and consultants. There were conflicting instructions in the medical and nursing notes for 22 of these 61 patients, and we identified several patients who were classified as for cardiopulmonary resuscitation but who would almost certainly not have survived attempts at resuscitation; this has been reported before.^{2,3} Including or excluding such patients would have a major effect on the success rate of cardiopulmonary resuscitation in this hospital, regardless of the competence and training of the resuscitation team.

Secondly, the survey does not show, as the authors suggest, that age alone adversely affects survival after cardiopulmonary resuscitation. What it does show is that those elderly patients selected for cardiopulmonary resuscitation did worse than selected younger ones. This is not surprising as elderly patients are more likely to have the sorts of conditions that make them poor candidates for successful cardiopulmonary resuscitation, but it has not yet been satisfactorily shown that age alone adversely affects outcome. In our view it is wrong to suggest that age should be used as a factor when making decisions on whether to resuscitate.

We await further results from this study with interest.

KEVIN STEWART
ADRIAN WAGG

Department of Geriatric Medicine,
Newham General Hospital,
London E13 8RU

1 Tunstall-Pedoe H, Bailey L, Chamberlain DA, Marsden AK, Ward ME, Zideman DA. Survey of 3765 cardiopulmonary resuscitations in British hospitals (the BRESUS study): methods and overall results. *BMJ* 1992;304:1347-51. (23 May.)

2 Keatinge RM. Exclusion from resuscitation. *J R Soc Med* 1989;82:402-5.

3 Stewart K, Abel K, Rai GS. Resuscitation decisions in a general hospital. *BMJ* 1990;300:705.

EDITOR,—Hugh Tunstall-Pedoe and colleagues report that 21% of their attempts at cardiopulmonary resuscitation in hospital resulted in survival to discharge.¹ The proportion may be higher than that in other studies for two methodological reasons.

Firstly, the potential for underreporting is mentioned. My impressions from a similar study were that attempts at resuscitation were less likely to be recorded if they were in some way embarrassing and that this applied particularly to cases in which the clinical team had simply not got around to telling the nurses that a dying patient was not for resuscitation.² Tunstall-Pedoe and colleagues identified arrests by designating a member of the resuscitation team to report the arrest. The success rate may therefore be greater than would have been reported if the records of dead patients had been checked for evidence of an attempt at resuscitation.

Secondly, multiple attempts at resuscitation may have been entered into the study during one episode of care (only recurrences within 24 hours were excluded). This may either increase or decrease the rate of survival to discharge compared with the rate in a study that recorded only the first attempt in an episode.

The authors comment that "hospital resuscitation is seen to be cost effective." The report contains no mention of costs; I hope that some factual basis for this assertion will be provided in due course. Unpublished data suggest that the costs are appreciable, arising mostly from treatment after resuscitation. The resources used on patients who survive for a short period with an extremely poor quality of life form an element of this cost that may be avoided with improvements to the present policy on resuscitation. The actual

attempts at resuscitation, equipment, and training may be relatively cheap.

R KEATINGE

London SE1 7RB

1 Tunstall-Pedoe H, Bailey L, Chamberlain DA, Ward ME, Zideman DA. Survey of 3765 cardiopulmonary resuscitations in British hospitals (the BRESUS study): methods and overall results. *BMJ* 1992;304:1347-51. (23 May.)

2 Keatinge RM. Exclusion from resuscitation. *J R Soc Med* 1989;82:402-5.

EDITOR,—Hugh Tunstall-Pedoe and colleagues' study of cardiopulmonary resuscitation rightly used survival as its main outcome measure.¹ It would be interesting to know, however, how many of the discharged survivors were functionally impaired and whether there were many patients who survived but required a long term stay in hospital or nursing care. Quality of life is surely as important as survival itself.

Roughly a quarter of the patients studied were over 75. The initial success rates were lower in this elderly group, consistent with the findings of a recent study.² This raises the question of the appropriateness of the attempt at resuscitation in these cases. Premorbid factors such as pneumonia, hypotension, renal failure, cancer, and a home bound lifestyle are associated with a reduced likelihood of successful resuscitation³ and are more commonly present in the elderly population. This should not discourage resuscitation of elderly people, but it underlines the importance of giving careful consideration to their resuscitation status.

ELIZABETH SMITH

Department of Geriatric Medicine,
St George's Hospital Medical School,
London SW17 0RE

1 Tunstall-Pedoe H, Bailey L, Chamberlain DA, Marsden AK, Ward ME, Zideman DA. Survey of 3765 cardiopulmonary resuscitations in British hospitals (the BRESUS study): methods and overall results. *BMJ* 1992;304:1347-51. (23 May.)

2 O'Keefe S, Redahan C, Keane P, Daly K. Age and other determinants of survival after in-hospital cardiopulmonary resuscitation. *Q J Med* 1991;81:1005-10.

3 Bedell SE, Delbanco TL, Cook EF, Epstein GH. Survival after cardiopulmonary resuscitation in the hospital. *N Engl J Med* 1983;309:569-76.

EDITOR,—It is understandable but far from excusable when authors make hyperbolic claims on the basis of research they have performed. It is neither understandable nor excusable when journal editors and referees allow authors' unjustified assertions to appear in print. With absolutely no prior reference to the costs entailed, cost effectiveness, or comparisons with other interventions Hugh Tunstall-Pedoe and colleagues conclude the paper reporting their survey of cardiopulmonary resuscitation in British hospitals with the sentence: "Compared with other life saving interventions hospital resuscitation is seen to be cost effective and a procedure where improved speed of reaction, training, and competence should produce improved results."¹

The totally unsubstantiated assertion contained in the first part of the sentence and the slightly guarded prediction made in the second part are afforded such importance by the authors that they provide the basis (or at least I assume that they provide the basis as I can find no other in the article) for a recommendation concerning resource allocation which is found only in the abstract: "Hospital resuscitation is life saving and cost effective and warrants appropriate attention, training, coordination, and equipment."

It is all very well for the *BMJ* to be strict about ethics, sexism, methodological rigour, appropriate statistical tests, and correct referencing, but it should not forget to ensure that authors' conclusions are justified by the evidence presented, or that the abstract is abstracted from the contents of