Asthma and open cast mining

EDITOR,—We agree with J M F Temple and A M Sykes that there is widespread public concern about industrial emissions and that the health effects of major industrial developments should be assessed.¹ It is important that such studies should be epidemiologically sound, and we have grave misgivings concerning both the design of this study and the conclusions drawn from it. There are many aspects of design and analysis on which comment could be made, but we shall restrict ourselves to two fundamental areas.

To say that a hypothesis has been proved in a study one must first exclude bias. The authors dismiss observer bias as impossible as they were unaware of the opening date of the open cast mine, but bias could have arisen in other ways. The authors may have been blind to the starting of open cast operations, but it is difficult to believe that the entire local population was unaware of such a major operation. The *a priori* hypothesis of the study was that the weekly number of new episodes of asthma treated by the town's general practitioners would increase when the new mine started. In view of the fact that the practice had stated, at a public inquiry, that open cast mining would aggravate its patients' asthma, it is not surprising that the weekly number of new episodes increased.

The second issue is that of statistical association and inference of causation. Although there seems to be a statistical association between the start of mining and an increase in asthma, this does not mean that the two are causally related. Some of the more pertinent issues such as dose-response relation cannot be estimated because of a lack of information. Although the diagnosis of asthma was made using a "standard definition," no details are given of what the definition was, or whether cases included patients with newly diagnosed asthma or exacerbation of asthma in existing patients. Also, although the environmental effects of the open cast mine are likely to change from day to day and week to week depending on the operations and prevailing weather, the increase in asthma remained remarkably constant. The lack of direct environmental measurements is a flaw in the study that cannot easily be overcome, and without such measurements exposure cannot simply be assumed.

Finally, the authors are quite wrong in suggesting that there are few studies of the effect of industries on the general community. A recent study by Symington *et al* of respiratory symptoms in children at schools near a foundry² is one of many environmental studies which have been performed using adequate epidemiological methods. There are also more recent cancer studies than those cited.^{3,4}

In environmental exposures, which are generally associated with small increases in risk, adequacy of study design and unbiased reporting are essential, neither of which is a feature of this study.

DAVID MCBRIDE
JERRY BEACH
IAN CALVERT

Institute of Occupational Health,
University of Birmingham,
Birmingham B15 2TT

1 Temple JMF, Sykes AM. Asthma and open cast mining. *BMJ* 1992;305:396-7. (15 August.)

2 Symington P, Coggon D, Holgate S. Respiratory symptoms in children at schools near a foundry. *Br J Ind Med* 1991;48:588-91.

3 Lloyd O, Ireland E, Tyrrell H, Williams F. Respiratory cancer in

Advice to authors

Priority will be given to letters that are less than 400 words long and are typed with double spacing. All authors should sign the letter. Please enclose a stamped addressed envelope for acknowledgment.

a Scottish industrial community: a retrospective case-control study. *J Soc Occup Med* 1986;36:2-8.

4 Brown LM, Pottern LM, Blot WJ. Lung cancer in relation to environmental pollutants emitted from industrial sources. *Environmental Research* 1984;34:250-61.

EDITOR,—J M F Temple and A M Sykes report an increase in frequency of new episodes of asthma among patients in the Glynneath practice during 1990-2 and relate this in time to open cast coal mining at the Derlwyn site.¹ We are not convinced that the increase is due to dust from the open cast mine.

The criteria for making the diagnosis of asthma are not stated. In view of well known inconsistencies in clinical diagnosis of the condition, and resulting opportunities for subconscious bias, we need to know how new episodes of asthma were identified and defined. How many individuals suffered the increased episodes of asthma—that is, did these occur as an increased frequency of attack in people already diagnosed as asthmatic or as first occurrences in people previously unaffected, or both?

The authors would have been well aware of the start of this mining activity since Dr Temple's interest in asthma in relation to open cast mining was well publicised locally (including his involvement in the public inquiry). Has this interest encouraged asthmatic patients to attend his surgery more readily?

The *cusum* technique of statistical presentation used is in principle fine for the data being illustrated but is unconventional in medical journals and epidemiology. Those unfamiliar with *cusum* methods might (wrongly) conclude that the situation was deteriorating, because the graph generally rises from October 1990 onwards. But the rise is steeper between October and (say) April than in the following six months, April till October—the drier months, when more dust might have been expected to be generated. The levels of attendance during these six months are not very different from those during the period before mining started.

Any dust generated by mining is heavily diluted as it leaves the site. Measurements indicated that emissions of respirable dust from Derlwyn were insignificant in comparison with dust borne on winds from the Continent.² Coarse (inhalable) dust is released in very low concentrations, all measured perimeter concentrations being less than 15% of the occupational exposure standard.³ Extensive research in the United Kingdom and elsewhere over more than 40 years has not shown asthmatic effects of exposure to coal mine dust from deep mining. Nor is there any history of anecdotal evidence in coal mining communities of excess asthma in miners or their families.

On general grounds, open cast mining could be expected to improve community health by bringing jobs and increased prosperity to the area. Un-

warranted conclusions attributing asthma to open cast mining only serve to provide speculative news to the media and cause unnecessary anxiety to the public.

A S AFACAN

Director of Medical Service,
British Coal Corporation,
Eastwood,
Nottinghamshire NG16 3EB

1 Temple JMF, Sykes AM. Asthma and opencast mining. *BMJ* 1992;305:396-7. (15 August.)

2 British Coal Corporation, Operations Department. *Assessment of dust emission from surface coal mining operations*. Luxembourg: Commission of the European Communities, Directorate General for Employment, Industrial Relations and Social Affairs, Health and Safety Directorate, 1992. (Final report of ECSC Research Project No 7263-02/077/08.)

3 Health and Safety Executive. *Occupational exposure limits 1992, guidance note EH 40/92*. London: HMSO, 1992.

EDITOR,—We thank David McBride and colleagues and A S Afacan for their interest in our short report. Their major concern is that of bias, suggesting that our prediction of an increased level of asthma once mining activities started was a self fulfilling prophecy. We recognise the seriousness of this suggestion but note five points: the data gathering was performed blind until 31 January 1991; once "sight was restored" no increase in rate was observed; the change in rate occurred some three months after preliminary site operations started and it coincided with the start of actual mining; although there was week to week variability (see the *cusum* chart) the increase was maintained from the start of mining activities for at least 16 months; and the site, though within 2 km of the centre of Glynneath, has been carefully designed to minimise impact on the local community.

In view of these points, it is unlikely that any bias effect of the public inquiry and local attitudes to open cast mining could result in such a sustained and immaculately timed increase.

A further concern of our critics is the lack of definition of an episode of asthma. This was omitted for reasons of space. We emphasise that the details of the definition are less important than the knowledge that it was applied consistently throughout the period of the study.

We are also taken to task on the vexed issue of statistical association and causation, though we have assumed no causative mechanism, nor do we need to do so to test the hypothesis stated. Our analysis clearly enables us to reject the null hypothesis but of course does not indicate how the change was effected.

Afacan suggests that readers of the *BMJ* would wrongly conclude from the *cusum* chart that "the situation was deteriorating," whatever that might mean. The message of the *cusum* chart is clear and concise: the initial zero gradient indicates variation around the reference mean (the initial rate of asthma episodes). Subsequent (after mining) positive gradients indicate rates always in excess of the reference mean.

Afacan states that on site measurements indicate that emissions of respirable dust are insignificant in comparison with dust borne on the winds from the Continent. We understand (from a helpful meeting with representatives from British Coal) that this is based on on site measurements from four monitors that were subject to breakdown and calibration problems. We look forward to examining these data ourselves and appraising this assertion. Of course,