Asthma and open cast mining

EDITOR,—We agree with J M F Temple and A M Sykes that there is widespread public concern about industrial emissions and that the health effects of major industrial developments should be assessed.¹ It is important that such studies should be epidemiologically sound, and we have grave misgivings concerning both the design of this study and the conclusions drawn from it. There are many aspects of design and analysis on which comment could be made, but we shall restrict ourselves to two fundamental areas.

To say that a hypothesis has been proved in a study one must first exclude bias. The authors dismiss observer bias as impossible as they were unaware of the opening date of the open cast mine, but bias could have arisen in other ways. The authors may have been blind to the starting of open cast operations, but it is difficult to believe that the entire local population was unaware of such a major operation. The *a priori* hypothesis of the study was that the weekly number of new episodes of asthma treated by the town's general practitioners would increase when the new mine started. In view of the fact that the practice had stated, at a public inquiry, that open cast mining would aggravate its patients' asthma, it is not surprising that the weekly number of new episodes increased.

The second issue is that of statistical association and inference of causation. Although there seems to be a statistical association between the start of mining and an increase in asthma, this does not mean that the two are causally related. Some of the more pertinent issues such as dose-response relation cannot be estimated because of a lack of information. Although the diagnosis of asthma was made using a "standard definition," no details are given of what the definition was, or whether cases included patients with newly diagnosed asthma or exacerbation of asthma in existing patients. Also, although the environmental effects of the open cast mine are likely to change from day to day and week to week depending on the operations and prevailing weather, the increase in asthma remained remarkably constant. The lack of direct environmental measurements is a flaw in the study that cannot easily be overcome, and without such measurements exposure cannot simply be assumed.

Finally, the authors are quite wrong in suggesting that there are few studies of the effect of industries on the general community. A recent study by Symington *et al* of respiratory symptoms in children at schools near a foundry² is one of many environmental studies which have been performed using adequate epidemiological methods. There are also more recent cancer studies than those cited.^{3,4}

In environmental exposures, which are generally associated with small increases in risk, adequacy of study design and unbiased reporting are essential, neither of which is a feature of this study.

DAVID MCBRIDE
JERRY BEACH
IAN CALVERT

Institute of Occupational Health,
University of Birmingham,
Birmingham B15 2TT

1 Temple JMF, Sykes AM. Asthma and open cast mining. *BMJ* 1992;305:396-7. (15 August.)

2 Symington P, Coggon D, Holgate S. Respiratory symptoms in children at schools near a foundry. *Br J Ind Med* 1991;48:588-91.

3 Lloyd O, Ireland E, Tyrrell H, Williams F. Respiratory cancer in

Advice to authors

Priority will be given to letters that are less than 400 words long and are typed with double spacing. All authors should sign the letter. Please enclose a stamped addressed envelope for acknowledgment.

a Scottish industrial community: a retrospective case-control study. *J Soc Occup Med* 1986;36:2-8.

4 Brown LM, Pottern LM, Blot WJ. Lung cancer in relation to environmental pollutants emitted from industrial sources. *Environmental Research* 1984;34:250-61.

EDITOR,—J M F Temple and A M Sykes report an increase in frequency of new episodes of asthma among patients in the Glynneath practice during 1990-2 and relate this in time to open cast coal mining at the Derlwyn site.¹ We are not convinced that the increase is due to dust from the open cast mine.

The criteria for making the diagnosis of asthma are not stated. In view of well known inconsistencies in clinical diagnosis of the condition, and resulting opportunities for subconscious bias, we need to know how new episodes of asthma were identified and defined. How many individuals suffered the increased episodes of asthma—that is, did these occur as an increased frequency of attack in people already diagnosed as asthmatic or as first occurrences in people previously unaffected, or both?

The authors would have been well aware of the start of this mining activity since Dr Temple's interest in asthma in relation to open cast mining was well publicised locally (including his involvement in the public inquiry). Has this interest encouraged asthmatic patients to attend his surgery more readily?

The *cusum* technique of statistical presentation used is in principle fine for the data being illustrated but is unconventional in medical journals and epidemiology. Those unfamiliar with *cusum* methods might (wrongly) conclude that the situation was deteriorating, because the graph generally rises from October 1990 onwards. But the rise is steeper between October and (say) April than in the following six months, April till October—the drier months, when more dust might have been expected to be generated. The levels of attendance during these six months are not very different from those during the period before mining started.

Any dust generated by mining is heavily diluted as it leaves the site. Measurements indicated that emissions of respirable dust from Derlwyn were insignificant in comparison with dust borne on winds from the Continent.² Coarse (inhalable) dust is released in very low concentrations, all measured perimeter concentrations being less than 15% of the occupational exposure standard.³ Extensive research in the United Kingdom and elsewhere over more than 40 years has not shown asthmatic effects of exposure to coal mine dust from deep mining. Nor is there any history of anecdotal evidence in coal mining communities of excess asthma in miners or their families.

On general grounds, open cast mining could be expected to improve community health by bringing jobs and increased prosperity to the area. Un-

warranted conclusions attributing asthma to open cast mining only serve to provide speculative news to the media and cause unnecessary anxiety to the public.

A S AFACAN

Director of Medical Service,
British Coal Corporation,
Eastwood,
Nottinghamshire NG16 3EB

1 Temple JMF, Sykes AM. Asthma and opencast mining. *BMJ* 1992;305:396-7. (15 August.)

2 British Coal Corporation, Operations Department. *Assessment of dust emission from surface coal mining operations*. Luxembourg: Commission of the European Communities, Directorate General for Employment, Industrial Relations and Social Affairs, Health and Safety Directorate, 1992. (Final report of ECSC Research Project No 7263-02/077/08.)

3 Health and Safety Executive. *Occupational exposure limits 1992, guidance note EH 40/92*. London: HMSO, 1992.

EDITOR,—We thank David McBride and colleagues and A S Afacan for their interest in our short report. Their major concern is that of bias, suggesting that our prediction of an increased level of asthma once mining activities started was a self fulfilling prophecy. We recognise the seriousness of this suggestion but note five points: the data gathering was performed blind until 31 January 1991; once "sight was restored" no increase in rate was observed; the change in rate occurred some three months after preliminary site operations started and it coincided with the start of actual mining; although there was week to week variability (see the *cusum* chart) the increase was maintained from the start of mining activities for at least 16 months; and the site, though within 2 km of the centre of Glynneath, has been carefully designed to minimise impact on the local community.

In view of these points, it is unlikely that any bias effect of the public inquiry and local attitudes to open cast mining could result in such a sustained and immaculately timed increase.

A further concern of our critics is the lack of definition of an episode of asthma. This was omitted for reasons of space. We emphasise that the details of the definition are less important than the knowledge that it was applied consistently throughout the period of the study.

We are also taken to task on the vexed issue of statistical association and causation, though we have assumed no causative mechanism, nor do we need to do so to test the hypothesis stated. Our analysis clearly enables us to reject the null hypothesis but of course does not indicate how the change was effected.

Afacan suggests that readers of the *BMJ* would wrongly conclude from the *cusum* chart that "the situation was deteriorating," whatever that might mean. The message of the *cusum* chart is clear and concise: the initial zero gradient indicates variation around the reference mean (the initial rate of asthma episodes). Subsequent (after mining) positive gradients indicate rates always in excess of the reference mean.

Afacan states that on site measurements indicate that emissions of respirable dust are insignificant in comparison with dust borne on the winds from the Continent. We understand (from a helpful meeting with representatives from British Coal) that this is based on on site measurements from four monitors that were subject to breakdown and calibration problems. We look forward to examining these data ourselves and appraising this assertion. Of course,

even if it is true on site, the nature of the dusts involved and the quantities affecting the local population could well be different.

Afacan's allusion to research into occupational exposures in the mining industry is irrelevant in the community setting and his reliance on occupational exposure standards is unwise. The numbers of people employed in open cast mining is low, so it is unlikely to improve community health by bringing jobs and increased prosperity to the area. Chambers has eloquently expressed the feelings that the "spectre of open cast mining" brings to a community.¹

We would welcome a major epidemiological study to evaluate the size of the risk and the nature of the mechanisms involved, but in 1989 the Secretary of State for Wales felt that no such study was required.

J M F TEMPLE

Glynneath,
West Glamorgan SA11 5AL

A M SYKES

University College,
Swansea

1 Chambers J. Paradise postponed. *BMJ* 1988;297:1688-90.

Exercise, fitness, and health

EDITOR,—Daphne Gloag's editorial confirms what all exercise enthusiasts have been saying for years: that exercise is good for us and that most people do not take enough.¹ Gloag concludes that all doctors should ask about exercise when they see patients and advise on suitable exercise and local facilities. Doctors may also, however, unwittingly reduce some patients' capacity to take exercise by prescribing drugs that may interfere with exercise metabolism. Endurance exercise has specific benefits by favourably modifying lipid profiles.² Any endurance exercise of moderate intensity (about 50% maximal oxygen uptake) relies heavily on oxidation of fat for resynthesis of ATP and production of energy, thereby preserving carbohydrate stores. Drugs that modify the metabolism of fat and reduce the availability of fatty acids for oxidation would therefore be expected to have an impact on a patient's capacity to take exercise. Our recently completed trial suggests that a lipid lowering drug (a derivative of nicotinic acid) considerably reduced oxidation of fat, a fibrate had an intermediate effect, and a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor had no effect.

A suitably designed exercise regimen is often prescribed as part of a rehabilitation programme for patients suffering from hypertension or angina or recovering from myocardial infarction. Concurrent drug treatment for such patients may include a β blocker, which has a cardioprotective effect.³ Our observations suggest that in comparable doses β_1 selective agents may have less of an impact on total oxidation of fat than non-selective agents. During one hour's exercise the mean proportion of energy supplied by fat in eight healthy men was 41.5% (SD 10.9%) with a placebo, 34.2% (10%) with a β_1 selective drug, and 28.6% (8.2%) with a non-selective drug. Furthermore, β_1 selective drugs reduce the unwanted β_2 mediated effects on glucose metabolism and peripheral vasoconstriction associated with non-selective drugs and should therefore have less of an impact on exercise metabolism. Doctors might also give advice on the timing of exercise as with most β blockers peak plasma concentrations are reached one to three hours after an oral dose, with exercise being more difficult during this period.

Thus care should be taken that concurrent prescription of drugs and exercise are not in conflict. A patient taking a non-selective β blocker and a fibrate (a not uncommon combination) might have considerable difficulty in taking modest

exercise such as a brisk walk for one to two hours. Further studies are needed to find therapeutic regimens that reduce patients' risk of suffering from the effects of coronary artery disease but also allow exercise which may have the same goals and, in addition, improve the quality of life.

MARTIN J KENDALL
ANTHONY HEAD

Clinical Investigation Unit,
Department of Medicine,
Queen Elizabeth Hospital,
Birmingham B15 2TH

- 1 Gloag D. Exercise, fitness, and health. *BMJ* 1992;305:377-8. (15 August.)
- 2 Dufaux B, Assman G, Hollman W. Plasma lipoproteins and physical activity. *Int J Sports Med* 1982;3:123-36.
- 3 Olsson G, Wikstrand J, Warnold I, Manger Cats V, McBoyle D, Herlitz J, et al. Metoprolol-induced reduction in postinfarction mortality: pooled results from five double blind randomised trials. *Eur Heart J* 1992;13:28-32.

EDITOR,—Daphne Gloag believes that there is no dispute that exercise is beneficial.¹ Although over 800 deaths have occurred in accidents during sports and leisure exercise in the last five reported years (1986-90)² as well as other deaths during exercise from pre-existing disease, the benefits of exercise with regard to mortality are well established, at least from the public health point of view. From this vantage, discounting the future health gains derived from fitness to set against the present hazards may not be appropriate.³ Last Saturday, however, when I played cricket in a team without its regular wicket keeper, who was in hospital recovering from a heart attack that occurred during a game of squash, against a team missing a player who had died after being struck by lightning during a cricket match, the balance was not so clear.

Furthermore, taking morbidity into account leaves matters still less clear cut. In 1988 over half a million people aged 16 and over and 200 000 children attended accident and emergency departments in the United Kingdom after sports accidents occurring outside the home (unpublished data from home accident surveillance system 1988, Consumer Safety Unit of Department of Trade and Industry). In 1987 the general household survey included 184 adults who had had sports accidents that resulted in contact with a doctor.⁴ These patients took 670 days off work as a result. Translating these small numbers into national estimates yields an estimated 1.5 million injuries related to exercise needing contact with a doctor and 5.5 million days lost from work. A report, commissioned by the Sports Council of England, on a national survey of injuries and illnesses related to sports and exercise in 17 500 adults, which should be published later this year, shows that even these numbers underestimate the problem. In addition, many non-trivial injuries occur which are treated by physiotherapists, dentists, complementary therapists, nurses, and others, and there is the possibility of chronic conditions developing later in life.

Thus as well as indisputably conferring health benefits, exercise indisputably incurs substantial health costs. As always, the trick is to weigh these carefully against each other to determine whether exercise is, in fact, beneficial.

JON NICHOLL

Medical Care Research Unit,
Department of Public Health Medicine,
Medical School,
Sheffield S10 2RX

- 1 Gloag D. Exercise, fitness, and health. *BMJ* 1992;305:377-8. (15 August.)
- 2 Office of Population Censuses and Surveys. Fatal accidents occurring during sporting and leisure activities 1986-1990. London: HMSO, 1988-91. (DH4 88/3, 88/6, 89/4, 90/4, 91/4.)
- 3 Discounting health care: only a matter of timing [editorial]. *Lancet* 1992;340:148-9.
- 4 Office of Population Censuses and Surveys. *General household survey, 1987 data*. London: HMSO, 1990.

Validating the SF-36

EDITOR,—We would like to draw attention to a number of errors and anomalies of interpretation in Brazier and colleagues' paper on validating the SF-36 health survey questionnaire.¹

Firstly, it is inappropriate to attempt to validate a health survey questionnaire (the SF-36) against a measure of perceived distress (the Nottingham health profile). In addition to measuring different concepts, the items in the two measures were generated by different methods. In the SF-36, the items represent professionals' assumptions about issues of relevance to health status. The items in the Nottingham health profile were derived directly from patients and other lay people, describing their perceptions of ill health. It has also been made clear in the major publication on the measure that the Nottingham health profile is not suitable for use in general population or general practice surveys. The instrument works best with selected groups, particularly chronically ill and elderly patients.² Furthermore, presenting two similar measures together is likely to lead to response bias.

Secondly, the paper does not report one of the Nottingham health profile's sections at all: sleep. Since sleep problems are common in ill health this would appear to be an important omission from the SF-36, possibly resulting from the method by which items were generated.

Thirdly, the authors claim that the test-retest values obtained for the SF-36 indicate excellent reliability. This is in fact incorrect, as the values are lower than those generally accepted as indicating that a scale is reliable.³

Lastly, if a questionnaire cannot be shown to be reliable then there is little value in testing its validity. However, even in the authors' own terms the validity of the SF-36 was not shown. The socioeconomic class gradients are not as consistent as would be expected on several of the dimensions. Two of the subscales do not correspond to the presence of chronic physical problems, and the correlations between dimensions do not accord with logic (vitality and mental health correlated 0.69).

A close inspection of the data presented in the paper suggests that they do not support either the reliability or the validity of the SF-36 in the United Kingdom. It is noteworthy that nothing has been published on the SF-36 in its country of origin.

We welcome this attempt to find a much needed measure for assessing health in the general population. However, the choice of a measure, the techniques by which it is tested, and the interpretation of data require more careful attention.

SONJA M HUNT
STEPHEN P McKENNA

Care Outcomes Group,
Rheumatology and Rehabilitation Research Unit,
School of Medicine,
University of Leeds,
Leeds LS2 8NZ

- 1 Brazier JE, Harper R, Jones NMB, O'Cathain A, Thomas KJ, Usherwood T, et al. Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ* 1992;305:160-4. (18 July.)
- 2 Hunt SM, McEwen J, McKenna SP. *Measuring health status*. London: Croom Helm, 1986.
- 3 McDowell I, Newell C. *Measuring health: a guide to rating scales and questionnaires*. New York: Oxford University Press, 1987.

EDITOR,—It is vital that the validity and reliability of health status measures are thoroughly investigated. Test-retest reliability is particularly important because measurement error can make it more difficult to detect real differences between populations. Although Brazier and colleagues have carried out a large and well designed evaluation of their version of the SF-36 health survey questionnaire,¹ we found their analysis of test-retest reliability incomplete. This aspect of their paper is highlighted by the claim of a level of test-retest