and fertilisation rate ($p=0.001$) but not for indication ($p=0.08$) or age ($p=0.9$). The group who stopped treatment had a higher rate of total fertilisation failure, a lower fertilisation rate, and a generally more unfavourable indication for in vitro fertilisation. These results indicate that the assumption of the life table method is not met because patients who decide to stop treatment are likely to have less chance of conceiving by in vitro fertilisation than patients who continue.

Few studies giving cumulative pregnancy rates after several cycles of in vitro fertilisation evaluate possible selection bias in the couples who continue treatment. To avoid too high expectations people should be aware of the possibility of such selection, particularly in a study with the objective "to provide reliable prognostic information for couples seeking assisted conception."

ERTEVELDE
JKOUDSTAAL

Section of Reproductive Medicine,
Division of Obstetrics and Gynaecology,
University Hospital Utrecht,
POB 85500,
3508 GA Utrecht,
Netherlands

JMEIMERS

Department of Public Health and Social Medicine,
Erasmus University,
Rotterdam

1 Hull MGR, Eddowes HA, Fahy U, Abuzeid MI, Mills MS, Cahill DJ, *et al.* Expectations of assisted conception for infertility. *BMJ* 1992;304:1465-9. (6 June.)

Reaccrediting general practice

EDITOR,—John Russell reports a small number of problems with assessment visits in general practice about which he has been informed anecdotally.¹ Against this, however, must be set the view of over 24 000 general practitioners, who responded "Disagree or strongly disagree" to the statement "There is no need for the profession to consider the possibility of a system of professional accreditation and re-accreditation at the present time" as follows: general practitioners with "no involvement with training"—49.1%; general practitioners with "training provided in the practice"—60.9%; general practitioners who "provide training personally"—70.1%.² It is therefore still logical to conclude that those general practitioners who have had personal experience of the trainer system, which involves about a quarter of all general practices in Britain, are more in favour of reaccreditation than those who have not.

Chris Nancollas emphasises the importance of consultation.³ The General Medical Services Committee represents all general practitioners in the NHS and regularly communicates with all local medical committees in the country. Similarly, the Royal College of General Practitioners is committed to its local faculties, which cover the whole of Britain.⁴

Developments in medicine worldwide have placed greater emphasis on continuing education and assessment or reaccreditation, or both, for all health professionals. It seems wise for the medical profession to continue to work on this, and it is encouraging that both Russell and Nancollas also believe that assessment of general practitioners through peer review is preferable to other methods, such as re-examination.

DENIS PEREIRA GRAY

Exeter EX2 4TJ

1 Russell J. Reaccrediting general practice. *BMJ* 1992;305:835. (3 October.)

2 General Medical Services Committee. *Your choices for the future. A survey of GP opinion. UK report.* London: Electoral Reform Ballot Services, 1992.

3 Nancollas C. Reaccrediting general practice. *BMJ* 1992;305:835. (3 October.)

4 Royal College of General Practitioners. *A college plan. Priorities for the future.* London: RCGP, 1990. (Occasional paper 49.)

Guidance on guidelines

EDITOR,—Andrew Haines and Gene Feder's editorial on clinical guidelines draws attention to issues that need to be resolved if Britain is to avoid the American experience of producing thousands of guidelines of unproved efficacy.¹ Many of these issues were raised last year at a conference to assess the progress made in developing guidelines in Britain.^{2,3} Three crucial issues surfaced: how guidelines can be produced and owned locally but not be perceived as being inferior to guidelines published by national bodies; how to ensure that they are updated regularly; and how to ensure their effective dissemination and use.

A national "clearing house" may seem to be an attractive means of collating and disseminating information. The experience of the King's Fund Centre in setting up a centralised system for collating details of good practice in medical audit, however, suggests that this process in isolation is of limited value (personal communication). The research and development initiative provides an alternative model, in which the provision is organised at a regional level.⁴

Guidelines are an important component of the development of explicit standards of care and thus are intimately connected with medical audit. It may be opportune, therefore, to integrate the "guideline industry" with the medical audit initiative; a strong medical audit structure already exists at regional level. In South West Thames, on the basis of 13 years' experience of producing hospital guidelines, we have established a treatment guidelines centre at St George's Hospital Medical School. This unit will disseminate information on national and local guidelines and also provide a publishing service for individual units. Its aim is to facilitate the development of local guidelines rather than to standardise practice across the region.

P LITTLEJOHNS
J COLLIER
S HILTON

St George's Hospital Medical School,
London SW17 0RE

1 Haines A, Feder G. Guidance on guidelines. *BMJ* 1992;305:785-6. (3 October.)

2 Collier J, Ismail I, Littlejohns P. Guidelines for treatment in NHS hospitals. *Medical Audit News* 1992;2:141-3.

3 Richardson J. Treatment guidelines and medical audit. *Lancet* 1991;338:877.

4 *Research for health: a research and development strategy for the NHS.* London: HMSO, 1991.

The death penalty

EDITOR,—In his editorial Richard J Bonnie mentions a recommendation of a BMA working party that certification of death should always take place away from the site of an execution.¹

That is not enough. The certifying doctor has to be removed from the execution with regard to both place and time (so as not to monitor the execution by, for example, electronically transmitted electrocardiography). Certification should be issued for medicolegal purposes only at a time and place where there is no longer any non-medical doubt about death having occurred.

The reason for this requirement is the sad fact that doctors are probably always present at executions (in the United States and elsewhere). Their "certification of death" amounts in many cases to active participation in the execution itself (monitoring heartbeats, etc and then accordingly ordering more voltage or more intravenous poison if deemed necessary) until they can certify that death has occurred and the executional process can be stopped, thereby making them de facto the chief executioners.

The request for certification was meant to be for the issuing of a medicolegal (written) statement

that death had irrevocably occurred at some earlier time. Instead, many doctors have twisted "certification" into meaning "certain making," from the original Latin meaning of the word *certificare*, thereby allowing them the actions mentioned above.

The Council of Judicial and Ethical Affairs of the American Medical Association is working on a new and stricter resolution, hopefully much along the lines of the resolution adopted by the Nordic medical associations in 1986, which in part reads: "For a physician to prepare, administer or monitor any procedure with a view to injuring a human being or to train others to do so would be a perversion of medical knowledge and skill and of the physician's responsibility to and role in society. Thus the medical associations of the Nordic countries... declare it indefensible for any physician to participate in any act connected to and necessary for the administration of capital punishment."

SOEREN GANES

9850 Hirtshals,
Denmark

1 Bonnie RJ. The death penalty. *BMJ* 1992;305:381-2. (15 August.)

Health checks for people over 75

EDITOR,—We were disturbed to read about the inadequacies in the system of annual health checks for people over 75, especially those relating to the detection of dementia.^{1,2} The shortcomings included lack of follow up of people not responding to the offer of a health check, the fact that inexperienced nurses performed the assessments, and a lack of national guidelines.

Detecting early dementia is important as it probably lessens the need for crisis intervention for patients' and relatives' support and because treatments likely to be available soon will depend on early diagnosis for best effect. Regular screening of elderly people is potentially useful for detecting dementia. We recently canvassed 37 local practices to examine what cognitive assessment was carried out in their checks of people over 75.

We received replies from 26 of the 37 practices (70%): three were completed by general practitioners, who indicated that they did most of their practice's assessments, and the rest by practice nurses. All 26 respondents reported assessing mental function, 16 informally and 10 formally. Of those who assessed it formally, four used a test and six used specific questions, one going on to use a test if this was indicated. Three different tests were being used, most commonly the abbreviated mental test score.³ This was usually provided by the Alzheimer's Disease Society. In an open question only three of the respondents who formally tested cognition indicated that the presence of physical or sensory problems affected their interpretation of performance. Five respondents spontaneously requested advice regarding the use of suitable tests.

Detection of dementia is unlikely to be achieved effectively by informal assessment by untrained staff. Screening instruments would enhance detection if training was given in their administration and interpretation. The abbreviated mental test score is appropriate as it is short and is effective at detecting dementia⁴ and at discriminating between organic and functional illness,⁵ which is probably important. Patients who perform poorly or whose test score has decreased should subsequently be assessed by their general practitioner. Cognitive screening of people over 75 in an average practice of 2000 people would produce around 17 people needing medical assessment,⁶ though in the second year of a screening programme the number should be lower.