18% (n = 23) in the relaxation training group, and 15% (n = 20) in the control group. At six months the proportion of participants who thought that their neck had fully or considerably recovered was 53% (n = 72) in the dynamic muscle training group, 23% (n = 30) in the relaxation group, and 17% (n = 22) in the control group compared with 26% (n = 35), 23% (n = 30), and 12% (n = 16), respectively, at 12 months.

Discussion

Dynamic muscle training or relaxation training for chronic neck pain in female office workers had no effect on the intensity of pain, neck disability, or sick leave over 12 months. The training groups reported better subjective recovery than the control group, and there was a slight improvement in the cervical range of motion, but this was not clinically relevant.

Systematic reviews of conservative treatments for neck pain have looked at only a few randomised clinical trials on dynamic muscle exercises and none on relaxation exercises. None of the studies included the dynamic exercises we used. The methods most similar to ours were group instructional strategies with light gymnastics or stretching, which did not reduce pain when compared with no treatment.^{3-5,9} In one study, individual proprioceptive exercises, relaxation, and behavioural support produced an alleviation of neck pain at three months but not at 12 months.¹⁸ High intensity exercise has not been found more effective than low intensity exercise for chronic neck pain.¹⁹ Our training comprised dynamic exercises for large muscle groups using dumbbells followed by stretching of the exercised muscles, and the intensity of the exercises was increased progressively during the training programme. Studies assessing the effectiveness of muscle strengthening exercises have also had small sample sizes and weak methods.⁷⁻⁹

The large number of patients in our study led to both good comparability between the treatment groups and strong statistical power. Minor imbalances in characteristics at baseline were controlled for in the analyses. Adherence during the first three months was not complete, but the attendance rate was probably at least as good as in clinical practice. The amount and content of the interventions also complied with current practice.

We thank the physicians, physiotherapists, and nurses at the occupational healthcare centres in Tampere for their contribution.

Contributors: See bmj.com

Funding: This work was supported by a grant from the Finnish work environment fund (project No 96243). The guarantors accept full responsibility for the conduct of the study, had access to the data, and controlled the decision to publish.

Competing interests: None declared.

Ethical approval. This study was approved by the ethics committee of the Finnish Institute of Occupational Health.

What is already known on this topic

Dynamic muscle training and relaxation training are often prescribed for chronic neck pain

Reliable data on the effectiveness of these interventions compared with ordinary activity are lacking

What this study adds

Dynamic muscle training and relaxation training do not have more favourable effects on chronic neck pain over advising patients to be active

Ordinary activity leads to an outcome similar to that of dynamic muscle training or muscle relaxation training

- Kjellman GV, Skargren EI, Oberg BE. A critical analysis of randomized clinical trials on neck pain and treatment efficacy. A review of the literature. *Scand J Rehabil Med* 1999;31:139-52.
- Borghouts JA, Koes BW, Bouter LM. The clinical course and prognostic factors of non-specific neck pain. A systematic review. *Pain* 1998;77:1-13.
- Levoska S, Keinänen-Kiukaanniemi S. Active or passive physiotherapy for occupational cervicobrachial disorders? A comparison of two treatment methods with a one-year follow-up. *Arch Phys Med Rehabil* 1993;74:425-30.
- Dyrssen T, Svedenkrans M, Paasikivi J. Muskelträning vid besvär i nacke och skuldror: effektiv behandling för att minska smärtan. *Läkartidningen* 1989;86:116-20.
- Van Tulder MW, Goossens M, Hoving JL. Nonsurgical treatment of chronic neck pain. In: Nacheimson AL, Jonsson E, eds. *Neck and back pain*. Philadelphia: Lippincott Williams and Wilkins, 2000.
- Toivanen H, Helin P, Hanninen O. Impact of regular relaxation training and psychosocial working factors on neck-shoulder tension and absenteeism in hospital cleaners. *J Occup Med* 1993;35:1123-30.
- Mosston M, Ashworth S. *Teaching physical education*. 4th ed. New York: MacMillan, 1994.
- Rinne M, Toropainen E. How to lead group—practical principles and experiences of conducting a promotional group in health-related physical activity. *Patient Educ Counsel* 1998;334(suppl 1): 69-76.
- Hertling D, Jones D. Relaxation. In: Hertling D, Kessler R, eds. *Management of common musculoskeletal disorders*. Philadelphia: Lippincott, 1990:144-65.
- Denner I, Denner A, Agnischock A. Retest reliability on the CMS. Deutsche Sporthochschule Köln, Institut für Leichtathletik und Turnen 1992.
- Salokangas RK, Poutanen O, Stengård E. Screening for depression in primary care. Development and validation of the depression scale, a screening instrument for depression. *Acta Psychiatr Scand* 1995;92:10-6.
- Elo A-L. Assessment of mental stress factors at work. In: Zenz C, Dickerson OB, Hovarth EP Jr, eds. *Occupational medicine*. 3rd ed. Saint Louis, MO: Mosby-Year Book, 1994:945-59.
- Altman D, Gardner M. Calculating confidence intervals for means and their differences. In: Gardner M, Altman D, eds. *Statistics with confidence—confidence intervals and statistical guidelines*. London: British Medical Journal, 1989:20-7.
- Taimela S, Takala E-P, Asklöf T, Seppälä K, Parviainen S. Active treatment of chronic neck pain. *Spine* 2000;25:1021-7.
- Randlov A, Ostergaard M, Manniche C, Kryger P, Jordan A, Heegaard S, et al. Intensive dynamic training for females with chronic neck/shoulder pain. A randomised controlled trial. *Clin Rehabil* 1998;3:200-10. (Accepted 3 June 2003)

Endpiece

A warning

Don't scrape your insides with much roughage as it is more likely to do harm than good. Vegetarianism is harmless enough though it is apt to fill a man with wind and self-righteousness.

Sir Robert Hutchison (1871-1960) in an address to the BMA in Winnipeg, Canada, 1930

Fred Charatan, retired geriatric physician, Florida

- Bovim G, Schrader H, Sand T. Neck pain in the general population. *Spine* 1994;19:1307-9.
- Côté P, Cassidy JD, Carroll L. The Saskatchewan Health and Back Pain Survey, the prevalence of neck pain and related disability in Saskatchewan adults. *Spine* 1998;23:1689-98.
- Gross AR, Aker PD, Goldsmith CH, Peloso P. Conservative management of mechanical neck disorders. A systematic overview and meta-analysis. In: *Cochrane Library*, Issue 4. Oxford: Update Software, 2000.
- Gross AR, Aker PD, Goldsmith CH, Peloso P. Patient education for mechanical neck disorders. A systematic overview and meta-analysis. In: *Cochrane Library*, Issue 4. Oxford: Update Software, 2000.