(61, 62, and 63). Also, three of our patients who were operated on had small aneurysms initially that grew by over 1 cm in six months, a far quicker rate of growth than that described by others in non-hypertensive patients.²

On the basis of our initial findings we believe that screening should be offered to male hypertensive patients from the age of 60 and should be done regularly—for example, every three to five years—in this group.

In the 388 women we have screened eight aneurysms have been detected (only two over 3.5 cm), confirming the low incidence of aneurysms in women described by others.³ One woman, aged 72, has required an elective repair of an aneurysm of 6.1 cm. Many more hypertensive women will require screening to determine more precisely the lower incidence of aneurysms in this group. Offering screening in some form to this group may well, however, prove to be just as economical as breast cancer screening, in terms of quality adjusted life years at least.

Numerous studies have suggested an association between hypertension and abdominal aortic aneurysms. We believe that hypertensive people also "require special provision for screening irrespective of any plans to screen apparently healthy people."¹

MARTIN SOWTER
MICHAEL LEWIS

Department of Surgery,
East Glamorgan Hospital,
Pontypridd,
Mid-Glamorgan CF38 1AB

- 1 Harris PL. Reducing the mortality from abdominal aortic aneurysms: need for a national screening programme. *BMJ* 1992;305:697-8. (19 September.)
- 2 Delin A, Ohlson H, Swedenborg J. Growth rates of abdominal aortic aneurysms as measured by computed tomography. *Br J Surg* 1985;72:530-2.
- 3 Collin J. The epidemiology of abdominal aortic aneurysms. *Br J Hosp Med* 1988;39:64-7.

EDITOR,—I am sympathetic to P L Harris's objective of trying to reduce mortality from ruptured abdominal aortic aneurysms.¹ I have recently screened 678 (97.6%) of the 695 patients aged 60-79 in our practice for aneurysms. Twenty six were found to have an aneurysm (range 3.0-8.3 cm external sagittal diameter), and 13 were referred for a surgical opinion. The screening programme has exposed some of the dilemmas in current management of aneurysms.

Patients deserve to know of important risks associated with repair of an aneurysm. Harris's statement that in the best centres elective repair carries an "operative risk of under 5%" cannot be generally assumed, and published mortality statistics may not reflect the risk for an average patient.² Morbidity associated with elective repair of an aneurysm has not been widely published, but in series of mixed elective and emergency repairs it has been considerable.^{3,4} Without reference statistics on mortality and morbidity the balance of when to operate for a particular size of aneurysm and risk to the patient becomes uncomfortably difficult. For individual patients local results will be most pertinent unless distant referral is considered.

Harris rightly directs attention to aneurysms of 4.0-5.0 cm, for which management is contentious; most aneurysms detected by screening fall into this category. Surgery has been advocated for aneurysms of >4 cm,⁵ but such an aggressive policy is not supported by recent prospective^{6,7} and retrospective⁸ studies of the natural course of aneurysm. Rarely, small aneurysms will rupture fatally, but I believe that relatives find unlikely natural tragedy easier to bear than tragedy after well intentioned surgery. A more conservative approach to surgery tips the risk-benefit balance towards benefit, and Scott *et al*'s study exemplifies how such a policy has worked successfully.⁷

With regard to the psychological consequences

of detecting aneurysms by screening, will patients with small aneurysms be able to maintain a fair perspective of a low risk of rupture or will their predominant perception be of a time bomb waiting to explode within? The predicament of those with large aneurysms who are considered to be unfit for surgery is particularly unfortunate. The anxiety an aneurysm can generate should not be underestimated or disregarded.

If a low mortality and morbidity associated with elective surgery, a conservative approach to intervention, and adequate counselling of patients can be combined then I believe that a local screening policy for aneurysms could make good ethical and economical sense. That such criteria apply nationally is doubtful, and currently I do not favour a national screening programme.

Lastly, β blockade has shown promise in the management of aneurysms.¹⁰ As aneurysms are common, whether physiological β_1 adrenergic antagonism can retard their expansion or reduce the rate of rupture is of great practical importance. An extension of the Medical Research Council's small aneurysm study to address this issue would be expedient.

J S ROSE

The Surgery,
Lindford,
Hampshire GU35 0QJ

- 1 Harris PL. Reducing the mortality from abdominal aortic aneurysms: need for a national screening programme. *BMJ* 1992;305:697-9. (19 September.)
- 2 Vella V, Shandall A, Duthie G, Shute K. Aortic aneurysms—who should do them? *Ann R Coll Surg Engl* 1990;72:215-7.
- 3 Giorgetti PL, Arpesani A, Bortolani EM, Rampoldi V, Rignano A, Ghildardi G, *et al*. Abdominal aortic aneurysm surgery in over 75 year old patients. *Panminerva Med* 1991;33:19-25.
- 4 Branchereau A, Nazet J, Colavolpe J, Scotti L. Combined mortality and morbidity of direct surgical treatment of abdominal aortic aneurysms. *Ann Vasc Surg* 1990;4:10-4.
- 5 Olsen P, Schroeder T, Agerskov K, Røder O, Sørensen S, Perko M, *et al*. Surgery for abdominal aortic aneurysms. *J Cardiovasc Surg* 1991;32:636-42.
- 6 Collin J. Point of view. Elective surgery for small abdominal aortic aneurysms. *Lancet* 1987;i:909.
- 7 Scott RAP, Ashton HA, Kay DN. Abdominal aortic aneurysm in 4237 screened patients: prevalence development and management over 6 years. *Br J Surg* 1991;78:1122-5.
- 8 Guirguis EM, Barber GG. The natural history of abdominal aortic aneurysms. *Am J Surg* 1991;162:481-3.
- 9 Nevitt MP, Ballard DJ, Hallett JW. Prognosis of abdominal aortic aneurysms. *N Engl J Med* 1989;321:1009-14.
- 10 Leach SD, Toole AL, Stern H, De Natale RW, Tilson MD. Effect of β -adrenergic blockade on the growth rate of abdominal aortic aneurysms. *Arch Surg* 1988;123:606-9.

Domiciliary thrombolysis by general practitioners

EDITOR,—The results of the Grampian region early anistreplase trial¹ need to be seen within the context of the burden of myocardial infarction that is carried by a community. The general practitioners who participated clearly, and effectively, performed a great deal of selection. Recruitment of one patient every 11 months means that most patients with myocardial infarction were not entered into the trial. A local estimate for Plymouth Health Authority is of eight to 10 myocardial infarctions per general practitioner each year; on the basis of the lower figure, and assuming a 30% death rate if medical help is not called, these general practitioners' patients would have suffered 1537 myocardial infarctions, but only 311 were recruited.

Another way of looking at this is to consider the total number of deaths ascribed to myocardial infarction among patients of the doctors in the study. Extrapolation from local data for Plymouth gives an estimate of 1060 deaths over the period of the study. A considerable proportion of these will have been sudden deaths; this still leaves many more deaths than those noted during the study.

Any strategy for implementing a new advance needs to take into account the whole range of

presentations of conditions; for thrombolysis this means not only patients with classical myocardial infarction diagnosed by general practitioners but also, for example, people with atypical chest pain and those who do not perceive their symptoms as serious.

ALISON ROUND

Plymouth Health Authority,
Plymouth PL6 5XN

- 1 GREAT Group. Feasibility, safety, and efficacy of domiciliary thrombolysis by general practitioners: Grampian region early anistreplase trial. *BMJ* 1992;305:548-53. (5 September.)

EDITOR,—I am surprised that in the Grampian region early anistreplase trial no patients were diagnosed as having unstable angina, which is the most common differential diagnosis and the most difficult to make in the early stages of a myocardial infarction.¹ It is likely that the patients in the diagnostic groups "possible myocardial infarction" and "ischaemic heart disease" in fact had unstable angina. If only definite and probable myocardial infarctions are counted the diagnostic accuracy of the general practitioners was 57% (and of the hospital doctors 66%). This may also account for the lower mortality and fewer Q wave infarctions in the domiciliary group.

As there is no evidence that thrombolytic treatment is of benefit in unstable angina² it seems that nearly half the patients in the study received thrombolytic treatment inappropriately and were needlessly exposed to the risks of haemorrhage. Colleagues and I found similar figures in a study in Somerset, where the general practitioners accurately diagnosed myocardial infarction on clinical grounds (without electrocardiography in most cases) in 45% of cases (S Rule *et al*, unpublished work). Again this was largely because many patients with unstable angina were thought to be in the early stages of myocardial infarction.

Diagnosing myocardial infarction at the onset can be difficult, but at a minimum a good history should be obtained and an electrocardiogram properly interpreted. In the Grampian study the general practitioner was required to record an electrocardiogram but not to interpret it, which seems pointless. It is the electrocardiogram, however, that causes problems for many general practitioners as individually they will see few cases of myocardial infarction each year. The higher diagnostic accuracy in hospital may relate to this.

JOHN E SANDERSON

Taunton and Somerset Hospital,
Taunton,
Somerset TA1 5DA

- 1 GREAT Group. Feasibility, safety, and efficacy of domiciliary thrombolysis by general practitioners: Grampian region early anistreplase trial. *BMJ* 1992;305:548-53. (5 September.)
- 2 UNASEM Collaborative Group. Thrombolytic agents in unstable angina. *J Am Coll Cardiol* 1990;15:65A.

EDITOR,—The Grampian region early anistreplase trial provides a valuable contribution to the debate on pre-hospital thrombolysis, particularly in view of the importance of minimising the delay to treatment.¹ Several points merit additional discussion. Although the authors state that about 60% of eligible patients were recruited into the study, they do not state the proportion of all patients with myocardial infarction. The narrow time window for entry to the trial selected patients presenting early. Indeed, the median patient delay in an earlier community study by the same authors was two hours,² compared with 45 minutes in this study. Thus the improvements in outcome may not necessarily apply to patients presenting later. The median delay to presentation in recent large scale studies has been substantially longer (57% beyond four hours in the second international study of infarct survival³).

The high accuracy of diagnosis achieved in this

study in the absence of electrocardiographic criteria is not necessarily generally applicable. Unless thrombolysis is restricted to those presenting early and with classic symptoms of infarction, the proportion of alternative diagnoses (2.2%) is unlikely to be substantiated. For example, phase 1 of the myocardial infarction triage and intervention project found only one in six confirmed infarctions among those evaluated before admission to hospital.¹

The statement that "even in an urban area there would be a temporal advantage in the general practitioner giving thrombolytic therapy in the home" is untested and cannot be extrapolated from the present study. A 999 call and shortening of the delays in hospital would have reduced the difference between home and hospital treatment substantially. We have shown that in an urban area the time of administration of thrombolytic treatment after the onset of symptoms was reduced to a median of 150 minutes by the introduction of a "fast track" system.⁴

Until these issues are resolved it may be premature to advise the widespread implementation of pre-hospital thrombolysis without electrocardiographic confirmation.

A C H PELL
K A A FOX

Cardiovascular Research Unit,
Edinburgh University,
Edinburgh EH8 9XP

- 1 GREAT Group. Feasibility, safety, and efficacy of domiciliary thrombolysis by general practitioners: Grampian region early anistreplase trial. *BMJ* 1992;305:548-53. (5 September.)
- 2 Rawles JM, Haines NE. Patient and general practitioner delays in acute myocardial infarction. *BMJ* 1988;296:882-4.
- 3 ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17 187 cases of suspected acute myocardial infarction: ISIS-2. *Lancet* 1988;ii:349-60.
- 4 Weaver WD, Eisenberg MS, Martin JS, Litwin PE, Schaeffer SM, Ho MT, et al. Myocardial infarction triage and intervention project—phase 1: patient characteristics and feasibility of prehospital initiation of thrombolytic therapy. *J Am Coll Cardiol* 1990;15:925-31.
- 5 Pell ACH, Miller HC, Robertson CE, Fox KAA. Effect of "fast track" admission for acute myocardial infarction on delay to thrombolysis. *BMJ* 1992;304:83-7.

EDITOR.—Much publicity has been, and will be, given to the finding of the Grampian region early anistreplase trial that patients who received thrombolytic treatment (anistreplase) at home had 49% fewer deaths than those who received it in hospital.¹ Unfortunately, the trial was really too small to estimate reliably any reduction in mortality, and so significance could be achieved only if (because of either chance or bias) an implausibly large treatment difference was observed. In such circumstances a bayesian analysis provides a useful interpretation by setting a surprising finding in the context of more cautious prior belief.²

First one expresses prior belief about the proportionate reduction in mortality due to thrombolysis at home. Given the known benefits of early thrombolysis^{3,4} and the average two hours saved in time to treatment, it could be argued that a 15-20% reduction in mortality is highly plausible, while the extremes of no benefit and a 40% reduction are both unlikely. The figure (a) shows such a distribution of prior belief. This prior is compatible with the results of the European myocardial infarction project, in which the same drug was given to over 5000 patients.

In the Grampian region early anistreplase trial 23 of the 148 patients who received home thrombolysis died within three months compared with 13 of the 163 who received hospital thrombolysis. This is displayed in the figure (b). The observed 49% reduction is the mode of this distribution, and the 2% tail area beyond no effect indicates $p=0.02$ one sided. The widely spread distribution illustrates the inevitable uncertainty with only 36 deaths in total.

Using Bayes's theorem, we have combined the prior belief and likelihood to produce a posterior belief distribution (figure (c)). This quantifies how opinion on the efficacy of home thrombolysis should be affected by the limited amount of highly positive data in the Grampian region early anistreplase trial. The peak of the posterior distribution is a 25% reduction in mortality, with a 95% confidence interval from no effect to a 43% reduction. Thus belief is shifted in a positive direction, but not by much, and, specifically, a halving of mortality remains implausible.

Bayesian analysis of data from Grampian region early anistreplase trial

Perhaps the Grampian region early anistreplase trial was just lucky. For instance, based on the figure (a) a difference of 23 versus 13 deaths or more should occur with probability 0.1. We are also concerned, however, about the emphasis on three month mortality (not a predefined end point), the lack of independent monitoring of data, the randomisation method, and the early stopping of the trial.

Overall, such an important therapeutic issue requires larger scale trials which can quantify the treatment effect precisely. Here we seem faced with publication bias. A small positive trial (the Grampian region early anistreplase trial) gets emphasised while another larger trial of the same issue (the European myocardial infarction project) remains unpublished. On a broader note, we would encourage a wider use of bayesian methods in reports of clinical trials, especially when a small trial is claiming a large treatment benefit.

STUART J POCOCK

Medical Statistics Unit,
London School of Hygiene and Tropical Medicine,
London WC1E 7HT

DAVID J SPIEGELHALTER

MRC Biostatistics Unit,
Institute of Public Health,
Cambridge CB2 2SR

- 1 GREAT Group. Feasibility, safety, and efficacy of domiciliary thrombolysis by general practitioners: Grampian region early anistreplase trial. *BMJ* 1992;305:548-53. (5 September.)
- 2 Pocock SJ, Hughes MD. Practical problems in interim analyses, with particular regard to estimation. *Controlled Clinical Trials* 1989;10:209-215.
- 3 Gruppo Italiano per lo Studio della Streptochinasi nell'infarto Miocardico (GISSI). Effectiveness of intravenous thrombolytic treatment in acute myocardial infarction. *Lancet* 1986;ii:397-401.
- 4 ISIS-2 (Second International Study of Infarct Survival) Collaborative Group. Randomised trial of intravenous streptokinase, oral aspirin, both, or neither among 17 187 cases of suspected myocardial infarction: ISIS-2. *Lancet* 1988;ii:349-60.

On site medical services at major incidents

EDITOR.—Matthew W Cooke¹ and D G Nancekivill² emphasise the need for better organisation and training for hospital staff in providing on site medical services when a major incident occurs.

A hospital coping with a deluge of casualties from a major incident might be overstretched in providing one or more appropriate teams as well as a doctor senior enough to be the medical incident officer (the Department of Health has abandoned the term site medical officer). Cooke highlights the paucity of training in this role. Wide ranging discussions have taken place in London with representatives of the London accident and emergency consultants' group, the London Ambulance Service, the British Association for Immediate Care, and health emergency planning officers from each Thames regional health authority with the aim of creating a cadre of 40-50 trained and accredited medical incident officers. This scheme relieves the main receiving hospital of the onerous duty of providing all the resources required at the site. The scheme has been approved by all participants, but, in view of its variation from guidance from the Department of Health, individual units will retain the option of making their own arrangements.

Two established training courses for doctors are available nationally. A one day course is run by the British Association for Immediate Care each year in Cambridge, and a three day course on the medical management of major incidents is run jointly by the Royal Postgraduate Medical School and the British Association for Immediate Care at Hammersmith Hospital. This course is multi-disciplinary and combines lectures, seminars, and practical training for NHS staff called on to work as medical incident officers or with mobile medical and nursing teams. In the two years that the course has been run, 102 people have been trained. The participants undertake an assessment at the end of the course, a major function of which is to allow the course organisers to assess the effectiveness of the training offered in key principles.

Though advanced trauma life support courses offer excellent training in clinical aspects, specific training is required for all prehospital care, including elements of safety and working with the emergency services.

ROBERT A COCKS

Royal Postgraduate Medical School,
Hammersmith Hospital,
London W12 0HS

- 1 Cooke MW. Arrangements for on scene medical care at major incidents. *BMJ* 1992;305:748. (26 September.)
- 2 Nancekivill DG. On site medical services at major incidents. *BMJ* 1992;305:726-7. (26 September.)
- 3 Department of Health. *Emergency planning in the NHS: health service arrangements for dealing with major incidents*. London: HMSO, 1990. (HC/90/25.)

EDITOR.—We agree with D G Nancekivill that both medical incident officers and site medical teams for major incidents need to be trained and to be familiar with the procedures of the other emergency services.¹ We disagree that training is a problem. The British Association for Immediate Care has been training doctors in this work for many years.

The association produced its first guide to managing major incidents in 1985,² and the skills of doctors trained by the association were recognised in the report on the railway accident at Clapham.³ The association's interservice and disaster liaison committee has been working with the ambulance, police, and fire services and the armed forces, coastguard, mountain rescue services, and, latterly, the Home Office adviser on civil emergencies on all aspects of managing major incidents. The association hopes to produce guidelines on the medical aspects of managing major