that heterosexual transmission of HIV is understood and that manifestations of AIDS and HIV infection that are particular to women are recognised—notably, disease of the female genital tract. But the overwhelming evidence shows that most manifestations of HIV infection are gender non-specific in type and incidence.

Ford is unfair and, indeed, incorrect in citing the recent clinical teaching text by my colleagues Parkin and Peters.² Their case oriented book, based on actual patients, uses a problem solving approach to help clinicians learn about AIDS. Most cases of AIDS in the developed world have been in men, and so real patients are described as man/he/him; it would be offensive to those men gratuitously to change their gender. Sadly, as heterosexual spread of HIV and ignorance and denial of it continue more cases will be seen among women, but the delay between infection and disease means that this spread will not be apparent as clinical disease for some years.

Current cumulative figures on AIDS in the United Kingdom show 5215 cases in men and 341 in women. In the vast majority of case histories, radiographs, and histological material, and most clinical slides, however, gender would be indistinguishable. Furthermore, Ford says that only two of 153 presentations in this book are of women, but, in fact, of 138 cases, 10 are in women and for 27 the gender is not specified; of 63 clinical photographs, six are of women and one shows a family.

Parkin was a coauthor of a report on one of the first women in the United Kingdom (a monogamous English housewife) to acquire AIDS through heterosexual transmission.¹ Parkin and other members of my team have repeatedly emphasised the importance of heterosexual spread over many years and have learnt much from patients; we have endeavoured to pass this on to our colleagues and the public in many forums. Ford's attack on young and enthusiastic colleagues is unwarranted.

ANTHONY J PINCHING

Department of Immunology,
Medical College of St Bartholomew's Hospital,
London EC1A 7BE

- 1 Ford C. Heterosexual AIDS epidemic. *BMJ* 1992;305:772. (26 September.)
2 Parkin JM, Peters BS. *Differential diagnosis in AIDS*. London: Wolfe, 1991.
3 Jenkins P, Malthouse SR, Baughan CA, Parkin JM, Pinching AJ. AIDS: the African connection. *BMJ* 1985;290:1284-5.

EDITOR,—We agree with Chris Ford that services for diagnosing, treating, and preventing HIV infection in women have lagged behind those targeted at people whose major risk behaviour is male homosexual contact.¹ The pattern of risk behaviour has changed in recent years; of the 263 HIV seropositive heterosexuals who attended St Stephen's Clinic to the end of 1991, 110 were female and 49 of these presented after January 1990. In response to this change a clinic for women only was initiated in April 1991 to attract women with a range of risk behaviours who were not using existing facilities. This clinic is staffed exclusively by women for women and has proved to be popular.² Similarly, a clinic providing genitourinary medicine services for lesbians has been instituted in the district as the sexual health of this group is underresearched and lesbians have had difficulty in gaining access to appropriate medical services.

Ford also cites a short article by one of us as an example in which "information about the disease is still male dominated."³ This article was intentionally written without specific references to either the sex or orientation of the infected subjects and to apply equally to all with HIV infection or AIDS. We hope that HIV infection will be regarded as a multi-system disorder that may affect people from any walk of life but is amenable to risk reduction—

the adoption of measures such as safer sex and not sharing drug using equipment.

RAK NANDWANI
JAYNE KAVANAGH
SIMON MANSFIELD

St Stephen's Clinic,
London SW10 9TH

- 1 Ford C. Heterosexual AIDS epidemic. *BMJ* 1992;305:772. (26 September.)
2 McCarthy G, Cockell A, Beevor A, Kell P, Boag F. An audit of a women only clinic for HIV, genitourinary medicine and substance misuse. *Gentourin Med* (in press).
3 Mansfield S. How should fungal infections be treated in HIV patients? *MIMS Magazine* 1992 June 30:48.

Tachycardia during cisapride treatment

EDITOR,—Sten Olsson and I Ralph Edwards report seven cases of tachycardia or palpitations associated with treatment with cisapride.¹ These presumably represent all or most of these events reported spontaneously to four national centres over three years.

We have studied cisapride in 13 233 patients and have compared the rates of palpitations, tachycardia, and extrasystoles with data collected under substantially identical conditions from 240 381 patients in 23 recent prescription event monitoring studies. This database, which comprises about 40% of our records, includes 1960 records of tachycardia, palpitations, or extrasystoles, 595 of which were noted to have occurred during the first month of observation (table).

Number of cases of palpitations, tachycardia, and extrasystoles recorded during first month* in 23 prescription event monitoring studies: rates per thousand in descending order

Drug	No	Rate/1000
Nicardipine	160	14.6
Lisinopril	99	8.0
Xamoterol	35	6.5
Enalapril	72	4.7
Fluvoxamine	44	4.0
Buspirone	43	3.9
Diltiazem	25	2.5
Fluoxetine	29	2.3
Betaxolol	2	1.3
Ornepazole	20	1.2
Terodiline	14	1.1
Cisapride	11	0.8
Nedocromil	10	0.6
Flunitrazepam	6	0.8
Misoprostol	9	0.7
Tenoxicam	4	0.4
Etozolac	3	0.3
Famotidine	3	0.3
Nizatidine	2	0.3
Nabumetone	2	0.2
Acrivastine	1	0.1
Cetirizine	1	0.1
Loratadine	0	0
All studies	595	2.5

*Includes patients who had stopped treatment with relevant drug.

The rate of these events during the first month was 0.8/1000 for cisapride compared with a mean rate of 2.5/1000 in all the studies. Even allowing for the fact that several studies were of drugs used to treat cardiovascular conditions in which tachycardia would be expected to be common, our results provided no epidemiological support for Olsson and Edwards's hypothesis. We therefore examined all relevant records in our study of cisapride individually. There were 104 such records—67 of palpitations, 23 of tachycardia, five of extrasystoles, and nine of heart block. Fifty five of the events occurred during the follow up period, up to a year after the patient had stopped using cisapride. Only eight patients using cisapride experienced one of these events during the first

month of treatment. We consider that all these events were coincidental and almost certainly unrelated to cisapride. For all 104 patients, no doctor stopped treatment with cisapride because of these events or attributed them to the drug.

The few reports to the World Health Organisation's programme suggest an extremely low risk, even if there has been considerable underreporting. We believe that these reports represent only a small fraction of the events of this type that would be expected with or without treatment with cisapride. Palpitations and tachycardia are common and have occurred coincidentally in all of the groups we have studied. Spontaneous reporting systems were designed to detect previously unsuspected reactions that were too infrequent to be identified during premarketing clinical trials. When events are reported that commonly occur in the absence of any recognisable association with treatment it is inappropriate to draw conclusions from them, especially if alternative causes have not been eliminated.

WILLIAM INMAN
KIYOSHI KUBOTA

Drug Safety Research Unit,
Southampton SO3 8BA

- 1 Olsson S, Edwards IR. Tachycardia during cisapride treatment. *BMJ* 1992;305:748-9. (26 September.)

EDITOR,—Sten Olsson and I Ralph Edwards report that treatment with cisapride may induce tachycardia.¹ They conclude that the mechanism of this action is not obvious but may be a procainamide-like effect. A tachycardic response to cisapride has, however, been reported previously,² and we suggest that activation of serotonin-4 receptors is a more likely explanation for this effect.

The existence of serotonin-4 receptors has only recently been established.¹ These receptors are positively coupled to adenylyl cyclase¹ and are present on pig heart,⁵ where they mediate a chronotropic response. Experiments have also shown the presence of serotonin-4 receptors on human right atrium, where they mediate an inotropic response.⁶ In this atrial tissue the response to serotonin was associated with raised levels of cyclic AMP⁷; thus, although the rate response was not measured, it is likely that, as in pig heart, stimulation of serotonin-4 receptors mediates a chronotropic as well as an inotropic response. Furthermore, it is well documented that cisapride is an agonist at serotonin-4 receptors,¹ and, in particular, exhibits partial agonist activity at the serotonin-4 receptor found in human right atrium.⁸

We conclude that the tachycardic response to cisapride in humans may be due to activation of atrial serotonin-4 receptors. Possibly this occurs infrequently in clinical practice because cisapride is a particularly weak partial agonist at serotonin-4 receptors in the human heart.⁹

P P A HUMPHREY

Glaxo Institute of Applied Pharmacology,
Department of Pharmacology,
University of Cambridge,
Cambridge CB2 1QJ

K T BUNCE

Glaxo Group Research Limited,
Gastrointestinal Pharmacology Department,
Ware,
Hertfordshire SG12 0DP

- 1 Olsson S, Edwards IR. Tachycardia during cisapride treatment. *BMJ* 1992;305:748-9. (26 September.)
2 Bateman DN. The action of cisapride on gastric emptying and the pharmacodynamics and pharmacokinetics of oral diazepam. *Eur J Clin Pharmacol* 1986;30:205-8.
3 Bockaert J, Fozard JR, Dumuis A, Clarke DE. The 5-HT₄ receptor: a place in the sun. *Trends Pharmacol Sci* 1992;13:141-5.
4 Dumuis A, Sebben M, Bockaert J. The gastrointestinal prokinetic benzamide derivatives are agonists at the non-classical 5-HT receptor (5-HT₄) positively coupled to adenylyl cyclase in neurons. *Naunyn Schmiedeberg Arch Pharmacol* 1989;340:403-10.
5 Villalon CM, den Boer MO, Heiligers PC, Saxena PR. Further