Published in final edited form as:

Biochem Biophys Res Commun. 2007 April 6; 355(2): 444–450.

Cardiac-Specific Overexpression of *SCN5A* Gene Leads to Shorter P Wave Duration and PR Interval in Transgenic Mice

Teng Zhang, Sandro L. Yong, Xiao-Li Tian, and Qing K. Wang*

Department of Molecular Cardiology, Lerner Research Institute and Center for Cardiovascular Genetics, Department of Cardiovascular Medicine, Cleveland Clinic, 9500 Euclid Ave, Cleveland, OH 44195, USA Department of Molecular Medicine, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, 9500 Euclid Ave, Cleveland, OH 44195, USA

Abstract

SCN5A plays a critical role in cardiac electrophysiology and its mutations, either gain- or loss-of-functions, are associated with lethal arrhythmias. In this study we investigated the effect of overexpression of *SCN5A* on the cardiac phenotype in a transgenic mouse model (TG-WT L10). Compared to NTG mice, heart rate, QRS duration and QT intervals remained unchanged in TG-WT mice. Moreover, no spontaneous ventricular arrhythmias were detected in TG-WT hearts. Despite these results, a mild, irregular cardiac phenotype was observed in TG-WT mice. The P wave and PR interval were significantly shorter in TG-WT compared with NTG mice (P, 8.8 ± 0.8 ms vs. 12.6 ± 0.9 ms; PR, 12.5 ± 2 ms vs. 33.5 ± 0.7 ms). Furthermore, spontaneous premature atrial contractions were often detected in TG-WT mice. These results suggest that the expression level of the *SCN5A* gene is a determinant for the length of the P wave duration and PR interval on electrocardiograms (ECG).

Keywords

cardiac sodium channel gene *SCN5A*; long QT syndrome (LQTS); ventricular arrhythmias or tachycardia (VT); transgenic mice; electrocardiogram (ECG); P wave and PR interval; sodium current

Introduction

The cardiac voltage-sensitive sodium channel $\text{Na}_v1.5$ is responsible for rapid influx of sodium ions that initiates and propagates the cardiac action potential in the heart [1-6]. The large peak sodium influx is responsible for excitability and conduction in the myocardium and special conduction tissues. It is encoded by the *SCN5A* gene which consists of 28 exons spanning over 80 kb of DNA on chromosome 3p21 [7]. To date, hundreds of mutations have been identified in *SCN5A* and linked to a number of heart diseases including long QT syndrome (LQTS), Brugada syndrome (BrS), cardiac conduction disease, sudden infant death, idiopathic ventricular fibrillation, and congenital sick sinus syndrome [1;4;5;8;9]. Three *SCN5A* mutations have been studied *in vivo* in transgenic mice, including transgenic overexpression of the N1325S mutation associated with LQTS in the heart (TG-NS/LQT3) [3], knockin of the ΔKPQ associated with LQTS [10], and knockin of the 1795insD mutation associated with

*Corresponding author. Fax: +1 216 636 1231. E-mail address: wangq2@ccf.org (Q.K. Wang).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

bradycardia, conduction disease, LQTS and BrS [11]. These mice recapitulate the major clinical features that correspond to human patients.

Knockout mice for *SCN5A* have also been generated. Homozygous mice died at embryonic day 10.5 [12]. Heterozygous mice showed prolongation of the P wave and the PR interval on the ECG, which was indicative of slow conduction [13;14]. However, the effect of overexpression of the *SCN5A* gene on the electrophysiology of the heart has not been explored. To further investigate the physiological contribution of $\text{Na}_v1.5$ to cardiac function, we created a transgenic mouse model with overexpression of wild type human *SCN5A* gene in the mouse heart (TG-WT). The phenotypes of the transgenic mice were compared to non-transgenic littermate control mice (NTG). We demonstrated that the P wave duration and PR interval on the ECG in TG-WT mice were significantly shorter than in NTG mice. TG-WT mice also showed a high frequency of spontaneous premature atrial contractions (PACs). No other cardiac abnormalities were identified. Our results provide the first assessment of the effects of increased $\text{Na}_v1.5$ expression on cardiac function *in vivo*.

Material and methods

Generation of transgenic mice and genotyping

The transgenic mice with selective cardiac expression of human wild type *SCN5A* gene was generated by placing *SCN5A* under the control of the mouse myosin heavy chain- α promoter (TG-WT). The transgenic construct was similar to what we used to generate TG-NS/LQT3 L3 and L12 except that the wild type human *SCN5A* cDNA was used [3;15]. Quantitative Southern blot analysis was performed to estimate the copy number of the transgene [3;15]. The mice were genotyped by multiplex PCR which simultaneously amplified the mouse endogenous gene *AGGF1* (Pe10: 5'-GAGCTCACCTCCGCCTCGAT-3' and Pe11: 5'-CCTCCTTACTTAGTGCTGGAC-3': 784 bp) and human *SCN5A* transgene (Pmcn-3nd: 5'-CCTCTGACAGAGAAGCAGGCACTTT-3 and Pscn5a-5nd: 5'-CTGAACCGGAAGATGGTC TTGCCTTT-3', 523 bp) [3;15;16]. The amplification of mouse *AGGF1* served as an indicator for successful PCR to avoid mis-genotyping.

Quantification of the transcript of human *SCN5A* transgene in TG-WT mouse hearts

Transcript of human *SCN5A* transgene was quantified by quantitative reverse-transcript PCR (RT-PCR) [17]. RT-PCR primers were 5'-CATGCTCATCATGTGCACCAT-3' and 5'-GGTGCGTAAGGCTGAGACATT-3'. This pair of primers amplified a 259 bp PCR fragment from both the mouse and human *SCN5A* genes. The internal loading control was 18S rRNA with amplicon size of 274 bp by primers 5'-CGTCTGCCCTATCAACTTT-3 and 5'-CGGCTGCTGGCACCAGACTT-3 as described previously. In brief, 5 μg of total RNA from a cardiac ventricular tissue sample was pre-treated with 5 μg of DNase I to remove contaminated genomic DNA (37°C for 30 min), followed by 90°C for 5 min to inactivate DNase I. No genomic DNA (transgene) contamination in the pre-treated RNA was confirmed by PCR without RT. Quantitative PCR was performed using cDNA template reverse transcribed with random hexamer. Various numbers of PCR cycles were empirically determined to ensure that PCR amplification was in the logarithm range (18-25 cycles for 18S rRNA; 25-31 cycles for *SCN5A*). The relative amount of *SCN5A* transcripts was expressed as a ratio of band densities of *SCN5A* to 18S multiplied by 1000.

Protein expression of human *SCN5A* transgene in TG-WT mouse hearts

Western blot analysis was carried out to estimate the protein expression of human *SCN5A* transgene in TG-WT mouse hearts. Total protein was extracted from the isolated hearts using lysis buffer containing 1 M Tris-HCl (pH 8.0), 5 M NaCl, and 0.1 M EDTA and 10% NP-40.

The protein concentration was measured using the BioRad protein assay kit, BSA was used as a standard, and all samples were normalized to the same concentration of 5 or 10 µg/ul. 30-75 µg or protein extracts was separated by SDS-PAGE gels, and transferred to polyvinylidene difluoride membranes (Immun-Blot PVDF Membrane, BioRad). The membrane was blocked in 5% nonfat dry milk in PBST (phosphate-buffered saline (PBS), 0.05% (v/v) Tween-20 (Sigma), pH 7.4), and then probed with an anti-human Na_v1.5 antibody [6], followed by a horseradish peroxidase-conjugated secondary antibody. The protein signal was visualized using the ECL Western Blotting Detection kit and Hyperfilm ECL (Amersham Biosciences). The member was re-probed using an anti-GAPDH antibody (Chemicon), which serves as loading control.

Electrophysiology

All experiments were conducted in accordance with the guidelines of the Cleveland Clinic Foundation Institutional Review Board on Animal Subjects and conformed to NIH guidelines. Details that describe the isolation procedure have been previously reported [18;19]. Hearts were harvested from age-matched (6-8 months) mice and enzymatically digested via the retrograde, Langendorff perfusion manner. Enzymatic digestion (Type II, 209 U/mg, Worthington) was conducted using a Ca²⁺-free Tyrode's buffer solution for 28 min. Ventricular and atrial appendages were separately removed and placed in separate incubation buffers (final Ca²⁺ concentration of 1.8 mM). Gentle titration of each was performed and cells were collected by gravity sedimentation. Prior to experimentation, cells were re-suspended in recording bath solution.

Recordings were conducted using a MultiClamp 700A amplifier interfaced to a Pentium computer equipped with Digidata 1322A and pClamp 9 software (Axon Instruments) [3;19]. Only single, quiescent, and rod-shaped myocytes were chosen. Recording electrodes were fabricated from borosilicate glass (FHC) with 1.8-2.3 MΩ resistance. Signals were low-passed filtered at 1 kHz and digitized at 10 kHz and current recordings were not corrected for leak. All recordings were made at room temperature and the recording bath and pipette solutions have been previously reported [3;19].

Telemetric electrocardiographic recording

ECG from non-transgenic and TG-WT L10 mice were recorded and observed from freely moving mice by the Telemetry system (DSI international) as described [3;18]. Male transgenic mice and age-matched littermates were used for this study (6-8 months, 6-8 from each genotype).

Langendorff perfusion of the mouse heart

The mouse was heparinized with 0.15 ml heparin (1000 U/ml) and anesthetized with 0.65 mg/g, i.p. avertin (20 mg/ml, ALDRICH) before sacrifice. The heart was then quickly excised, cannulated, and studied as described previously [20].

Statistical analysis

All values are expressed as mean ± SEM. The paired t-test was used to evaluate the ECG parameters and electrophysiological recording data. ANOVA was used to compare the expression of *SCN5A* and heart rate. $P < 0.05$ was set for statistical significance.

Results

Development of transgenic mice with cardiac overexpression of wild type human *SCN5A*

Two transgenic lines with overexpression of the wild type human *SCN5A* gene were developed and designated as TG-WT L5 and L10, respectively (Fig. 1A and 1B). By quantitative Southern blot analysis, we estimated that TG-WT L5 had 1-2 copies of transgene and L10 had approximately 10 copies of transgene (Fig. 1B). Two previously generated mouse models with overexpression of the LQTS-causing mutation N1325S in *SCN5A* were included in the present study as controls. One model is the transgenic line with 1-2 copies of transgene (TG-NS/LQT3 L3) and the other is the line with 10 copies of transgene (TG-NS/LQT3 L12). The latter has been characterized in detail and found to display characteristic clinical features of LQTS, including prolonged QTc and development of spontaneous VT/VF, as well as syncope and sudden death [3;19;20]. As shown in Fig. 1B, TG-WT L5 had a comparable copy number of transgene with TG-NS/LQT3 L3, and TG-WT L10 had a similar copy number to TG-NS/LQT3 L12. As the major goal of this study is to determine the effect of overexpression of *SCN5A* on cardiac function, the results in the present study focused only on TG-WT L10 mice, the line with the higher copy number of the transgene. The amount of the total *SCN5A* transcript in ventricular tissues of TG-WT L10 and TG-LQT3 L12 was comparable (1.3 ± 0.045 vs 1.1 ± 0.033 , $p > 0.05$), and both were significantly increased compared to NTG hearts (0.15 ± 0.003 , $p < 0.05$) as shown by quantitative RT-PCR (Fig. 1C). The protein expression level of $\text{Na}_v1.5$ in both TG-WT L10 and TG-LQT3 L12 hearts was also markedly increased compared to that in NTG mice as shown by Western blot analysis (Fig. 1D). These results indicate that the expression of the *SCN5A* gene is markedly increased in TG-WT L10 mice at both the mRNA and protein levels.

Phenotype of TG-WT mice

Mouse ECG were recorded from conscious and free-moving TG-WT L10 and NTG mice by a Telemetry monitoring system (5 mice per group). Representative ECG traces are shown in Fig. 2A for NTG mice and in Fig. 2B for TG-WT L10 mice. The major parameters on ECG were measured and shown in Fig. 2C-G. The TG-WT L10 mice showed a shorter PR interval than NTG mice (Fig. 2D). The P wave duration in the TG-WT mice was also shorter than that in the NTG mice (Fig. 2E). No differences in the RR interval (heart rate), QT interval corrected for the heart rate (QTc), and the QRS interval were detected between the TG-WT L10 and NTG mice (Fig. 2C, 2F, 2G).

Recently, we reported spontaneous ventricular arrhythmias (VT) in telemetrically-monitored TG-NS/LQT3 L12 mice [3] and in isolated Langendorff-perfused hearts from this line of mice [20]. However, the Telemetry system did not detect VT in the TG-WT L10 mice and NTG mice. No VT were detected in isolated Langendorff-perfused hearts either. Similar to the NTG mice, no sudden death was observed in TG-WT L10 mice. However, frequent spontaneous premature atrial contractions (PAC) were detected in TG-WT L10 mice. In contrast, no PACs were observed in NTG mice (Fig. 3). It is interesting to note that generation of PACs was associated with successive reduction of the PR interval with each beat close to the PAC (Fig. 3A).

Electrophysiology of isolated cardiomyocytes from TG-WT mice

Sodium current (I_{Na}) were elicited with a 60 ms ramp protocol from a holding potential of -80 to 0 mV (Fig. 3). Peak I_{Na} was expressed as pA/pF in which the cell capacitance was measured by calculating the area under the capacitive transient elicited by a -10 mV pulse. Sodium current densities for TG-WT and NTG ventricular cells were 14.5 ± 1.2 ($n = 19$) and 16.6 ± 2.3 pA/pF ($n = 10$; $P > 0.05$), respectively, and for atrial cells the densities were 6.4 ± 1.1 ($n =$

6) and 7.4 ± 0.8 pA/pF ($n = 8$; $P > 0.05$), respectively. For ventricular cells, $I_{Na,L}$ was estimated quantitatively by fitting the declining phase of I_{Na} with a biexponential function, $I/I_{max} = A_{1x}(1 - e^{-(t/\tau_1)}) + A_{2x}(1 - e^{-(t/\tau_2)})$ and plotting the slow time component, τ_2 ; for TG-WT and NTG cells the values were 1.8 ± 0.1 and 1.8 ± 0.3 ms, $P > 0.05$. Action potentials for both ventricular and atrial cells were elicited with a 1000-1500 pA pulse of 25-50 ms duration at a cycle length of 0.5 sec. Zero holding current was applied. The time to 90% repolarization (APD_{90}) and cell capacitance (see Fig. 3) were measured. APD_{90} values for TG-WT and NTG ventricular cells were 68.4 ± 5.0 ($n = 9$) and 69.5 ± 5.3 ms ($n = 11$, $P > 0.05$), respectively, and for atrial cells the values were 27.4 ± 3.2 ($n = 6$) and 27.5 ± 1.7 ms ($n = 6$, $P > 0.05$), respectively.

Discussion

Both the knockout and transgenic expression techniques are useful systems for exploring the physiological functions of a gene. Previous studies using knockout mice revealed that reduction of *SCN5A* expression by 50% resulted in prolongation of the PR interval [13;14]. In this study, we created transgenic TG-WT L10 mice that overexpress the human cardiac sodium channel gene *SCN5A*. Baseline ECG recordings by a Telemetry system revealed a shortened PR interval and P wave duration in TG-WT L10 mice compared to their littermate controls. As the AV node forms much of the PR segment on the ECG, these results suggest that the expression level of the *SCN5A* gene may alter AV nodal conduction and, as such, decrease the PR interval on the ECG.

Despite these unexpected findings on the P wave and PR interval, we observed no abnormalities in the electrophysiology of the ventricles. The QRS duration and QTc were normal in TG-WT and comparable to NTG mice. More importantly, no ventricular arrhythmias were detected in TG-WT mice. The sodium current densities and APDs from TG-WT ventricular cardiomyocytes were nearly identical to that from NTG cells. Equally important were the similarities in sodium current densities and APDs found in atrial cells. This lends support to our earlier finding that overexpression of *SCN5A* did not significantly increase the surface density of the sodium channels in ventricular [3] and atrial myocytes. Thus, it would appear that the risks associated with overexpression of *SCN5A* is minimal and the abnormalities that were previously reported in TGNS/LQT3 L12 mice [3] were directly associated to the N1325S mutation.

Similar to TG-WT L10 mice, TG-NS/LQT3 L12 mice also exhibited the short PR interval phenomenon [3]. This suggests that PR shortening may be a consequence of the overexpression of *SCN5A* alone and, not necessarily the direct effect of the N1325S mutation. Furthermore, the high incidence of arrhythmias, mortalities, and electrophysiological abnormalities that have been reported in TG-NS/LQT3 L12 mice, whereas no such cardiac abnormalities have been detected in TG-WT L10 mice, is further evidence that the cardiac dysfunctions are a direct consequence of the N1325S mutation and not from *SCN5A* gene overexpression. Such phenotypical comparison between TG-WT L10 and TG-NS/LQT3 L12 strongly validates our TG-NS/LQT3 mice as a suitable model for LQT3.

Overexpression of mutation R302Q in the *PRKAG2* gene in the mouse heart also exhibited a short PR interval [21]. In addition, these mice showed an increased QRS duration and an increased incidence of supraventricular arrhythmias, the features of which are associated with Wolff-Parkinson-White syndrome (WPW) [21]. The *PRKAG2* gene encodes the γ -2 subunit of AMP-activated protein kinase. Although the transgenic model serves as a suitable model for WPW, no clear molecular mechanism by which overexpression of R302Q causes PR interval shortening and/or induction of supraventricular arrhythmias was provided. Considering the similarities in the observations for PR intervals between our TG-WT L10 and

the *PRKAG2* mice, it is likely that the *PRKAG2* defect may affect the function and/or expression of the *SCN5A* gene, resulting in WPW.

The molecular mechanism for the P-wave and PR interval shortening in TG-WT L10 mice has not been clearly defined within the scope of this study. We speculate that expression of the cardiac sodium channel in the conductive tissues (i.e., SA node and AV node) may be increased such that conduction of electrical pulses in these tissues is facilitated. Such a scenario would lead to accelerations in atrial depolarization and conduction, resulting in a shorter P wave duration and PR interval. The same mechanisms may also explain the development of PACs in TG-WT L10 mice which presumably originate near or at the AV node as AV junctional rhythms. Historically and clinically, individuals with short PR intervals and PACs range from asymptomatic to a variety of complex arrhythmias, and our TG-WT L10 mice are at the lower risk range.

In conclusion, cardiac overexpression of the cardiac sodium channel gene *SCN5A* causes shortening of the P wave duration and PR interval.

Acknowledgments

This work was supported by an NIH grant R01 HL66251 (to Q.K.W.), The State of Ohio Wright Center of Innovation grant and Biomedical Research and Technology Transfer Partnership Award (Ohio's Third Frontier Project) (to Q.K.W.) and AHA-Ohio-Affiliate Postdoctoral Fellowship (to S.Y.). Q.K.W. is an Established Investigator of the American Heart Association (0440157N). We thank Yuanna Cheng and Mei-Ling Chnag Liao for help with the equipment for recording optical action potentials, and Qianqian Liu for technical help. We also thank Ying Ni, Shin Yoo and other members of Wang Laboratory and David van Wagoner for valuable discussion.

References

1. Chen Q, Kirsch GE, Zhang D, Brugada R, Brugada J, Brugada P, Potenza D, Moya A, Borggrefe M, Breithardt G, Ortiz-Lopez R, Wang Z, Antzelevitch C, O'Brien RE, Schulze-Bahr E, Keating MT, Towbin JA, Wang Q. Genetic basis and molecular mechanism for idiopathic ventricular fibrillation. *Nature* 1998;392:293–296. [PubMed: 9521325]
2. Gellens ME, George AL Jr, Chen LQ, Chahine M, Horn R, Barchi RL, Kallen RG. Primary structure and functional expression of the human cardiac tetrodotoxin-insensitive voltage-dependent sodium channel. *Proc.Natl.Acad.Sci.U.S.A* 1992;89:554–558. [PubMed: 1309946]
3. Tian XL, Yong SL, Wan X, Wu L, Chung MK, Tchou PJ, Rosenbaum DS, Van Wagoner DR, Kirsch GE, Wang Q. Mechanisms by which *SCN5A* mutation N(1325)S causes cardiac arrhythmias and sudden death in vivo. *Cardiovasc.Res* 2004;61:256–267. [PubMed: 14736542]
4. Wang Q, Shen J, Li Z, Timothy K, Vincent GM, Priori SG, Schwartz PJ, Keating MT. Cardiac sodium channel mutations in patients with long QT syndrome, an inherited cardiac arrhythmia. *Hum.Mol.Genet* 1995;4:1603–1607. [PubMed: 8541846]
5. Wang Q, Shen J, Splawski I, Atkinson D, Li Z, Robinson JL, Moss AJ, Towbin JA, Keating MT. *SCN5A* mutations associated with an inherited cardiac arrhythmia, long QT syndrome. *Cell* 1995;80:805–811. [PubMed: 7889574]
6. Wu L, Nishiyama K, Hollyfield JG, Wang Q. Localization of $Na_v1.5$ sodium channel protein in the mouse brain. *NeuroReport* 2002;13:2547–2551. [PubMed: 12499865]
7. Wang Q, Li Z, Shen J, Keating MT. Genomic organization of the human *SCN5A* gene encoding the cardiac sodium channel. *Genomics* 1996;34:9–16. [PubMed: 8661019]
8. Tan HL, Bink-Boelkens MTE, Bezzina CR, Viswanathan PC, Beaufort-Krol GCM, van Tintelen PJ, van den Berg MP, Wilde AAM, Balser JR. A sodium-channel mutation causes isolated cardiac conduction disease. *Nature* 2001;409:1047. [PubMed: 11234014]
9. Wolf CM, Berul CI. Inherited conduction system abnormalities--one group of diseases, many genes. *J.Cardiovasc.Electrophysiol* 2006;17:446–455. [PubMed: 16643374]
10. Nuyens D, Stengl M, Dugarmaa S, Rossenbacker T, Compennolle V, Rudy Y, Smits JF, Flameng W, Clancy CE, Moons L, Vos MA, Dewerchin M, Benndorf K, Collen D, Carmeliet E, Carmeliet P.

Abrupt rate accelerations or premature beats cause life-threatening arrhythmias in mice with long-QT3 syndrome. *Nat.Med* 2001;7:1021–1027. [PubMed: 11533705]

11. Remme CA, Verkerk AO, Nuyens D, van Ginneken AC, van BS, Belterman CN, Wilders R, van Roon MA, Tan HL, Wilde AA, Carmeliet P, de Bakker JM, Veldkamp MW, Bezzina CR. Overlap syndrome of cardiac sodium channel disease in mice carrying the equivalent mutation of human SCN5A-1795insD. *Circulation* 2006;114:2584–2594. [PubMed: 17145985]
12. Papadatos GA, Wallerstein PM, Head CE, Ratcliff R, Brady PA, Benndorf K, Saumarez RC, Trezise AE, Huang CL, Vandenberg JI, Colledge WH, Grace AA. Slowed conduction and ventricular tachycardia after targeted disruption of the cardiac sodium channel gene *Scn5a*. *Proc.Natl.Acad.Sci.U.S.A* 2002;99:6210–6215. [PubMed: 11972032]
13. Royer A, van Veen TA, Le BS, Marionneau C, Griol-Charhbil V, Leoni AL, Steenman M, van Rijen HV, Demolombe S, Goddard CA, Richer C, Escoubet B, Jarry-Guichard T, Colledge WH, Gros D, de Bakker JM, Grace AA, Escande D, Charpentier F. Mouse model of SCN5A-linked hereditary Lenegre's disease: age-related conduction slowing and myocardial fibrosis. *Circulation* 2005;111:1738–1746. [PubMed: 15809371]
14. van Veen TA, Stein M, Royer A, Le QK, Charpentier F, Colledge WH, Huang CL, Wilders R, Grace AA, Escande D, de Bakker JM, van Rijen HV. Impaired impulse propagation in *Scn5a*-knockout mice: combined contribution of excitability, connexin expression, and tissue architecture in relation to aging. *Circulation* 2005;112:1927–1935. [PubMed: 16172272]
15. Tian XL, Wang QK. Generation of transgenic mice for cardiovascular research. *Methods Mol Med* 2006;129:69–81. [PubMed: 17085805]
16. Tian XL, Kadaba R, You SA, Liu M, Timur AA, Yang L, Chen Q, Szafranski P, Rao S, Wu L, Housman DE, DiCorleto PE, Driscoll DJ, Borrow J, Wang Q. Identification of an angiogenic factor that when mutated causes susceptibility to Klippel-Trenaunay syndrome. *Nature* 2004;427:640–645. [PubMed: 14961121]
17. You SA, Archacki SR, Angheloiu G, Moravec CS, Rao S, Kinter M, Topol EJ, Wang Q. Proteomic approach to coronary atherosclerosis shows ferritin light chain as a significant marker: evidence consistent with iron hypothesis in atherosclerosis. *Physiol Genomics* 2003;13:25–30. [PubMed: 12644631]
18. Yong SL, Wang QK. Animal models for cardiac arrhythmias. *Methods Mol Med* 2006;129:127–148. [PubMed: 17085809]
19. Yong SL, Ni Y, Zhang T, Tester DJ, Ackerman MJ, Wang QK. Characterization of the cardiac sodium channel SCN5A mutation, N(1325)S, in single murine ventricular myocytes. *Biochem.Biophys.Res.Commun* 2007;352:378–383. [PubMed: 17118339]
20. Tian XL, Cheng Y, Zhang T, Liao ML, Yong SL, Wang QK. Optical mapping of ventricular arrhythmias in LQTS mice with SCN5A mutation N1325S. *Biochem.Biophys.Res.Commun* 2007;352:879–883. [PubMed: 17157817]
21. Sidhu JS, Rajawat YS, Rami TG, Gollob MH, Wang Z, Yuan R, Marian AJ, DeMayo FJ, Weilbacher D, Taffet GE, Davies JK, Carling D, Khoury DS, Roberts R. Transgenic mouse model of ventricular preexcitation and atrioventricular reentrant tachycardia induced by an AMP-activated protein kinase loss-of-function mutation responsible for Wolff-Parkinson-White syndrome. *Circulation* 2005;111:21–29. [PubMed: 15611370]

Fig 1.

Generation and characterization of the transgenic mouse model. (A) Transgenic construct for engineering of transgene, human wild type *SCN5A*, into the mouse genome. The *NotI* fragment consists of the mouse myosin heavy chain- α promoter (α -mMHCp), transgene *SCN5A*, and the human growth hormone poly(A) signal (*hGH pIA*). Probe (1.7 kb), a portion of α -mMHCp, was used as a marker for Southern blot analysis. PCR primer pairs P1-P2 and P3-P4 were used to identify transgenic mice during breeding. (B) Southern blot analysis. Mouse genomic DNA was digested with *BstEII* and hybridized with probe [see above]. The 6.0- and 3.1-kb fragments represent the transgene *SCN5A*, and the endogenous mouse myosin heavy chain- α gene, respectively. Two transgenic lines, TG-WT L5 and L10, were generated. The following genotypes are listed: *TGM(NS31)L3*: Lanes 1 and 3; *TGM(NS31)L12*: Lanes 4 and 6; and *non-transgenic mice*: Lanes 2 and 5. (C) Quantitative RT-PCR shows the amount of total *SCN5A* transcripts in ventricular tissues of TG-WT L10, TG-LQT3 L12 and NTG mice (D) Western blot analysis was performed using heart tissue protein extracts from TG-WT L10, TGM(NS31) L12 and NTG mice. A polyclonal anti-*SCN5A* antiserum was raised against a peptide located in the less-conserved N-terminus of *SCN5A*. The same filter was probed with a monoclonal anti-GAPDH antibody (Sigma-Aldrich) in order to calibrate the amount of protein extracts loaded in each lane.

Fig 2.

Representative ECG traces from NTG (A) and TG-WT mice (B). The various ECG parameters were measured and analyzed (C-G). The P wave duration and PR interval in TG-WT mice were significantly shorter than those of NTG mice ($n=5$, $P=0.017$ and 0.00086 , respectively). No differences were observed in RR, QTc, and QRS duration ($P>0.05$). NS, no statistical significance.

Fig 3.

Detection of premature atrial contractions (PACs) by a telemetry ECG recording system from TG-WT mice. (A) ECG traces showing PACs. A PAC is defined as an ectopic beat with abnormally shaped or absent P-wave before the QRS complex, but without any effect on QRS complexes. PACs are indicated by arrows. (B) The frequency of PACs in TG-WT and control NTG mice was analyzed from ECGs as the number of PACs per hour.

Fig 4.

Sodium currents and action potentials from atrial and ventricular myocytes. Sodium currents from TG-WT and NTG single atrial and ventricular myocytes were generated using the ramp step. Representative sodium current traces for each cell group are shown with the corresponding action potential tracing. Cell capacitance for ventricular cells was estimated as 224.0 ± 22.9 (TG-WT) and 220.2 ± 14.5 pF (NTG), $p > 0.05$, whereas for atrial cells the values were 194.9 ± 17.9 (TG-WT) and 144.9 ± 27.6 (NTG), $p < 0.05$. Sodium current densities and APD₉₀ values are listed in the Results section.