Corrigendum

Suppressor of cytokine signaling 3 controls lysosomal routing of G-CSF receptor

Mahban I Irandoust, Lambertus HJ Aarts, Onno Roovers, Judith Gits, Stefan J Erkeland and Ivo P Touw

The EMBO Journal (2007) **26**, 2822. doi:10.1038/sj.emboj.7601709

Correction to: *The EMBO Journal* (2007) **26**, 1782–1793. doi:10.1038/sj.emboj.7601640

The authors have identified an error in Figure 7B of the above article.

The error concerns lanes 3 (wt) and 4 (ev), where in the lower line + and – have been swapped.

Please find below the correct figure in its entirety.

The authors apologize for any inconvenience caused.

Figure 7 Ubiquitination of G-CSFR. (A) Phoenix E cells cotransfected with HA-tagged wt or K5R G-CSFR and FLAG-tagged Ub were incubated with (+) or without (–) G-CSF for 30 min, and HA immunoprecipitates were analyzed by Western blotting using anti FLAG (upper panel) or anti-HA antibodies (lower panel). (B) A similar setup as in (A), but instead of HA immunoprecipitation, cells were incubated with bio-G for 20 min at 16°C and for 30 min at 37°C and subsequently G-CSFR was pulled down with streptavidin coated magnetic beads.