We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer. Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.

LETTERS

PHOTOS.COM

PRESUMED CONSENT

First address informed consent in organ donation

English's arguments in favour of presumed consent for organ donation ignore serious problems with the current system of organ transplantation.¹ People who sign organ donation cards claiming that they will be donating organs "after my death" may not realise that organ donors declared dead by brain based criteria are not dead in the usual sense. Their hearts are beating, their bodies are warm, and they do not seem to be dead. Family members may be understandably reluctant to sign a form authorising what they believe to be the killing of their loved one by removal of vital organs.

In addition, there has been widespread questioning of both the philosophical and scientific viability of brain death criteria.^{2 3} Such questioning of brain death criteria by reputable physicians and philosophers should not be taken lightly and shows that presumed consent for a practice that may involve doctors killing patients is premature.

Finally, before any notion of presumed consent can be considered, the issue of adequate informed consent to organ donation must be addressed. Without adequate informed consent, the current system of voluntary consent for organ donation is problematic. Any system of presumed consent is also problematic.

Michael Potts professor of philosophy
Methodist University, Fayetteville, NC 28311-1498, USA
gratiaetnatura@yahoo.com

Competing interests: None declared.

- 1 English V. Is presumed consent the answer to organ shortages? Yes. *BMJ* 2007;334:1088. (26 May.)
- 2 Potts M, Byrne PA, Nilges RG, eds. *Beyond brain death: the case against brain-based criteria for human death*. Dordrecht: Kluwer, 2000.
- 3 De Mattei R, ed. *Finis vitae: is brain death still life?* Rome: Consiglio Nazionale delle Ricerche, 2006.

Is prognosis key in donation?

Potts (previous letter) highlights an often misunderstood aspect of organ donation: that death is not always what it seems, especially to relatives.¹ However, I wonder whether his concerns are predominantly motivated by theological considerations?

I suspect the key word is prognosis, as accepted by a wide consensus of peers. The heart may still be beating, but if the prognosis is bleak why not let someone else have a chance of a longer life?

Richard Bartley physiotherapist
Denbigh Infirmary, Denbigh, Clwyd LL16 3ES
monty.python@mac.com

Competing interests: None declared.

- 1 English V. Is presumed consent the answer to organ shortages? Yes. *BMJ* 2007;334:1088. (26 May.)

Tell public about brain death

Until fairly recently, the definition of death was cardiopulmonary death, not brain death.¹ There is a vast suspicion that doctors will take organs from those who are not "really dead"—this suspicion will seem to be confirmed if the heart is still beating and there seems to be undue haste to harvest the organs.

The public needs to be educated on what "brain dead" means, the difference between brain death and a coma, the need for speed in removal of the organs, and, most importantly, the criteria that must be met to confirm that someone is indeed "dead" before organs will be removed.

Joan McClusky medical writer
New York, NY 10003, USA joanmcnewyork@aol.com
Competing interests: None declared.

- 1 English V. Is presumed consent the answer to organ shortages? Yes. *BMJ* 2007;334:1088. (26 May.)

Add carrying a card to QOF

Why not make possession of a donor card a QOF (quality and outcomes framework) target?¹

A question about possession of a donor card is fairly simple but if the practice was required to discuss donation, and obtain

informed consent, this would require more resources.

The information could be available on the NHS spine, although I share concerns about awareness of contraindications to donation.

Anne Holmes general practitioner
Tithebarn Medical Centre, Stockton on Tees TS19 8RH
anne.holmes@ntlworld.com

Competing interests: None declared.

- 1 English V. Is presumed consent the answer to organ shortages? Yes. *BMJ* 2007;334:1088. (26 May.)

COSMETIC GENITOPLASTY

Surgical solution is becoming acceptable, as for birth

Liao and Creighton refer to idiosyncratic decisions about cosmetic genitoplasty in the absence of local and national guidelines.¹ Having worked as a clinical psychologist in women's health for several years, I have seen patients at various stages on the labial surgery pathway, and the reasons for referral to me have been varied.

Some women have had labial surgery and request further interventions, thus causing concern to their surgeons. Satisfaction with surgery has been professed, but it has not changed how they feel about their bodies. These women are faced either with the prospect of more surgery or the stark reality that such a solution may not offer everything they had hoped. Other referrals are to "cover all bases," check that the patient is in "sound mind," and rarely can a serious psychiatric diagnosis be invoked. Unfortunately, some of these patients have already been given a date for their surgery and think that it can proceed unless the psychologist says otherwise. Thus they are unlikely to embrace a psychological rather than a surgical solution.

My surgical colleagues seem to be reluctant to provide such interventions, but they respond to psychological distress and can be disempowered by the general rhetoric of "patient choice." Surgical solutions for various concerns about the body may be what patients seem to want, but increasing the availability of surgery can inhibit the visibility of other choices.

Some clinicians may believe that labial surgery is on the fringes of obstetrics and gynaecology. But history repeats itself. We have all seen the rise in caesarean section rates in the United Kingdom. This is reported to have happened because of concerns about litigation, but another process may be in operation—the sanitisation of a surgical solution to giving birth. This comes from the widespread availability of a surgical solution. Do we not already have a snapshot of how things will develop if we do not debate this issue now?

Competing interests: None declared.

Jacqueline Doyle clinical psychologist, Department of Clinical Psychology, Hillingdon Hospital, Uxbridge, Middlesex UB8 3NN jacqueline.doyle@thh.nhs.uk

- 1 Liao LM, Creighton SM. Requests for cosmetic genitoplasty: how should healthcare providers respond? *BMJ* 2007;334:1090-2. (26 May.)

PHARMACISTS IN PRIMARY CARE

Study shows wrong people, wrong skills, wrong tools . . .

Salter et al identify some deficiencies in a small sample of pharmacist consultations.¹ Their results say more about the need for consultation skills training and the context of the pharmacists and patients involved in this study than they do about the concept of pharmacist medication review.

The pharmacists involved in this study were the wrong people because they had no connection with the patient, the general practitioner, the local pharmacy, or the hospital department and therefore lacked credibility. They had the wrong skills because they scrupulously avoided exploring patients' ideas and beliefs and persisted in a predetermined agenda that patients did not identify with. They had the wrong tools because they did not have the medical records or any indication for the medicines. They were doing the wrong job because people who have just had their medicines reviewed are not likely to benefit from a further review. And the timing was wrong because older people just discharged from the turmoil of hospital need some time to settle and reflect before someone intervenes again.

This study says little about pharmacists' ability to conduct medication reviews. The HOMER study was not a realistic model for pharmacist medication review.² The discharge note must be reviewed in context of the clinical record and in discussion with the general practitioner. Only then should the patient be visited. Their paper is

therefore a lesson in the need to construct a useful intervention before setting out to test it. It also reminds us of the need to learn Osler's century old lesson "Listen to the patient, he's telling you the diagnosis."

Duncan R Petty lecturer practitioner

Arnold Zermansky

University of Leeds, Leeds LS2 9UT d.petty@leeds.ac.uk

Competing interests: None declared.

- 1 Salter C, Holland R, Harvey R, Henwood K. "I haven't even phoned my doctor yet." The advice giving role of the pharmacist during consultations for medication review with patients aged 80 or more: qualitative discourse analysis. *BMJ* 2007;334:1101-4. (26 May.)
- 2 Holland R, Lenaghan E, Harvey I, Smith R, Shephstone L, Lipp A, et al. Does home based medication review keep older people out of hospital? The HOMER randomised controlled trial. *BMJ* 2005;330:293.

ANAEMIA IN DEVELOPING COUNTRIES

Mass iron treatment is cheaper than routine deworming

Gulani et al say that routine administration of intestinal anthelmintic agents results in a marginal increase in haemoglobin (1.71 g/l).¹ What needs to be considered is whether this approach of mass anthelmintic therapy is actually economically feasible, especially in third world countries where iron deficiency anaemia is a major health issue.² This needs special consideration, given the fact that the primary cause of anaemia in third world countries is dietary malnutrition rather than intestinal infestation with helminths.³

A better and more economically feasible approach to thwart the "epidemic" of anaemia might be mass supplementation with iron supplements such as oral ferrous sulphate.⁴ The average cost of mebendazole treatment (100 mg three times a day for three days) is £15. According to Gulani et al, this regimen increases haemoglobin by 1.7 g/l. On the other hand, ferrous sulphate at a dose of 325 mg three times a day will increase haemoglobin by the same amount in about two weeks and cost £1.

Shailendra Kapoor resident physician

University of Illinois at Chicago, Chicago, Schaumburg, IL, 60195, USA shailendrakapoor@yahoo.com

Competing interests: None declared.

- 1 Gulani A, Nagpal J, Osmond C, Sachdev HP. Effect of administration of intestinal anthelmintic drugs on haemoglobin: systematic review of randomised controlled trials. *BMJ* 2007;334:1095-7. (26 May.)
- 2 Cook JD, Reusser ME. Iron fortification: an update. *Am J Clin Nutr* 1983;38:648-59.
- 3 Tatala S, Svanberg U, Mduma B. Low dietary iron availability is a major cause of anemia: a nutrition survey in the Lindi District of Tanzania. *Am J Clin Nutr* 1998;68:171-8.
- 4 Mitra AK, Akramuzzaman SM, Fuchs GJ, Rahman MM, Mahalanabis D. Long-term oral supplementation with iron is not harmful for young children in a poor community of Bangladesh. *J Nutr* 1997;127:1451-5.

MTAS

Lessons from the disaster

Some lessons must be swiftly learnt from the Medical Training Application Service (MTAS) experience.¹ The BMA and the Academy of Medical Royal Colleges have been made to look feeble and ineffectual after entering into "partnership" roles with the Department of Health. The postgraduate deans have been notably silent and have behaved as willing accomplices in the promotion of MTAS. The deaneries have declined as an independent force in medical training and are struggling to fulfil their correct role in providing quality educational leadership because of over-dependence on political approval linked to their funding mechanism. The reputation of UK medical training has taken a damaging hit.

The royal colleges, threatened by loss of power and influence and undermined by the emergence of the Postgraduate Medical Education and Training Board (PMETB), seem to have been all too easily lured into partnership agreements, using a set of desirable motherhood and apple pie objectives that were seductively easy to sign up to. The trap was then sprung and the poorly drafted and unworkable operational details released deliberately late in the process. The timetable of the action plan became a higher priority than the quality of the project itself. Website design and selection procedures were unfinished, and consultation on details was token or non-existent. Bullying tactics created an unstoppable momentum for MTAS implementation, regardless of the obvious problems piling up and the well based objections of a majority of consultants. The colleges are protesting that they have been misrepresented, although manipulated would be more accurate with junior doctors feeling disconnected and unsupported until it was too late. The government, anxious to displace blame elsewhere, insists they were fully on board.

These "partnership" arrangements have become a damaging form of pseudo collaboration. The end result has been a major system crash between the Department of Health and the profession, which is now much deeper than the single catastrophe of MTAS.

John J Turner consultant physician, University Hospital Aintree, Liverpool L9 7AL john.turner@aintree.nhs.uk

Competing interests: JJT is a consultant and educational supervisor.

- 1 Godlee F. The future of specialist training. *BMJ* 2007;334:1067-8. (26 May.)