We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer. Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.

LETTERS

COMSTOCK COMPLETE

DOCTORS AND ALCOHOL

Investing in alcohol is no longer responsible

"Doctors are stalwart drinkers," says Richards, calling on doctors to do more to tackle alcohol abuse.¹ Doctors are also stalwart investors and must begin to use their financial clout as shareholders to remove alcohol from their portfolios.

Last autumn, the multinational drinks industry used its political, business, and financial lobbying muscle to subvert the European Commission's attempts to bring forward an alcohol strategy based on public health principles.² In the past, the public health agenda has relied solely on evidence to effect political change. This worked when elected politicians had a stronger influence on society. The business corporations have now become bigger players where the measure of success is profit. Health policy has to accommodate to this change and will have to engage shareholders directly to get them to avoid alcohol at both manufacturing and the sometimes forgotten retailing levels.

Leadership is required from doctors, who are often the first witnesses of alcohol related harm. This can be exercised by refusing to hold shares in alcohol companies and by instructing fund managers to set up and seek non-alcohol based portfolios. It is time to include the manufacture and importantly the sale of alcohol as an unworthy and unethical way of making money for the individual investor. Most doctors would not knowingly invest in tobacco companies, and, although no level of tobacco use is safe, alcohol is no ordinary commodity either³ and even small amounts can cause catastrophic harm.

Advocacy budgets will never come

remotely near industry budgets, but doctors interested in the health of the public have to develop expertise on the ways of big business and use this expertise in informing individual shareholders on the wider societal implications of their investments.

Joseph M Barry clinical professor, Department of Public Health and Primary Care, Trinity College Centre for Health Sciences, AMNCH Tallaght, Dublin 24, Republic of Ireland
joebarry@tcd.ie

Tom O'Dowd professor of general practice

Competing interests: None declared.

- 1 Richards T. Europe on the rocks. *BMJ* 2007;334:1142. (2 June.)
- 2 McKee M. A European alcohol strategy: will the opportunity be missed? *BMJ* 2006;333:871-2.
- 3 Babor T, Caetano R, Casswell, S, Edwards G, Giesbrecht N, Graham K, et al. *Alcohol: no ordinary commodity. Research and public policy.* Oxford and London: Oxford University Press, 2003.

SAFER PLAY IN RUGBY

Time to address safety in the tackle

As an amateur rugby referee who spent the 2005 season refereeing in New Zealand and took part in the RugbySmart campaign I was interested to read the analysis by Quarrie et al of its impact on spinal injuries.¹ As they point out, most of the reduction in injuries was accounted for by a reduction in scrum related injuries. Whether a similar campaign would be equally effective in Britain is unclear, given the recent 2006 changes in the scrum engagement procedure; but as pointed out in the accompanying editorial it can certainly do no harm.²

The paper highlighted another area of concern; seven out of eight spinal injuries occurred at the tackle. Over the past 10 years there has been an increase in the high impact, chest high "ball and all" tackle. This area of the game is now associated with the most injuries,³ and this type of tackle puts the tackling player at risk of a head on torso collision and spinal injury.

Instead of further debates over banning the scrum we should concentrate on improving the safety at the tackle area by legislation and educational initiatives similar to those described by Quarrie et al.

Alastair P Greystoke research fellow, medical oncology
Paterson Institute of Cancer Research, University of

Manchester, Manchester M20 4BX
agrey@picr.man.ac.uk

Competing interests: APG is a rugby referee who took part in the RugbySmart campaign.

- 1 Quarrie KL, Gianotti SM, Hopkins WG, Hume PA. Effect of nationwide injury prevention programme on serious spinal injuries in New Zealand rugby union: ecological study. *BMJ* 2007;334:1150-3. (2 June.)
- 2 Noakes TD, Draper CE. Preventing spinal cord injuries in rugby union. *BMJ* 2007;334:1122-3. (2 June.)
- 3 Fuller CW, Brooks JH, Cancea RJ, Hall J, Kemp SP. Contact events in rugby union and their propensity to cause injury. *Br J Sports Med* 2007; May 18. [Epub ahead of print] PMID: 17513332.

PTSD IN NORTHERN IRELAND

Tell us more

Duffy et al provide evidence for the effectiveness of cognitive therapy in post-traumatic stress disorder (PTSD) in the context of terrorism and civil conflict in Northern Ireland.¹ More information would have been helpful to interpret the results.

No patients were started on medications during the trial. However, 52% in the immediate therapy group were taking antidepressants already. When were these initiated in relation to the trial? Also, were any changes to the antidepressant dose allowed during the trial? Over 70% in the immediate therapy group had comorbid major depression. The effect of antidepressant initiation just before the trial or dose changes may be partly responsible for the improvement in this group's symptoms.

The percentages for the overall effectiveness of cognitive therapy are the combined scores of the immediate treatment and waiting list control groups. This makes them uncontrolled scores. The authors are not comparing two groups of patients, one receiving therapy and the other not receiving therapy.

The follow-up mean scores in table 3 have been taken at either four or 12 months. As a clinician, I would be particularly interested in information about the maintenance of gains at 12 months. This is not clear from the table. If gains seen at four months are lost by 12 months, this then raises questions about whether booster sessions are indicated.

Finally, the therapist effect is important. It would be interesting to look at whether

this difference in patient scores is related to the type of qualification in cognitive therapy that the therapists had. Recent research has shown that formal post-qualification training in cognitive therapy is associated with competence.²

Adarsh Shetty specialist registrar in general adult psychiatry
Psychiatric Unit, Derby City Hospital, Derby DE22 3NE
dradarshshetty@yahoo.co.in

Competing interests: None declared.

- 1 Duffy M, Gillespie K, Clark DM. Post-traumatic stress disorder in the context of terrorism and other civil conflict in Northern Ireland: randomised controlled trial. *BMJ* 2007;334:1147-50. (2 June.)
- 2 Brosan L, Reynolds S, Moore RG. Factors associated with competence in cognitive therapists. *Behav Cognitive Psychother* 2007;35:179-90.

SKIN BIOPSY

Watch out for digital nerves

As a hand surgeon, I was concerned about the bottom right picture in figure 3 of the article by Lauria and Lombardi, which seems to show that a skin biopsy had been taken from the radiovolar skin of the left index finger.¹ In the summary points box it was suggested that a skin biopsy was easy and almost painless. Furthermore, nowhere was there any indication that potential injuries could be caused by the use of this technique. Skin biopsy as indicated above carries a high risk of causing a digital nerve injury leading to a neuroma causing chronic pain, which may require reconstructive surgery. If volar digital skin biopsies are required then it is much safer to harvest these in the midline of the finger, but patients should be warned of the potential nerve injury and neuroma risk as important nerves are only a few millimetres under the skin surface on fingers.

Bo Povlsen consultant orthopaedic surgeon, Guy's Hospital, London SE1 7EH bo@manusmedical.com

Competing interests: None declared.

- 1 Lauria G, Lombardi R. Skin biopsy: a new tool for diagnosing peripheral neuropathy. *BMJ* 2007;334:1159-62. (2 June.)

ROSIGLITAZONE AND HEART DEATHS

Glycaemic control is a myth

Tight control of HbA1c levels has been enshrined in the QOF (quality and outcomes framework) of the new general practitioner contract as being an evidence based proposal. Multiple drugs are licensed on the understanding that they reduce HbA1c levels and that this is a good thing. So should it surprise us that a meta-analysis of trials of rosiglitazone shows a raised risk

of myocardial infarction and an increase in cardiovascular deaths?¹

The reduction in diabetes related end points, mortality and stroke from using metformin is not explicable on the basis of glycaemic control.²

The data of UKPDS 33, which compared tight glycaemic control with sulphonylureas or insulin with conventional treatment, showed little benefit from tight control.³ The outcomes that did show some clinical benefit were cataract extractions, retinal photocoagulation, and non-fatal myocardial infarction and all cause mortality—that is, if you can call absolute risk reductions of between 1 and 3 per 1000 patient years as being clinically relevant.

The wonder with drug licensing is that we continue to accept surrogate end points in trials to license new treatments for conditions for which we already have treatments. Show me better data or accept that the control of blood glucose means metformin—anything else is merely for symptom control.

Illtyd R Thomas general practitioner, Swansea SA1 5LF
illtyd.thomas@gp-w98027.wales.nhs.uk

Competing interests: IRT has coauthored with Adrian Edwards, Glyn Elwin, and Rhys Williams a paper on explaining risk information over the internet to patients with diabetes, which was funded by the BMJ Group.

- 1 Tanne JH. Study indicates diabetes drug linked to cardiovascular death. *BMJ* 2007;334:1073. (26 May.) doi: 10.1136/bmj.39224.364630.DB
- 2 Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet* 1998;352:854-65.
- 3 Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risks of complications in patients with type 2 diabetes (UKPDS 33). *Lancet* 1998;352:837-53.

HELPING THE WORLD'S POOR

Let's help doctors work in the third world

Surely it is not beyond the wit of our profession to devise a scheme that will make it easier for UK health professionals to work overseas either as missionaries, volunteers with non-governmental organisations (NGOs), or in other “bona fide” set-ups. Some of the issues Mabey raises also affect UK doctors wanting to take career breaks in the United Kingdom.¹

We need to devise a way that UK doctors working overseas can be part of some form of revalidation. We have an oversupply of doctors at the moment trying to get into the Modernising Medical Careers (MMC) process. Allowing some to opt out to work overseas should be encouraged by recognising the experience they will have

gained and facilitating their re-entry into the system.

At the moment UK doctors working overseas cost the UK government nothing. The government could recognise this by helping to pay off debts from student loans and help with the General Medical Council's registration fees, etc. This money could come out of the overseas aid budget.

Some organisations such as the BMA and Medical Protection Society recognise overseas service by offering either free or heavily discounted membership to medical missionaries and NGO volunteers.

Andrew L Perkins former medical missionary, Westoning, Bedfordshire MK45 5JD alperkins@doctors.org.uk

Competing interests: ALP was a medical missionary in Mali, West Africa.

- 1 Mabey D. Improving health for the world's poor. *BMJ* 2007;334:1126. (2 June.)

TRIALS AND ELECTRONIC RECORDS

A frightening industry proposal

The National Care Record Service (CRS), if it is ever deployed, certainly offers amazing potential for pharmaceutical research. The whole COX2/NSAID debacle could have been rapidly resolved by access to the complete prescription records of 55 million people.

But this article sent shudders down my spine about how the Association of the British Pharmaceutical Industry would like to use the service.¹ Dr Barker is quoted as saying it would allow drug companies to easily identify patients fitting a trial's inclusion and exclusion criteria.

Not using anonymised data, it wouldn't. I suppose it might tell you how many potential candidates there were, but that shouldn't be too hard to figure out anyway. Anonymised data use for adverse events surveillance is one thing; non-anonymised data to identify potential trial candidates, presumably followed by a direct approach to invite them to join a drug trial, are another matter entirely.

This use is clearly in conflict with current data protection legislation—but Dr Barker should realise this. So why did he propose it?

Matthew L Grove consultant rheumatologist, North Tyneside General Hospital, North Shields NE29 8NH
matthew.grove@nhct.nhs.uk

Competing interests: None declared.

- 1 Butcher J. UK will lose clinical trials if electronic records system is delayed, ABPI warns. *BMJ* 2007;334:1132. (2 June.)