Published in final edited form as:

Biochem Biophys Res Commun. 2007 April 27; 356(1): 97–101.

Oxidation of Calmodulin Alters Activation and Regulation of CaMKII

AJ Robison^{a,†}, Danny G. Winder^a, Roger J. Colbran^a, and Ryan K. Bartlett^{a,‡}

^a*Vanderbilt University Medical Center, Department of Molecular Physiology and Biophysics, Nashville, TN 37232, USA*

Abstract

Increases in reactive oxygen species and mis-regulation of calcium homeostasis are associated with various physiological conditions and disease states including aging, ischemia, exposure to drugs of abuse, and neurodegenerative diseases. In aged animals, this is accompanied by a reduction in oxidative repair mechanisms resulting in increased methionine oxidation of the calcium signaling protein calmodulin in the brain. Here, we show that oxidation of calmodulin results in an inability to: 1) activate CaMKII; 2) support Thr²⁸⁶autophosphorylation of CaMKII; 3) prevent Thr^{305/6}autophosphorylation of CaMKII; 4) support binding of CaMKII to the NR2B subunit of the NMDA receptor; 5) compete with α -actinin for binding to CaMKII. Moreover, oxidized calmodulin does not efficiently bind calcium/calmodulin-dependent protein kinase II (CaMKII) in rat brain lysates or *in vitro*. These observations contrast from past experiments performed with CaMox and the Plasma Membrane Calcium ATPase, where CaMox binds to, and partially activates the PMCA. When taken together, these data suggest that oxidative stress may perturb neuronal and cardiac function via a decreased ability of oxidized calmodulin to bind, activate, and regulate the interactions of CaMKII.

Introduction

Calmodulin (CaM) is a ubiquitous intracellular calcium (Ca^{2+}) sensor essential for rapid and coordinated responses of a variety of enzymes, channels, and receptors to both local and global Ca^{2+} transients. CaM contains two globular EF-hand domains on each terminus, each capable of binding two Ca^{2+} ions, connected by a central flexible helix. Binding of Ca^{2+} to the EF-hands conveys a conformational change resulting in the N- and C-terminal domains exposing hydrophobic binding pockets that allow CaM to bind amphipathic α -helices in target molecules. CaM can also bind targets in the apo or partially saturated (2 Ca^{2+} ions bound) forms. These targets also include IQ motifs, such as those found in the L-type calcium channel, neuromodulin, and many others.

Neuronal Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII) is a ubiquitous target of CaM [1]. Under basal conditions, the kinase exists as a 12-subunit holoenzyme, with individual CaMKII subunits inhibited by binding of the autoinhibitory domain (AID) at the catalytic cleft. Ca^{2+} -dependent binding of CaM to a region overlapping the AID relieves this autoinhibition,

[‡]Author to whom correspondence should be addressed: ryan.bartlett@vanderbilt.edu, Vanderbilt University Medical Center, 724 RRB, Nashville, TN 37232, Ph: +1 615-322-4389

[†]Current Address: UT Southwestern Medical Center, Dallas, TX 75390

Funding: This work was funded by NIH #MH63232

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

allowing kinase activity. Ca^{2+} /CaM-dependent Thr²⁸⁶ autophosphorylation between adjacent subunits in the holoenzyme enhances the affinity for Ca^{2+} /calmodulin and confers autonomous (Ca^{2+} -independent) activity after CaM dissociates. Thr^{305/306} autophosphorylation occurs only in the absence of bound Ca^{2+} /CaM and blocks subsequent CaM binding. The calcium saturated carboxy-terminal lobe of CaM can partially activate CaMKII [2].

CaMKII is essential for diverse physiological processes such as Ca^{2+} -dependent long-term potentiation in the brain [3,4] and L-type Ca^{2+} -channel regulation in the heart [5]. Ca^{2+} homeostasis in these and other regions of the body are adversely affected by oxidative stress, due to ischemia, disease, or aging [6]. Multiple methionine residues within CaM can become oxidized to methionine sulfoxide (CaMox) during normal aging in the rat, causing conformational changes that alter normal association with target peptides [7]. For example, CaM isolated from the aged rat only partially activates, but still binds, the Plasma Membrane Ca^{2+} ATPase (PMCA) [7,8]. Thus, deleterious effects of oxidative stress may occur in part due to the inability of CaMox to properly bind and activate multiple target proteins. Previous studies show that mutation of specific methionine residues to glutamine effects CaMKII autophosphorylation [9], however the glutamine mutation does not always mimic methionine oxidation, and more detailed studies are required to look at all aspects of CaMKII activation and regulation by CaMox. Here, we show that progressive oxidation of CaM causes an inability to bind, activate, and regulate the interactions of CaMKII with various CaMKII associated proteins (CaMKAPs).

Materials and Methods

Materials

H_2O_2 , CaM-agarose, and glutathione agarose: Sigma (St. Louis, MO). Phenyl Sepharose CL-4B FF: Pharmacia (Piscataway, NJ). Goat α -CaMKII α antibody was previously described [10]. Alkaline phosphatase-conjugated secondary antibody: Jackson ImmunoResearch (West Grove, PA). Phospho-T²⁸⁶CaMKII antibody: Promega (Madison, WI). Phospho-T³⁰⁵ CaMKII antibody: Gift from A.J. Silva [11].

Protein purification

Recombinant CaM, CaMKII α , T286D CaMKII, and glutathione-S-transferase (GST) fusion proteins containing NR2B and α -actinin fragments were expressed and purified as described previously [12-14]. GST-NR2B contains amino acids 1260-1339 of NR2B, with Ser¹³⁰³ mutated to alanine, and GST-actinin contains amino acids 819-894 of α -actinin-2. All protein concentrations were quantified via extinction coefficients and confirmed with Pierce BCA protein assay (GST fusions) or Bradford protein assay (CaMKII).

CaM Oxidation

CaM oxidation was performed using H_2O_2 as previously described [15]. For progressive oxidation, aliquots of the oxidation reaction were removed and placed immediately into dialysis at the indicated times (see Figure 2). CaM agarose was oxidized using the same protocol, except that H_2O_2 was removed via extensive washing of the resin with equilibration buffer.

Brain Lysate Preparation

Rat forebrain lysates were prepared by homogenization in 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 5 mM CaCl_2 , 0.25% Triton X-100, protease inhibitor cocktail (Sigma, St. Louis, MO), 1 mM Benzamide, 1 mM PMSF, and 1 μM microcystin. Equal aliquots of lysate were loaded onto the CaM agarose, or CaMox agarose (2 mL packed resin equilibrated with 50 mM Tris-

HCl pH 7.5, 150 mM NaCl, 5 mM CaCl₂(Buffer A)). The resins were then washed with buffer A (20 mL), and eluted with 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 5 mM EDTA.

CaMKII Activity Assays and Autophosphorylation

Kinase assays were performed using the CaMKII model substrate Syntide-2 [16]. CaMKII α was autophosphorylated at T²⁸⁶ and T^{305/6} as described [10,12].

GST-Cosedimentation Assays

GST-cosedimentation assays were performed as described [12]. Proteins on the nitrocellulose membrane were visualized by staining with Ponceau stain and identities were confirmed by Western blot (data not shown).

Results

Progressive oxidation of CaM prevents CaMKII activation

Ca²⁺-dependent binding of CaM to CaMKII activates both T²⁸⁶ autophosphorylation and phosphorylation of exogenous substrates. In order to determine the effects of CaM oxidation, CaM was exposed to H₂O₂ for times ranging from 1 minute to 3 days (total). This method of CaM oxidation has been proven in the past to only target methionines and that methionine oxidation is progressive under these conditions [15,17,18]. Analysis by SDS-PAGE revealed progressive increases in apparent molecular weight over the incubation time consistent with progressive oxidation (Figure 1A) [19]. At least five distinct modified forms of CaM were visible, presumably due to the presence of mixed populations of CaM with distinct methionine residues, or combinations of residues, converted to methionine sulfoxide. The ability of partially oxidized CaM to activate Ca²⁺-dependent CaMKII activity was then determined. Increasing oxidation of CaM caused a progressive rightward shift in the CaMKII activation curve, and over-night exposure of CaM to H₂O₂ completely prevented CaM activation of CaMKII (Figure 1B). This data contrasts data collected in the same manner using the PMCA. CaM that is fully oxidized is still capable of binding to and partially activating the PMCA [15].

In order to determine whether oxidized CaM supports Thr²⁸⁶ autophosphorylation, CaMKII was incubated on ice with non-oxidized or fully oxidized CaM in the presence of Ca²⁺, Mg²⁺, and [γ -³²P]-ATP [10]. Incubation with oxidized CaM resulted in no detectable CaMKII autophosphorylation as assessed by autoradiography or phospho-Thr²⁸⁶-specific immunoblotting (Figure 2A,B). CaMKII autophosphorylation was not detected even when these incubations were repeated at 30°C (Figure 2B).

Ca²⁺/CaM binding to CaMKII blocks autophosphorylation of Thr³⁰⁵ and Thr³⁰⁶ within the CaM binding domain (Figure 2C). Even though CaMox cannot activate CaMKII, it is possible that it retains the capacity to block autophosphorylation at Thr^{305/306}. Therefore, we incubated CaMKII α [T286D] with ATP, Mg²⁺, and Ca²⁺ in the presence of either CaM or CaMox. The T286D mutant was used because constitutive activity of this kinase enhanced the CaM-dependent autophosphorylation at Thr^{305/306} and other sites (not shown), resulting in a form of CaMKII with reduced mobility on SDS-PAGE and strong reactivity toward a P-Thr³⁰⁶-specific antibody. Addition of CaM substantially reduced this autophosphorylation reaction, but CaMox appears to have no detectable effect on this form of autophosphorylation (Figure 2C).

CaMox does not properly regulate CaMKII-CaMKAP interactions

CaM differentially modulates CaMKII interaction with a variety of proteins [12,20,21]. For example, Ca²⁺/CaM binding promotes interaction of non-autophosphorylated CaMKII with

the NR2B subunit of the N-methyl-D-aspartate (NMDA) receptor, whereas Ca^{2+} /CaM competes with α -actinin for binding to CaMKII [21]. However, CaMox fails to support CaMKII binding to NR2B in cosedimentation assays (Figure 3A). CaMox also failed to compete with GST-actinin for binding to CaMKII (Figure 3B). These data indicate that CaMox is unable to properly regulate the interactions of CaMKII with neuronal binding partners.

CaMKII does not bind fully oxidized CaM

In order to directly assay whether CaMox binds CaMKII, oxidized or control CaM-agarose was incubated with purified CaMKII in the presence or absence of Ca^{2+} , washed extensively, and collected by centrifugation. Bound protein was eluted directly into SDS sample buffer, and SDS-PAGE analysis revealed purified CaMKII only in the non-oxidized CaM-agarose pellet, in a Ca^{2+} -dependent manner, and no stable binding at all to the oxidized CaM-agarose was observed (Figure 4A).

CaMKII from rat brain lysate was also tested for binding to CaMox. Soluble extract was prepared from rat homogenized adult rat brain and passed over columns of fully oxidized or control CaM-agarose in the presence of Ca^{2+} [22]. Resin was washed, and proteins stably bound to the CaM-agarose were eluted with EDTA. Fractions containing the eluted proteins were analyzed by SDS-PAGE and western blot. CaMKII bound to the control CaM column and was eluted with EDTA. However, CaMKII was not detected in fractions eluted from the CaMox-agarose. (Figure 4B).

To demonstrate that oxidized CaM agarose is intact and able to interact with some proteins, total elution from reduced CaM agarose and oxidized CaM agarose after incubation with rat brain lysate are shown. Even after oxidation, several neuronal proteins are clearly capable of binding to the oxidized CaM agarose (Figure 4).

Discussion

We show here that progressive oxidation of CaM prevents stable binding to CaMKII. The effects of this are four-fold: 1) Ca^{2+} -dependent activity toward exogenous substrates is not stimulated by CaMox; 2) autophosphorylation at Thr²⁸⁶ is not stimulated by CaMox; 3) autophosphorylation at Thr^{305/6} is not prevented by CaMox; and 4) CaMox cannot support binding to NR2B or disrupt the interaction with α -actinin.

Previous studies have shown that all nine methionine residues of CaM can be oxidized *in vitro*, and at least two residues can be oxidized *in vivo* during normal aging. The reduction in CaMKII activation by progressively oxidized CaM could not be attributed to reduction or accumulation of any single band of CaM as visualized by SDS-PAGE. It is therefore possible that either the reduction in CaMKII activation results from the progressive oxidation of a series of methionine residues within CaM, or that shifts in apparent molecular weight of CaM represent a variety of separate, overlapping patterns of methionine oxidation, with oxidation of only one or two specific residues contributing to the reduction in CaMKII activation. Further experiments, including mutation of methionine residues to non-oxidizable amino acids individually and in groups, will be required to resolve this issue [15]. These effects of CaM oxidation on CaMKII are distinct from previous reports of the effects of oxidation on other targets of CaM. For example, the PMCA is partially activated by fully oxidized CaM with little change in apparent binding affinity [15]. In contrast, oxidation reduces the apparent affinity of CaM for CaMKII and also abolishes CaMKII activation (Fig 1). The differential effect of CaM oxidation in binding to these targets presumably results from differences in the mechanism of interaction.

Blockade of CaMKII activity has deleterious effects in multiple physiological processes. For example, pharmacological blockade of CaMKII activity in the hippocampi of mice prevents multiple types of memory formation [23,24]. Additionally, in the mouse model, blockade of CaMKII activity in the heart can prevent sarcoplasmic reticulum Ca^{2+} release, a process known to be involved in triggered arrhythmias [5,25]. Thus physiological processes involving the generation of reactive oxygen species may result in concomitant deleterious effects on heart function or memory formation in part due to the oxidation of CaM. This oxidation in turn could result in loss of CaMKII activity as well as potential mislocalization due to improper regulation of CaMKII/CaMKAP interactions. Thus, CaM may act as a metabolic redox sensor [6], differentially effecting downstream targets, to modulate the cell's overall metabolic activity during normal aging or under ischemic conditions. In this model, one could envision CaMox acting to negatively regulate the activity of CaMKII toward targets that directly affect calcium feedback during times of disrupted calcium homeostasis.

Additional in depth experiments will be required to specifically identify individual oxidized methionines responsible for the loss of the ability of CaM to activate CaMKII. This type of information is crucial for interpreting disrupted signaling cascades during times of oxidative stress.

Literature Cited

1. Hudmon A, Schulman H. Structure-function of the multifunctional Ca^{2+} /calmodulin-dependent protein kinase II. *Biochem J* 2002;364:593–611. [PubMed: 11931644]
2. Shifman JM, Choi MH, Mihalas S, Mayo SL, Kennedy MB. Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII) is activated by calmodulin with two bound calciums. *Proc Natl Acad Sci U S A* 2006;103:13968–13973. [PubMed: 16966599]
3. Lisman J, Schulman H, Cline H. The molecular basis of CaMKII function in synaptic and behavioural memory. *Nat Rev Neurosci* 2002;3:175–190. [PubMed: 11994750]
4. Colbran RJ, Brown AM. Calcium/calmodulin-dependent protein kinase II and synaptic plasticity. *Curr Opin Neurobiol* 2004;14:318–327. [PubMed: 15194112]
5. Zhang R, Khoo MS, Wu Y, Yang Y, Grueter CE, Ni G, Price EE Jr, Thiel W, Guatimosim S, Song LS, Madu EC, Shah AN, Vishnivetskaya TA, Atkinson JB, Gurevich VV, Salama G, Lederer WJ, Colbran RJ, Anderson ME. Calmodulin kinase II inhibition protects against structural heart disease. *Nat Med* 2005;11:409–417. [PubMed: 15793582]
6. Squier TC, Bigelow DJ. Protein oxidation and age-dependent alterations in calcium homeostasis. *Front Biosci* 2000;5:D504–526. [PubMed: 10799358]
7. Anbanandam A, Bieber Urbauer RJ, Bartlett RK, Smallwood HS, Squier TC, Urbauer JL. Mediating molecular recognition by methionine oxidation: conformational switching by oxidation of methionine in the carboxyl-terminal domain of calmodulin. *Biochemistry* 2005;44:9486–9496. [PubMed: 15996103]
8. Gao J, Yao Y, Squier TC. Oxidatively modified calmodulin binds to the plasma membrane Ca-ATPase in a nonproductive and conformationally disordered complex. *Biophys J* 2001;80:1791–1801. [PubMed: 11259292]
9. Chin D, Means AR. Methionine to glutamine substitutions in the C-terminal domain of calmodulin impair the activation of three protein kinases. *J Biol Chem* 1996;271:30465–30471. [PubMed: 8940012]
10. McNeill RB, Colbran RJ. Interaction of autophosphorylated Ca^{2+} /calmodulin-dependent protein kinase II with neuronal cytoskeletal proteins. Characterization of binding to a 190-kDa postsynaptic density protein. *J Biol Chem* 1995;270:10043–10049. [PubMed: 7730306]
11. Elgersma Y, Fedorov NB, Ikonen S, Choi ES, Elgersma M, Carvalho OM, Giese KP, Silva AJ. Inhibitory autophosphorylation of CaMKII controls PSD association, plasticity, and learning. *Neuron* 2002;36:493–505. [PubMed: 12408851]

12. Robison AJ, Bass MA, Jiao Y, MacMillan LB, Carmody LC, Bartlett RK, Colbran RJ. Multivalent interactions of calcium/calmodulin-dependent protein kinase II with the postsynaptic density proteins NR2B, densin-180, and alpha-actinin-2. *J Biol Chem* 2005;280:35329–35336. [PubMed: 16120608]
13. Brickey DA, Bann JG, Fong YL, Perrino L, Brennan RG, Soderling TR. Mutational analysis of the autoinhibitory domain of calmodulin kinase II. *J Biol Chem* 1994;269:29047–29054. [PubMed: 7961870]
14. Brickey DA, Colbran RJ, Fong YL, Soderling TR. Expression and characterization of the alpha-subunit of Ca²⁺/calmodulin-dependent protein kinase II using the baculovirus expression system. *Biochem Biophys Res Commun* 1990;173:578–584. [PubMed: 2175600]
15. Bartlett RK, Bieber Urbauer RJ, Anbanandam A, Smallwood HS, Urbauer JL, Squier TC. Oxidation of Met144 and Met145 in calmodulin blocks calmodulin dependent activation of the plasma membrane Ca-ATPase. *Biochemistry* 2003;42:3231–3238. [PubMed: 12641454]
16. Colbran RJ. Inactivation of Ca²⁺/calmodulin-dependent protein kinase II by basal autophosphorylation. *J Biol Chem* 1993;268:7163–7170. [PubMed: 8385100]
17. Yao Y, Yin D, Jas GS, Kuczer K, Williams TD, Schoneich C, Squier TC. Oxidative modification of a carboxyl-terminal vicinal methionine in calmodulin by hydrogen peroxide inhibits calmodulin-dependent activation of the plasma membrane Ca-ATPase. *Biochemistry* 1996;35:2767–2787. [PubMed: 8611584]
18. Yin D, Kuczera K, Squier TC. The sensitivity of carboxyl-terminal methionines in calmodulin isoforms to oxidation by H₂O₂ modulates the ability to activate the plasma membrane Ca-ATPase. *Chem Res Toxicol* 2000;13:103–110. [PubMed: 10688534]
19. Ferrington DA, Sun H, Murray KK, Costa J, Williams TD, Bigelow DJ, Squier TC. Selective degradation of oxidized calmodulin by the 20 S proteasome. *J Biol Chem* 2001;276:937–943. [PubMed: 11010965]
20. Colbran RJ. Targeting of calcium/calmodulin-dependent protein kinase II. *Biochem J* 2004;378:1–16. [PubMed: 14653781]
21. Robison AJ, Bartlett RK, Bass MA, Colbran RJ. Differential modulation of Ca²⁺/calmodulin-dependent protein kinase II activity by regulated interactions with N-methyl-D-aspartate receptor NR2B subunits and alpha-actinin. *J Biol Chem* 2005;280:39316–39323. [PubMed: 16172120]
22. Yan P, Xiong Y, Chen B, Negash S, Squier TC, Mayer MU. Fluorophore-assisted light inactivation of calmodulin involves singlet-oxygen mediated cross-linking and methionine oxidation. *Biochemistry* 2006;45:4736–4748. [PubMed: 16605242]
23. Wolfman C, Fin C, Dias M, Bianchin M, Da Silva RC, Schmitz PK, Medina JH, Izquierdo I. Intrahippocampal or intraamygdala infusion of KN62, a specific inhibitor of calcium/calmodulin-dependent protein kinase II, causes retrograde amnesia in the rat. *Behav Neural Biol* 1994;61:203–205. [PubMed: 8067974]
24. Mayford M, Bach ME, Huang YY, Wang L, Hawkins RD, Kandel ER. Control of memory formation through regulated expression of a CaMKII transgene. *Science* 1996;274:1678–1683. [PubMed: 8939850]
25. Picht E, Desantiago J, Huke S, Kaetzel MA, Dedman JR, Bers DM. CaMKII inhibition targeted to the sarcoplasmic reticulum inhibits frequency-dependent acceleration of relaxation and Ca(2+) current facilitation. *J Mol Cell Cardiol*. 2006

Abbreviations

CaM	calmodulin
CaMKII	Calcium Calmodulin dependent Kinase II
GST	glutathione-S-transferase

Figure 1. Oxidation of calmodulin blocks CaMKII activation

A. Purified calmodulin was exposed to 100 mM H_2O_2 for various times and then analyzed by SDS-PAGE on a 4-20% acrylamide gel and Coomassie stained. Calmodulin unexposed to H_2O_2 (Non) or totally oxidized (Tot.) were included as controls. B. CaMKII was incubated with Ca^{2+} , Mg^{2+} , $[\gamma^{32}\text{P}]\text{ATP}$, and the peptide substrate syntide-2 to look at the ability of CaM vs. CaMox to activate CaMKII in a calcium-dependent manner.

Figure 2. Oxidized calmodulin does not regulate CaMKII autophosphorylation

A. CaMKII was incubated with Ca^{2+} , Mg^{2+} , $[\gamma^{32}\text{P}]\text{ATP}$, and oxidized or control calmodulin. Reactions were stopped at indicated times by the addition of SDS sample buffer. Samples were separated by SDS-PAGE and analyzed for total CaMKII by Ponceau stain (top), for total incorporation of radioactive phosphate by autoradiography (middle), and for specific autophosphorylation at Thr²⁸⁶ by Western blot with a phospho-specific antibody (bottom). **B.** Samples from panel A were quantified for total incorporation of radioactive phosphate into CaMKII by scintillation counter. Error bars represent standard error of the mean, $n=3$. **C.** CaMKII[T286D] was incubated with or without oxidized or control calmodulin under conditions which stimulate Thr^{305/6} autophosphorylation. Samples were separated by SDS-PAGE and analyzed for Thr^{305/6} autophosphorylation by Western blot with a phospho-specific antibody.

Figure 3. Oxidized calmodulin does not properly regulate CaMKII interaction with synaptic proteins

A. CaMKII was incubated with GST-NR2B and glutathione agarose in the presence or absence of oxidized or control calmodulin, 2 mM Ca²⁺, or 100 μ M ADP, washed extensively, and eluted in SDS sample buffer. Supernatant (Super) and eluate (Pellet) were separated by SDS-PAGE and analyzed by Ponceau stain. **B.** GST pulldown of CaMKII as in panel A using GST-actinin in the absence of nucleotides.

Figure 4. Oxidized calmodulin does not efficiently bind to CaMKII

A. Purified CaMKII was incubated with oxidized (OX, +) or control (OX, -) calmodulin agarose in the presence of 2 mM Ca²⁺ (Ca²⁺, +) or 2 mM EDTA (Ca²⁺, -), washed extensively, and eluted in SDS sample buffer. Supernatant (Super) and eluate (Pellet) were separated by SDS-PAGE and analyzed by Ponceau stain. B. Control (top) or oxidized (bottom) calmodulin agarose was incubated with rat brain lysate in the presence of 2 mM Ca²⁺, washed extensively, and eluted in 200 μ l fractions with 5 mM EDTA. Fractions were separated by SDS-PAGE and analyzed by Western blot for CaMKII. C. Rat brain lysate was passed over reduced or oxidized CaM agarose. Resin was washed with 0.5 M NaCl buffer and eluted with EDTA. Total elution from reduced (-) and oxidized (+) resin is shown. M: broad range markers.