Current Practice

Anaemia in the Tropics

E. J. WATSON-WILLIAMS,* M.B., M.R.C.P.

[WITH SPECIAL PLATE FACING PAGE 43]

Brit. med. J., 1968, 4, 34-38

Anaemia is one of the commonest manifestations of ill-health in the tropics. The causes are numerous, and anaemia in most patients is contributed to by several independent and inter-dependent factors. Moreover, the importance of different nutritional, environmental, and genetic factors differs widely from area to area. It is impossible, therefore, to give a detailed review of the diagnosis and management of anaemia in a brief article. Instead I have attempted to describe how the practitioner with limited resources may assess the different factors responsible for anaemia and plan rational management of his patients.

Pattern of Anaemia in the Tropics

Anaemia may be secondary to acute infection. It is usually slight, with normochromic red cells, but if the nutritional status is marginal infection may precipitate folic-acid deficiency or less commonly iron deficiency. In severe infection a haemolytic element is seen, and this is particularly likely in persons deficient of glucose-6-phosphate dehydrogenase.

In infection with tuberculosis, brucellosis, and amoebiasis anaemia is the rule, and responds only when the primary condition is treated. Nutritional deficiencies of iron and folic acid are commonly associated with these infections and require treatment.

Malaria always results in increased red cell destruction, and severe anaemia of rapid onset follows infection of the non-immune with *Plasmodium falciparum*. In areas where *P. falciparum* malaria is stable anaemia attributable to the parasite is seen in children but not in adults unless their immunity is altered by residence abroad, previous suppressive therapy, splenectomy, or disease of the reticuloendothelial system. During pregnancy, particularly in primiparae, there is a change in malarial immunity, and severe anaemia with parasitaemia may occur. In areas where *P. falciparum* malaria is unstable little immunity develops, and anaemia in adults as well as in children may be attributable to infection.

Undernutrition is widespread and particularly evident in infancy for several reasons. Because of maternal deficiency the newborn may have reduced iron, folic-acid, and vitamin-B₁₂ stores. In many areas weaning is delayed beyond five months; breast milk provides insufficient haematinics as well as calories and protein. Further, nutritional requirements are increased by bacterial and protozoal infections. The anaemia in kwashiorkor may become obvious only after the initial treatment, and is usually due to a combination of iron and folic-acid deficiency. In some cases there is a haemolytic anaemia as a direct result of protein deficiency. Anaemia is usually slight in the marasmic child, but becomes severe once the diet improves in

calories. This is a result of lack of reserve haematinics to provide for the return to normal blood volume. Curiously the malnourished child, although susceptible to bacterial infections, is less prone to malaria than the healthy infant. Once the child has reached 10 years of age the anaemia of malnutrition, although still common, is usually less severe. After puberty the male generally meets his nutritional requirements, though reserves of iron and folic acid may remain low. On the other hand, the female, as a result of menstrual loss, may remain iron deficient and is likely to enter pregnancy already anaemic. Folic-acid deficiency may be added after the 20th week of gestation, and the blood will show the features of dimorphic anaemia.

It is evident that with this background of poor nutritional status any increase in blood loss or destruction cannot readily be compensated for by increased haemoglobin and red cell formation. Thus haemolysis is usually complicated by folic-acid deficiency, and hookworm infection or schistosomiasis results in iron-deficiency anaemia. The amount of blood loss in hookworm infection is directly proportional to the numbers of adult worms, and the frequency and severity of the resultant anaemia are related to this and to the iron content of the diet.

Whereas iron deficiency and folic-acid deficiency are common throughout the tropics, nutritional vitamin-B₁₂ deficiency is limited to those whose diet is practically devoid of animal protein. It is exceptionally rare in the Negro race, but is not uncommon among Asians, who for religious and social reasons may be strict vegetarians.

In addition to these environmental and socio-economic factors there are two types of genetic variations in high frequency in the tropics that are responsible for anaemia. These are the abnormal haemoglobin disorders and thalassaemia and glucose-6-phosphate dehydrogenase deficiency, which are discussed later.

Diagnosis of Anaemia

Clinical.—An experienced physician should have no difficulty in recognizing the anaemic patient, but even severe anaemia is easily missed by those unused to tropical practice. The history volunteered may be bizarre and is often no more than a complaint of "general bodily weakness" of uncertain duration. Pallor is masked by skin pigmentation, but is detected by examining the nail beds and mucous membranes. Some specific clinical signs that are helpful in differential diagnosis are listed below and in Table I.

The *tongue* may be red in nutritional iron deficiency, is often smooth at its edges in folic-acid deficiency, and is invariably so in vitamin-B₁₂ deficiency.

The *skin* is a good guide to nutritional status. The appearance in kwashiorkor is diagnostic. In iron deficiency it is often dull, and there may be some loss of pigmentation. In

* Senior Lecturer in Clinical Haematology, University of Manchester, the Royal Infirmary, Manchester 13.

TABLE I.—Physical Signs in the Diagnosis of Anaemia in the Tropics

	Fever	Cardio- megaly	Oedema	Tongue	Skin	Spleen	Liver	Jaundice	Urine			
									Albumin	Uro- bilinogen	Bilirubin	Haemo- globin
Iron deficiency ..					Koilony- chia							
Nutritional ..		(+)	+	Red	Dull		(+)					
Hookworm ..		(+)	+		Dull		(+)					
Haemolytic	(+)	++			Leg ulcers	{ Children ++ Adults — }	{ ++ + }	+		++	(+)	
Sickle cell ..		++					+	++		++		
Thalassaemia ..							(+)	++		++		+
G-6-PD. deficiency ..	+	+					+	+		+		
Other inherited ..					(Petechiaie)		+	+		+		(+)
Autoimmune ..	+	(+)					+	+		+	(+)	(+)
Malaria ..	++						+	+	+	++	(+)	(+)
Megaloblastic	(+)	+	+	(Smooth)	(Pigment)*	+	+	(+)		+		
Folic-acid deficiency		+		Smooth	Pigment*	(+)		+		+		
Vitamin-B ₁₂ deficiency												
Secondary to		(+)	(+)		Dull	(+)	+		+			
Tuberculosis ..	+	(+)					+		+			
Amoebiasis ..	++						+		+			
Schistosomiasis ..						++	(+)					Red cells
Acute infection ..	++					+	(+)	(+)	+			
Aplastic ..	(+)	+			Petechiaie		(+)					Red cells
Leukaemia etc. ..	++	(+)			Petechiaie	++	+	(+)	+			

(+) Sometimes present. * i.e., Increased pigmentation.

megaloblastic anaemia there may be increased pigmentation, and if due to vitamin-B₁₂ deficiency this is particularly marked on the palmar surface.

Nails.—When shoes are not worn koilonychia is usually more obvious in the feet than the hands.

Abdomen.—In infants massive splenomegaly suggests sickle-cell anaemia, thalassaemia, or malaria, whereas slight enlargement is common in acute infections. In adults splenomegaly suggests schistosomiasis, leukaemia, portal hypertension, visceral leishmaniasis, brucellosis, or malaria in unstable areas; it is occasionally seen in inherited haemoglobin disorders and autoimmune haemolytic anaemia. Massive unexplained splenomegaly with anaemia is discussed separately. The spleen is never palpable in uncomplicated iron deficiency.

Enlargement of the liver is seen in the same conditions as splenomegaly, as well as in tuberculosis, amoebiasis, and kwashiorkor, and persists into adult life in sickle-cell anaemia.

Fever.—Mild hyperpyrexia may occur in severe anaemia from any cause except iron deficiency. Higher constant levels are seen in falciparum malaria, typhoid and related fevers, amoebiasis, tuberculosis, and other infections, as well as in sickle-cell infarctive crises. Intermittent and relapsing fever suggest malaria, amoebic hepatitis, or one of the specific relapsing fevers.

Urine.—Albumin is present in the urine in most fevers, but not in uncomplicated anaemia. Urobilinogen in excess and without bilirubin indicates haemolysis. Haemoglobin shows that haemolysis is intravascular and suggests malaria, glucose-6-phosphate dehydrogenase deficiency, or autoimmune haemolytic anaemia. The presence of unexplained macroscopic blood may be due to sickle-cell trait or disease. The deposit should be examined for pus cells and schistosome ova. The specific gravity is always below 1115 in sickle-cell anaemia.

Faeces.—In hypochromic anaemia the faeces should be examined for blood (Hematest tablets are suitable) and for hookworm ova. In jaundice the colour of the faeces may help to distinguish between biliary obstruction and haemolysis.

Examination of the Blood

The estimation of haemoglobin and examination of a stained blood film should be a minimum requirement in every patient in tropical practice. While ancillary help may be available, the physician should be in the habit of examining films himself, as his knowledge of the clinical condition allows him more

easily to determine the significance of the blood findings. The notes that follow are intended to help the inexperienced rather than the expert. They are therefore rather dogmatic.

Estimation of Haemoglobin

Simple screening to detect severe and moderate anaemia is sometimes attempted with the Tahlquist papers. These are subject to many errors and generally are no more valuable than the examination of the mucous membranes. If a screening test is desired the use of solutions of copper sulphate is much more reliable. The principle is that the higher the haemoglobin level the greater the density of a drop of blood. The concentration of the copper sulphate solutions is varied, so that at any chosen haemoglobin level the blood will either float or sink. Although it is possible to construct a series of solutions that would enable the haemoglobin level of every specimen to be estimated, it is usually sufficient to have only two solutions adjusted so that one will detect haemoglobin below 10 g./100 ml. (68%) and the other below 7 g./100 ml. (48%). (Standards A and B.)

A stock solution of S.G. 1100 is made by placing 510 g. of crystalline copper sulphate (CuSO₄.5H₂O) into a 4-litre bottle. Distilled water is added in three amounts, each of 1,006 ml. (if the temperature is 25° C.). The stock solution is diluted: 47.2 ml. made up to 100 ml. with water gives an S.G. of 1048 (Standard A) and 41.2 ml. gives an S.G. of 1042 (B). It is convenient to put 20 ml. into each of five screw-capped universal containers and to use each bottle for no more than 20 tests. A drop of blood taken either direct from a finger prick or from a venous specimen with a teated pipette is dropped vertically from a height of 1 cm. into the solution. After a few seconds the blood will either rise or fall. If it rises the haemoglobin is less than the standard.

This method is the most reliable of the simple screening tests, but depends on the concentration of plasma protein as well as the haemoglobin. If the plasma protein is less than 6.5 g./100 ml., or more than 8.0 g./100 ml., errors of about 5% will occur.

Accurate haemoglobin measurement demands the use of an electric colorimeter or an instrument such as the M.R.C. grey-wedge photometer. These can easily be adapted to work with a 2-volt accumulator, and are therefore usable in any location. Although there are several differing haem pigments which have been used, the cyanmethaemoglobin method is recommended. The modification described overcomes the disadvantage of having to wait 10 minutes for full conversion of the pigment. The solution of cyanide used is not sufficiently concentrated to be a hazard. The great advantage of this method is the availability of a commercial standard with good keeping qualities that makes it possible to check the colorimeter each day. In humid climates the photo-electric cell may have a shortened life. It is advisable to keep a few spare cells in a desiccator.

Method

Stock Reagent.—Potassium ferricyanide 200 mg., potassium cyanide 50 mg., potassium dihydrogen phosphate 140 mg., 1.0 ml. Nonidet P40 (Shell Chemical Co.) in 1 litre of distilled water. Keep in a dark bottle at 4–20° C.

Procedure.—Make a dilution of 20 cu. mm. of whole blood in 4 ml. (1/201) or 5 ml. (1/251) of the stock solution. Read against a water blank at 540 nm. or using an Ilford 625 filter. The concentration of haemoglobin in g./100 ml. is determined from a standard curve derived from serial dilutions of a standard solution (B.S.3985) supplied by C. Davis Keeler Limited, B.D.H. Ltd., or Diagnostic Reagents Ltd. The greatest error is due to unskilled use of the 20 cu. mm. pipette. This can be circumvented by using the "unopette" system (Becton Dickinson Ltd.); number 5857 unopette contains 5 ml. diluent and automatically dilutes 0.020 ml. of blood.

The Blood Film

Use only new slides straight from the suppliers' package. A small drop of blood is placed in the centre about $\frac{1}{2}$ cm. from one end. A second slide is used to make the film. Its short edge is placed across the slide and just in front of the drop of blood. Held at an angle of 45° it is first pulled back, so that the blood spreads nearly to its edges, and then pushed forward at a steady speed to the end of the slide. Several films should be made and the best chosen for staining. A well-made film covers about two-thirds of the slide, has an even tail without ridges, and both edges free.

Of the Romanovsky stains Leishman's is the simplest to use. A suitable alcoholic solution and tablets to make buffered water are available from laboratory suppliers. Undiluted stain is used for 30–45 seconds and then twice the volume of distilled water (pH 6.8) is added and mixed. After 10 minutes the stain is tipped off and the film washed with distilled water until slightly pink. It is air dried and the back of the slide is cleaned with cotton-wool moistened with methanol.

The film should be scanned initially with a low-power objective. The number of white cells is assessed and an area chosen for high-power examination with the oil-immersion lens. This should be in the centre just behind the thinnest portion (the tail) of the film. The red cells should be discrete, equally spaced, and fill nearly the whole field. Normal red cells have only a faint tinge of haemoglobin in the centre (central pallor). If this is not present the cells are either spherocytic and small or just released from the marrow and bluish in colour (polychromasia). If there is no central colour at all the cells are hypochromic. If hypochromasia is marked the anaemia is due to iron deficiency or thalassaemia, which can be distinguished by the presence of target cells, polychromasia, nucleated red cells, and perhaps punctate basophilia in the latter. Moderate or slight hypochromasia is seen in anaemia due to acute or chronic infection and in some of the abnormal haemoglobin

diseases, as well as in iron-deficiency anaemia. If target cells are abundant thalassaemia has to be distinguished from homozygous haemoglobin C disease, homozygous haemoglobin E disease, and sickle-cell anaemia and their mixed syndromes. In the case of sickle-cell anaemia a careful search will always reveal typical sickled cells as well as polychromasia. *P. falciparum* malaria should be suspected if there is polychromasia with some target cells and spherocytes. It is confirmed by finding the ring parasites and typical lumps of brownish-gold pigment in the monocytes. Macrocytic red cells with a number of pear-shaped cells suggest megaloblastic anaemia, and this should be confirmed by the presence of polymorphonuclear leucocytes with hypersegmented nuclei (more than 30% having five or more lobes). Megaloblastic anaemia in the tropics is usually associated with iron deficiency, and the film then shows a mixture of hypochromic microcytic and normochromic macrocytic cells, but hypersegmented neutrophils will always be present and make the diagnosis easier. The typical blood film appearances in different conditions are summarized in Table II and are illustrated in Figures 1–8, Special Plate.

Further Investigation.—Although the blood film appearances are usually sufficiently characteristic to suggest the diagnosis and correct treatment of anaemia, before treatment is started it is good practice to take specimens of 10 ml. of clotted and of anticoagulated blood so that confirmatory investigations can be made. Often this will mean sending the blood to a well-equipped laboratory. For this purpose the vacutainer system (Becton Dickinson) is ideal. The specimens are sterile and suitable for most investigations even after 72 hours at a high ambient temperature. Suggested investigations are listed in Table III. Even visual inspection of the plasma is useful. It

TABLE III.—Suggested Additional Investigation of Anaemia in the Tropics

HYPOCHROMIC ANAEMIA		
Serum iron	If low	Examine faeces for hookworm ova and blood Examine for schistosomiasis. Examine bone marrow for stainable iron. Consider acute or chronic infection. Screen for thalassaemia or abnormal haemoglobin.
	If normal or raised	
DIMORPHIC ANAEMIA OR MACROCYTIC ANAEMIA		
Bone marrow	If megaloblastic	Serum B ₁₂ and folate. Gastric acidity after histamine. Screen for abnormal haemoglobin.
HAEMOLYTIC ANAEMIA		Estimate red cell glucose-6-phosphate dehydrogenase Consider malaria, bartonellosis, leishmaniasis, etc. Malignancy, including reticulo-endothelial diseases. Tuberculosis, amoebiasis, etc. Acute or chronic leukaemia. Sickle-cell crisis.
LEUKAEMOID PICTURE		
PANCYTOPENIA		
Bone marrow for		Megaloblastic anaemia. Regenerative crisis in sickle-cell anaemia, etc. Acute leukaemia. Aplastic anaemia.

is characteristically pale in protein deficiency and in iron deficiency, icteric in megaloblastic and haemolytic anaemia, and brown or red if haemolysis has been intravascular.

TABLE II.—Diagnosis of Anaemia in the Tropics from Blood Film

Anisocytosis	Poikilocytosis	Hypochromasia	Polychromasia	Macrocytosis	Microcytosis	Target Cells	Sickle Cells	Spherocytes	Nucleated Reds	Other Features	White Cells	Platelets	Probable Diagnosis
+	+	++	(+)	(+)	++	(+)					HS	+	Iron deficiency Iron deficiency with blood loss Iron deficiency with folic-acid deficiency
+	+++	++	++	(+)	++	++	+		++	P.B.	(+)		Thalassaemia Sickle-cell anaemia Other abnormal haemoglobin G-6-P.D. deficiency Familial spherocytosis Autoimmune haemolytic anaemia Malaria
+	++	+	++	(+)	+	+		+	++	Rings	+	(-)	
+		+	++	+		(+)			+		+	+	Acute haemorrhage Acute infection Amoebiasis Tuberculosis
+	+		+	(+)					(+)		+++	(-)	Chronic leukaemia Acute leukaemia Aplastic anaemia

(+) Sometimes present or increased. (-) Sometimes reduced. P.B., punctate basophilia. Rings, malaria parasites. P, malaria pigment. H.S., hypersegmented neutrophils.

Treatment of Anaemia

Treatment consists of the appropriate management of any infection or other precipitating disease, an adequate diet, the use of blood transfusion when necessary, and the correct use of iron, folic acid, and vitamin B₁₂.

Blood transfusion carries particular hazards in the tropics because of the increased risk of infection, the high incidence of virus hepatitis, and the risk of transmitting malaria and other blood-borne diseases. Further screening of donors should include the determination of haemoglobin type and the presence of glucose-6-phosphate dehydrogenase deficiency. For these reasons it should be reserved for life-saving situations, and should never be used without thought. Most cases of anaemia in adults can be treated without transfusion, but severe anaemia in infants and during pregnancy can often be managed successfully only where there are good facilities for blood transfusion. A rough guide to the indications is given in Table IV.

TABLE IV.—Indications for Blood Transfusions in the Tropics

Age	Disease	Haemoglobin Level
3/12-3 years ..	Anaemia with heart failure	< 7 g./100 ml.
	Anaemia with infection	< 5 g./100 ml.
	Anaemia uncomplicated	< 3 g./100 ml.
Above 3 years	Acquired haemolytic anaemia	< 7 g./100 ml.
	Anaemia with infection	< 5 g./100 ml.
	Anaemia with heart failure	
	Inherited haemolytic anaemia	
	Thalassaemia	< 3 g./100 ml.
	Hypoplastic anaemia	
In pregnancy ..	Anaemia with heart failure	
	Anaemic crisis in sickle-cell disease	
At term ..	Anaemia	< 8 g./100 ml.
Post partum ..	Anaemia with heart failure	< 7 g./100 ml.
	Sickle-cell crisis	

Iron.—There are so many oral preparations that selection other than by cost is invidious. For adults ferrous sulphate (B.P.) 200 mg. thrice daily is generally well tolerated and usually entirely sufficient. Where there is doubt about absorption this can be improved by administering it between meals or together with ascorbic acid 800 mg. once daily. For children the B.P.C. ferrous sulphate mixture, containing 25 mg. of iron in 10 ml., may be preferred. The dose should contain 3 mg. of iron per kg. daily.

Parenteral iron undoubtedly has attractions in tropical practice. It is known to have been given, and the risk of children swallowing a large number of iron tablets left about the dwelling by the patient is avoided. Imferon, given as a total dose infusion, is particularly suitable, as only one attendance at hospital is necessary. *However, before prescribing parenteral iron it is important to be sure that the anaemia is due to iron deficiency, and it must never be used in patients with sickle-cell disease or thalassaemia.* Imferon contains 50 mg. of iron per ml. and the amount required is calculated from the formula

$$\text{Fe} = \frac{W (\text{HbD} + 1)}{250}$$

[Fe is iron in g., W is the weight in kg., and HbD the haemoglobin deficit in g. per 100 ml.]

The cause of the iron deficiency must be treated. In many cases in the tropics this will be due to hookworm infestation. Treatment is with bephenium hydroxynaphthoate 5 g. daily for three days or tetrachlorethylene 0.1 ml./kg. (maximum 3 ml.) after a 12-hour fast.

Folic Acid.—Folic-acid deficiency is the cause of the great majority of instances of megaloblastic anaemia in the tropics. Because of the frequency of deficiency it is good practice to give it as a prophylactic to all infants with infection or malnutrition, to all patients with haemolytic anaemia from any cause, and during pregnancy. Five milligrams by mouth daily is always sufficient, but up to 20 mg. daily may be used in the treatment of anaemia in pregnancy. It can be given parenterally if desired.

Vitamin B₁₂.—Vitamin-B₁₂ deficiency is rare except among strict vegetarians. Nutritional deficiency is adequately cor-

rected by oral doses of 50 µg. daily. For pernicious anaemia it should be given in doses of 1 mg. monthly by intramuscular injection, though if this is impracticable 300 µg. daily by mouth is effective.

Prevention of Anaemia in the Tropics

While complete prevention of anaemia would require the eradication of malaria, hookworm, and other common diseases, as well as a greatly increased standard of nutrition, a considerable amount of morbidity could be saved by the prophylactic use of oral iron and folic acid in infancy and pregnancy. In malarious areas suppressive therapy is advisable throughout pregnancy and during any illness in infancy. If confined to these situations it will not materially retard the development of immunity.

Sickle-cell Anaemia

Sickle-cell anaemia is due to the homozygous inheritance of sickle-cell haemoglobin. It is a common clinical problem in West, Central, and East Africa, and the Middle East, where up to 1.5% of children are affected. Lower frequencies occur among African descendants in the Caribbean and South and North America, and in certain areas around the Mediterranean and in the Indian subcontinent. Presentation is usually between the ages of 6 months and puberty with haemolytic anaemia or painful bone crisis. Painful crises are due to intravascular sickling of red cells, which may occur in any site. Thus in addition to bone pain episodes of abdominal, pulmonary, and intracranial infarction occur. A particular hazard is that an abdominal crisis may closely mimic an acute surgical emergency, but anaesthesia and operation result in a high mortality. The diagnosis is easily made from the examination of a blood film. If sickled cells are seen it is imperative that operation is resisted and the child treated as for sickle-cell crisis.

Management.—Sickle-cell crisis is treated by rest in bed, analgesia, antibiotics, and intravenous isotonic saline, 10 ml./kg. initially, followed by 10 ml./kg. every 12 hours for at least 48 hours. The antibiotic of choice depends on the circumstances: usually tetracycline 250 mg. six-hourly is used, but if there is evidence of osteomyelitis chloramphenicol in the same dose is to be preferred because of the likelihood of salmonella infection. In pulmonary crisis intramuscular penicillin and streptomycin may be preferable. Folic acid 15 mg. daily should be given, and blood transfusion is indicated if the haemoglobin drops to 3 g./100 ml.

In addition to infarctive episodes anaemic crises occur. These are of three types: megaloblastic, due to folic acid deficiency; aregenerative with temporary bone-marrow failure; and acute sequestration crisis with enlargement of liver and spleen. The management is by blood transfusion if the haemoglobin is below 3 g./100 ml. and folic acid.

Between crises efforts should be towards the prevention of infection with malaria suppression and perhaps oral penicillin V, and continued folic acid 5 mg. daily.

Other forms of sickle-cell disease due to inheritance of sickle-cell haemoglobin, together with thalassaemia or haemoglobin C, D Punjab, etc., result in clinically similar but less severe disease. All forms of sickle-cell disease are a particular hazard during pregnancy.

Thalassaemia

Homozygous β-thalassaemia presents as severe hypochromic anaemia between 6 months and 10 years of age. Other forms of thalassaemia are much less significant and often not diagnosed until adulthood. Blood transfusion is indicated to maintain the haemoglobin above 3 g./100 ml. Folic acid 5 mg. daily and malaria prophylaxis if indicated should be given

permanently. Splenectomy may be beneficial, but requires careful consideration. Iron, particularly parenterally, is contraindicated.

Glucose-6-phosphate Dehydrogenase Deficiency

Deficiency of this enzyme is inherited as a sex-linked recessive, and occurs in high frequency (about 10% to 20%) in African males and results in haemolytic anaemia following the taking of certain drugs and during severe infections. Deficiency also occurs in about 10% of males of Sardinian, Greek, Oriental Jewish, Filipino, Chinese, and Hindu Indian extraction. In addition to susceptibility to drug-induced haemolysis affected members of these populations may also have neonatal haemolytic anaemia and favism. Very occasionally a haemolytic anaemia without precipitating cause is due to this enzyme deficiency. Thus it must be considered in the differential diagnosis of haemolytic anaemia in the tropics. A simple spot screening test is described by Fairbanks and Beutler,¹ but if more than a few tests are made the B.C.B. decolorization of Motulsky and Campbell-Kraut² will be found more convenient. Haemolysis is intravascular, and therefore haemoglobinuria is usual, and there is a consequent risk of anuria. The treatment consists of stopping the offending drug, giving copious fluids (unless there is anuria), and blood transfusion if the haemoglobin drops to below 7 g. per 100 ml. Generally recovery is spontaneous and rapid. Drugs that cause haemolysis include

primaquine and other 8 amino-quinolines, sulphones and sulphonamides, phenacetin, P.A.S., aspirin, and water-soluble analogues of vitamin K. Detailed information, including diagnostic techniques, is given by Motulsky.³

Idiopathic Tropical Splenomegaly Syndrome

In many areas of the tropics gross splenomegaly without obvious cause is seen in a small number of adults. It is invariably associated with anaemia and usually with hepatomegaly and lymphocytic infiltration of liver and bone marrow. The aetiology is obscure, but in Africa at least there is increasing evidence that it is related to an abnormal immune response to malaria. It is possible that in non-malarious areas it is related to a similar abnormal response to other protozoal infections. Treatment with proguanil 100 mg. daily is effective in malarious areas and should be continued permanently. Elsewhere splenectomy may be advisable. The subject will be considered at greater length in a later article in this series.

REFERENCES

- ¹ Fairbanks, V. F., and Beutler, E., *Blood*, 1962, 20, 591.
- ² Motulsky, A. G., and Campbell-Kraut, J. M., *Proceedings of the Conference on Genetic Polymorphisms and Geographic Variations in Disease*, ed. B. S. Blumberg, 1961. New York.
- ³ Motulsky, A. G., *Abnormal Haemoglobins in Africa*, a C.I.O.M.S. symposium, ed. J. H. P. Jonxis, 1965. Oxford.

MEDICINE TODAY

Drop Attacks

"*Medicine Today*" is the television series for doctors produced by the B.B.C. Advice on the preparation of the programme is given by the Association for the Study of Medical Education.

The programme on B.B.C. 2 on 1 October was on the subject of investigation of strokes. Printed below is an article prepared with the help of expert contributors to complement the television programme, which will be repeated on B.B.C. 1 on 9 October about 11.37 p.m.

Completely unheralded disturbances in which patients abruptly fall to the ground are graphically termed "drop attacks." Most patients claim that they are unaware of falling, but all too vividly remember the moment of impact. Broken teeth, spectacles, and bones often testify to the suddenness of onset and speed of falling. If consciousness is impaired it may be for a brief instant only. Usually the patient is a middle-aged woman who has no other symptoms; fear of recurrence of these catastrophic upsets when crossing the road or out in the street often provokes understandable feelings of anxiety and insecurity.

A careful history paying particular attention to possible precipitating factors and circumstantial details normally excludes the many conditions in which falls may occur. For example, frail elderly patients often collapse unexpectedly; mild confusion and inattention, particularly in unfamiliar surroundings, are usually responsible. Those disabled by osteo-arthritis of the hip and knee occasionally complain that their legs let them down and that they frequently trip, but the significance of orthopaedic disorders is usually fairly evident. Any of the numerous causes of syncope, whether from hypotensive drugs or psychogenic faints, may cause patients to fall to the ground, but the characteristic premonitory symptoms and the duration

of loss of consciousness usually leave little doubt as to what really happened. Similarly, vertigo of sudden onset may cause patients to stagger and fall, but again little diagnostic confusion is aroused.

During attacks of insufficiency of the vertebrobasilar arterial system transient disorders of brain stem function frequently occur. True drop attacks may be associated with vertigo, tinnitus, ataxia, nystagmus, and dysaesthesiae, particularly of the face and tongue. However, it must be emphasized that in this common—or at least commonly diagnosed—condition drop attacks do not frequently occur. Certain epileptic patients, usually children, in addition to the more obvious manifestations of epilepsy experience drop attacks in which consciousness does not seem to be impaired. They are usually capable of picking themselves up from the ground immediately. These epileptic variants have been termed "akinetic" epilepsy. The pathophysiology of drop attacks is unknown, but their occurrence in brain stem disturbances suggests a paroxysmal instability of mechanisms controlling posture and muscular tone.

In cataplexy, vigorous emotional reactions, such as a hearty laugh, excitement, or anger, may provide an abrupt buckling of the legs and the patient may crash to the ground. The curious circumstances preceding the attack reveal the true diagnosis, and most of these unfortunate patients are also distressed by narcolepsy—a tendency to excessive sleep. A comprehensive list of the possible organic explanations why patients may fall abruptly would be tedious, but mention might be made of three uncommon but remedial disorders. Intermittent complete heart block may cause standing patients to crash to the ground; unstable subluxations of the atlanto-axial joint, as in advanced rheumatoid arthritis, may be associated with abrupt

E. J. WATSON WILLIAMS: ANAEMIA IN THE TROPICS

Photomicrographs of blood films showing diagnostic features. All $\times 1,000$.

1. Normal red cells. Note the central pallor.

2. Iron deficiency anaemia. The cells are very hypochromic and smaller than normal.

3. Megaloblastic anaemia. The cells are normochromic and macrocytic. Note the polymorphonuclear leucocyte has six nuclear lobes.

4. Dimorphic anaemia. The cells are microcytic and mostly hypochromic. The hypersegmented nucleus of the polymorphonuclear leucocyte shows that there is folic-acid or vitamin-B₁₂ deficiency as well as iron deficiency.

5. Sickle-cell anaemia. There are target cells, elongated cells with straight edges and one typical sickle cell.

6. The numerous target cells suggest an abnormal haemoglobin disorder and the spherocytes indicate that haemoglobin C is very likely present. This is from haemoglobin SC disease.

7. Thalassaemia. There is very marked anisocytosis and poikilocytosis. The majority of the cells are hypochromic, some target cells are present and there is a nucleated red cell.

8. *Plasmodium falciparum* malaria. The group of monocytes show typical dense spots of pigment. There are four ring parasites in the red cells.