¹A.G. Hayes, P.J. Birch, N.J. Hayward, M.J. Sheehan, H. Rogers, M.B. Tyers, *D.B. Judd, *D.I.C. Scopes & *A. Naylor

4 It is concluded that the seven novel compounds described are all potent and selective agonists for the κ -opioid receptor.

¹ Author for correspondence.

and DAMGO and U69593 in mouse vas deferens). In antagonist experiments, the tissues were incubated for 30 min with antagonists before the agonist concentration-response curve was re-determined. In the experiments with GR89696 and norbinaltorphimine, several antagonist concentrations were used in each tissue and the resulting dose-ratios were used to obtain a pA_2 value and Schild slope (Arunlakshana & Schild, 1959). Values from different tissues were then averaged together. In all the other antagonist experiments only a single antagonist concentration was used and a pK_B value was calculated from the equation: $pK_B = \log(\text{dose-ratio} - 1) - \log[\text{antagonist}]$.

In vivo experiments

Mouse experiments Male CRH mice (17–25 g) were used. Agonist drugs were administered subcutaneously (s.c.) in a dose volume of 10 ml kg^{-1} or orally in a dose volume of 20 ml kg^{-1} and 30 and 60 min respectively before testing. Naloxone was administered s.c. at 20 min before testing. Antinociception was measured by the acetylcholine-induced abdominal constriction test in the mouse (details of which are given in Hayes *et al.*, 1987a). Potential side-effects were assessed by studying drug effects in the rotarod test, on respiration rate and on gastrointestinal propulsion, all in the mouse (details of the measurement of these parameters are given in Hayes & Tyers, 1983).

Individual tests were carried out with dose groups of 6 mice. Either data from one individual experiment were used, or data were combined from 2 experiments i.e. $n = 12$. Each dose was randomised between cages and animals and drug solutions were colour coded such that the operators were unaware of which treatment the animals were receiving. The methods of Finney (1978) were used to determine regression slopes, linearity and parallelism for dose-response curves and also antinociceptive activities (ED_{50} values) and potency ratios, with 95% confidence limits.

Urine output of the rat Male PVG rats (125–200 g) were used. They were deprived of food overnight on the eve of an experiment and were water-loaded (25 ml kg^{-1}) 10 min before s.c. administration of the opioid agonist and/or antagonist, which were given in a dose volume of 4 ml kg^{-1} . On injection,

rats were placed in individual stainless steel metabolism cages and voided urine was collected in glass separators. Cumulative urine volumes were recorded hourly for up to 6 h; during this test period rats were deprived of food and water.

Drugs

Doses of salts refer to the weight of parent compound. The following drugs were used – GR38414 (1-[(3,4-dichlorophenyl)acetyl]-2-(1-pyrrolidinylmethyl) piperidine fumarate), GR89696 (methyl 4-[(3,4-dichlorophenyl)acetyl]-3-(1-pyrrolidinylmethyl)-1-piperazinecarboxylate fumarate), GR103545 ((R)-methyl 4-[(3,4-dichlorophenyl)acetyl]-3-(1-pyrrolidinylmethyl)-1-piperazinecarboxylate fumarate), GR45809 (8-[(3,4-dichlorophenyl)acetyl]-7-(1-pyrrolidinylmethyl)-1,4-dioxo-8-aza[4.5]spirodecane hydrochloride), GR91272 (5-[(3,4-dichlorophenyl)acetyl]-4,5,6,7-tetrahydro-4-[(3-hydroxy-1-pyrrolidinyl)methyl]furo[3,2-c]pyridine hydrochloride), GR44821 (1-[(3,4-dichlorophenyl)acetyl]-2-[(3-oxo-1-pyrrolidinyl)methyl]piperidine maleate) and GR86014 (2-[(3,4-dichlorophenyl)acetyl]-1,2,3,4-tetrahydro-1-(1-pyrrolidinylmethyl)-5-isoquinolinol maleate); all these compounds were synthesised in the Medicinal Chemistry Department, Glaxo Group Research Ltd., Ware. Ethylketocyclazocine (EKC) methane sulphonate (Sterling-Winthrop), morphine hydrochloride (Macfarlan-Smith), fentanyl citrate (Janssen), [D-Ala², D-Leu⁵] enkephalin (DADLE, Bachem), [D-Ala², N-MePhe⁴, Gly(ol)⁵] enkephalin (DAMGO, Bachem), naloxone hydrochloride (Sterling-Winthrop), naltrexone hydrochloride (Endo), 16-methyl cyprenorphine (M8008, synthesized in the Medicinal Chemistry Department, Glaxo Group Research Ltd., Ware), U69593 (5 α ,7 α ,8 β -(–)-N-methyl-N-(7-(1-pyrrolidinyl)-1-oxaspiro(4,5)dec-8-yl)benzeneacetamide) (Upjohn), norbinaltorphimine (norBNI, synthesized in the Medicinal Chemistry Department, Glaxo Group Research Ltd., Ware).

Results

Isolated tissue experiments

The data from the isolated tissue experiments are shown in Table 1. All of the compounds were extremely potent, full

Table 1 Effects of novel κ -agonists in the field-stimulated rabbit vas deferens, rat vas deferens and hamster vas deferens *in vitro*

Compound	Potency in isolated tissue (mean $IC_{50} \pm \text{s.e.mean}$) in nM		
	Rabbit vas deferens	Rat vas deferens	Hamster vas deferens
GR38414	10.6 ± 2.3 (6)	NT	NT
GR44821	0.2 ± 0.1 (3)	NSE at concentrations up to $10 \mu\text{M}$ (2)	
GR45809	0.1 ± 0.06 (3)	No agonist effect at concentrations up to $10 \mu\text{M}$	
		pK_B v DAMGO = 5.4 (2)	pK_B v DADLE = 5.2 (2)
GR86014	0.6 ± 0.4 (4)	$\left. \begin{matrix} 4,400 \\ 10,000 \end{matrix} \right\} (2)$	$\left. \begin{matrix} 4,200 \\ 12,000 \end{matrix} \right\} (2)$
GR89696	0.09 ± 0.03 (4)	No agonist effect up to $10 \mu\text{M}$	
		pK_B v DAMGO = 5.6 (2)	pK_B v DADLE = 6.1 (2)
GR103545	$\left. \begin{matrix} 0.02 \\ 0.02 \end{matrix} \right\} (2)$	No agonist effect up to $10 \mu\text{M}$	
		pK_B v DAMGO = 6.4 (2)	$\left. \begin{matrix} 1,900 \\ 160 \end{matrix} \right\} (2)$
GR91272	2.7 ± 2.3 (3)	NSE at concentrations up to $10 \mu\text{M}$ (2)	
U69593	$\left. \begin{matrix} 300 \\ 540 \end{matrix} \right\} (2)$	No agonist effect up to $10 \mu\text{M}$	
		pK_B v DAMGO = 5.2 (4)	NSE at concentrations up to $10 \mu\text{M}$

NT = not tested, NSE = no agonist effect and no antagonism of [D-Ala², N-MePhe⁴, Gly(ol)⁵]enkephalin (DAMGO) (RVD) or [D-Ala², D-Leu⁵]enkephalin (DADLE) (HVD). n values are given in parentheses.

Table 2 *In vivo* activity of novel κ -agonists in comparison with U69593

Compound	<i>ED</i> ₅₀ in abdominal constriction (95% confidence limits) (mg kg ⁻¹)	Rotarod <i>ED</i> ₅₀ (95% confidence limits) (mg kg ⁻¹)	Depression of respiration rate (mg kg ⁻¹)	Inhibition of gut propulsion (mg kg ⁻¹)	Effect on urine output (mg kg ⁻¹)
GR38414	0.03 s.c. (0.02–0.05) 1.2 p.o. (0.8–1.6)	2.7 s.c. (0.3–17)	Shallow DRC at 0.1–19 s.c. <i>E</i> _{max} = 28%	Shallow DRC at 0.1–10 s.c. <i>E</i> _{max} = 46%	Diuresis at 0.08–6 s.c.
GR44821	0.06 s.c. (0.03–0.09) 1.9 p.o. (1.3–3)	0.3 s.c. (0.002–1)	Shallow DRC at 0.3–3 s.c. <i>E</i> _{max} = 18%	Shallow DRC at 0.3–3 s.c. <i>E</i> _{max} = 46%	Diuresis at 0.01–2 s.c.
GR45809	0.001 s.c. (0.0003–0.002)	NT	NT	NT	NT
GR86014	0.003 s.c. (0.001–0.004)	NT	NT	NT	NT
GR89696	0.0005 s.c. (0.0003–0.0009) 0.04 p.o. (0.02–0.12)	0.052 s.c. (0.04–0.008)	Shallow DRC at 0.0004–0.1 s.c. <i>E</i> _{max} = 16%	Shallow DRC at 0.0004–0.1 s.c. <i>E</i> _{max} = 46%	Diuresis at 0.008–0.12 s.c.
GR103545	0.00025 s.c. (0.00012–0.00045) 0.012 p.o. (0.005–0.035)	0.022 s.c. (0.017–0.028)	NSE at doses up to 1 s.c.	Shallow DRC at 0.004–0.25 s.c. <i>E</i> _{max} = 53%	Diuresis at 0.001–0.03 s.c.
GR91272	0.002 s.c. (0.001–0.004)	0.05 s.c. (0.012–0.099)	NT	NT	Diuresis at 0.08–1.2 s.c.
U69593	0.14 s.c. (0.09–0.22)	1.2 s.c. (0.8–1.8)	NSE at doses up to 6 s.c.	Shallow DRC at 0.2–6 s.c. <i>E</i> _{max} = 28%	Diuresis at 2.5 and 10 s.c.

DRC = dose-response curve; NT = not tested; For all values $n > 6$.

agonists for inhibiting twitch responses in the rabbit vas deferens. The effects of (R,S)-GR89696 and its (R)-enantiomer, GR103545, in this tissue were antagonized by naloxone with mean pK_B values of 7.7 ± 0.04 ($n = 4$) and 7.5 ($n = 2$) respectively, indicating that these effects are indeed mediated via κ -receptors. None of the compounds showed any effects at μ -receptors in the rat vas deferens or at δ -receptors in the hamster vas deferens at concentrations less than $1 \mu M$, indicating that they are all very selective κ -agonists. Furthermore, GR89696 inhibited twitch responses of the mouse vas deferens with an IC_{50} of 0.12 ± 0.03 nM ($n = 8$), an effect which was antagonized by the κ -antagonist, norBNI, with a mean pA_2 value of 9.92 (slope = 0.95, $n = 2$); the mean pK_B for antagonism of U69593 by norBNI was 10.4 ($n = 2$) and for antagonism of DAMGO was 7.8 ($n = 2$).

In vivo experiments

The *in vivo* activities of the seven novel κ -opioid agonists are shown in Table 2. Their profiles of activity in the whole animal are very similar to that of the known κ -agonist U69593. Thus, they produced very potent antinociceptive

activity in the mouse abdominal constriction test, with GR103545 being the most potent ($ED_{50} = 0.25 \mu g kg^{-1}$ s.c.). The antinociceptive effect of GR38414 in the mouse was antagonized by naloxone (Table 3), at doses very similar to those necessary to antagonize the effect of the κ -agonist U69593 and considerably higher than those needed to antagonize the effects of the μ -agonist morphine. All the compounds produced marked sedative effects, as noted by visual observation and quantified with the mouse rotarod test. They produced very little depression of respiration rate or inhibition of gut propulsion in the mouse, any effects seen being of low maximum amplitude. The compounds which were tested for effects on urine output in the rat all produced a marked diuresis. The diuretic effect at higher doses was often preceded by an initial antidiuretic effect, a property that was also shared by U69593 (Figure 2). In an attempt to investigate the mechanism of this antidiuretic effect, its reversal by the opioid antagonists, naltrexone and M8008, was studied. Naltrexone, $5 mg kg^{-1}$ s.c., reduced both the initial decrease and the subsequent increase in urine output produced by U69593, indicating that both are mediated via opioid receptors. M8008 produced a small diuretic effect on its own, but did not antagonize the antidiuretic effects of U69593 or GR89696 at a

Table 3 Naloxone reversal of the antinociceptive effects of morphine, U69593 and GR38414 in the mouse abdominal constriction test

Naloxone dose (mg kg ⁻¹ s.c.)	Potency ratio for naloxone antagonism (95% confidence limits)		
	Morphine	U69593	GR38414
6	—	40.2 (24.1–29.3)	56.8 (35.3–92.9)
2	—	17.8 (10.3–29.6)	32.0 (19.0–54.8)
0.6	93.6 (38.6–232)	6.8 (4.0–11.3)	17.3 (10.7–28.1)
0.2	28.8 (11.8–71)	—	—
0.07	10.4 (4.3–25.8)	—	—

Figure 2 Reversal of the effects of (a and b) U69593, (c) fentanyl and (d) GR89696 on urine output in the rat by the opioid antagonists, naltrexone and M8008. U69593 was given at 10 mg kg^{-1} s.c., GR89696 at 0.03 mg kg^{-1} s.c., fentanyl at 0.9 mg kg^{-1} s.c., naltrexone at 5 mg kg^{-1} s.c. and M8008 at 4 mg kg^{-1} s.c. Each point represents the mean value calculated from at least 5 different rats; vertical lines show s.e.mean. (○) Represents vehicle-treated animals, (●) represents animals treated with antagonist alone, (▲) represents animals treated with agonist plus antagonist.

dose, 4 mg kg^{-1} s.c., which completely abolished the decrease in urine output produced by the μ -agonist, fentanyl.

Discussion

The novel compounds described in this paper were all extremely potent agonists in the rabbit vas deferens, a tissue which contains opioid receptors mainly of the κ -type (Oka *et al.*, 1981; Hayes & Kelly, 1985). However, the compounds showed very little activity in the rat vas deferens, a tissue which contains predominantly μ -opioid receptors (Smith & Carter, 1986; Sheehan *et al.*, 1988), and very little activity in the hamster vas deferens, a tissue which contains predomi-

nantly δ -opioid receptors (McKnight *et al.*, 1985; Miller & Shaw, 1985; Sheehan *et al.*, 1986). These data from isolated tissues thus suggest that the compounds are very potent and selective κ -opioid agonists. This was confirmed for one of the compounds, GR89696, in the mouse vas deferens in that its agonist effect was antagonized by the κ -selective antagonist, norBNI, with a pA_2 value within the range previously obtained for its action at κ -receptors (Birch *et al.*, 1987) and very similar to that obtained against U69593, but about 2 orders of magnitude higher than that obtained against the μ -agonist, DAMGO.

The potency and κ -agonist selectivity of the compounds was confirmed in the experiments *in vivo*. All of the compounds were potent antinociceptive agents, with GR103545 being the most potent κ -agonist described so far. In common with other κ -agonists, the compounds were generally highly depressant in the mouse. Sedation is also known to occur with κ -agonists in man (Peters *et al.*, 1987) and may be accompanied by psychotomimetic effects, although the latter are less well documented and do not appear to be detected in animals. GR89696 and GR103545 appeared to show a greater separation between sedative and antinociceptive doses in the mouse than the other compounds. However, this separation is not as good in other species (e.g. GR89696 has identical ED_{50} values in the rat paw pressure and rat rotarod tests – unpublished observations). The compounds produced very little respiratory depression or inhibition of gastrointestinal motility; the lack of such effects with κ -agonists is well documented in the literature (Tavani *et al.*, 1983; Porreca *et al.*, 1984; Hayes *et al.*, 1985b).

The compounds described here all produced large increases in urine output in the water-loaded rat. Such diuretic effects of κ -agonists have been extensively described previously (Leander, 1983; Slizgi *et al.*, 1984) and are thought to be due primarily to the presence of κ -receptors in the posterior pituitary, whose function is to inhibit the release of vasopressin (Bicknell *et al.*, 1987; Oiso *et al.*, 1988). Our own compounds and U69593 each produced an initial decrease in urine output before the large increase. Such decreases in urine output have generally been associated with activation of μ -receptors (Leander, 1984; Hayes *et al.*, 1987b), but this seems unlikely to account for the effect seen with these κ -agonists, as the anti-diuretic effect, unlike that of the μ -agonist fentanyl, was not antagonized by the μ -antagonist M8008. The antidiuretic effect of U69593 was reduced by naltrexone and is therefore opioid in origin; the most likely explanation is that it is a κ -receptor-mediated effect and may, for example, be a manifestation of the marked behavioural depression seen with these compounds.

In summary, we have described here a novel series of highly potent and selective κ -agonists. These compounds should be useful probes for ascertaining further the functional significance of the κ -opioid receptor.

We would like to thank Mrs Susan Harrison, Miss Jane Burrige and Mrs Rebecca Sargent for their excellent technical assistance, Mrs J. Lloyd for synthesizing the norBNI, and Dr A. McElroy for synthesizing the M8008.

References

- ARUNLAKSHANA, O. & SCHILD, H.O. (1959). Some quantitative uses of drug antagonists. *Br. J. Pharmacol.*, **14**, 48–58.
- BICKNELL, R.J., CHAPMAN, C. & ZHAO, B.-G. (1987). Opioid receptor subtypes mediating inhibition of oxytocin, vasopressin and noradrenaline release from the rat neurohypophysis *in vitro*. *J. Physiol.*, **338**, 9P.
- BIRCH, P.J., HAYES, A.G., SHEEHAN, M.J. & TYERS, M.B. (1987). Norbinaltorphimine: antagonist profile at κ -opioid receptors. *Eur. J. Pharmacol.*, **144**, 405–408.
- COSTELLO, G.G., MAIN, B.G., BARLOW, J.J., CARROLL, J.A. & SHAW, J.S. (1988). A novel series of potent and selective agonists at the opioid κ -receptor. *Eur. J. Pharmacol.*, **151**, 475–478.
- COWAN, A. & GMEREK, D.E. (1986). *In vivo* studies on κ -opioid receptors. *Trends Pharmacol. Sci.*, **7**, 69–72.
- FINNEY, D.J. (1978). *Statistical Method in Biological Assay*. 3rd Edition, London: Charles Griffin and Co.
- HAYES, A.G. & TYERS, M.B. (1983). Determination of receptors that mediate opiate side effects in the mouse. *Br. J. Pharmacol.*, **79**, 731–736.
- HAYES, A.G. & KELLY, A. (1985). Profile of activity of κ -receptor agonists in the rabbit vas deferens. *Eur. J. Pharmacol.*, **110**, 317–322.
- HAYES, A.G., SHEEHAN, M.J. & TYERS, M.B. (1985a). Determination of the receptor selectivity of opioid agonists in the guinea-pig ileum and mouse vas deferens by use of β -funaltrexamine. *Br. J. Pharmacol.*

- macol.*, **86**, 899–904.
- HAYES, A.G., SKINGLE, M. & TYERS, M.B. (1985b). Effect of β -funaltrexamine on opioid side-effects produced by morphine and U-50, 488. *J. Pharm. Pharmacol.*, **37**, 841–843.
- HAYES, A.G., SHEEHAN, M.J. & TYERS, M.B. (1987a). Differential sensitivity of models of antinociception in the rat, mouse and guinea-pig to μ - and κ -opioid receptor agonists. *Br. J. Pharmacol.*, **91**, 823–832.
- HAYES, A.G., SKINGLE, M. & TYERS, M.B. (1987b). Evaluation of the receptor selectivities of opioid drugs by investigating the block of their effect on urine output by β -funaltrexamine. *J. Pharmacol. Exp. Ther.*, **240**, 981–988.
- LAHTI, R.A., MICKELSON, M.M., MCCALL, J.M. & VON VOIGTLANDER, P.F. (1985). [3 H]-U69593 a highly selective ligand for the opioid kappa receptor. *Eur. J. Pharmacol.*, **109**, 281–284.
- LEANDER, J.D. (1983). A κ -opioid effect: Increased urination in the rat. *J. Pharmacol. Exp. Ther.*, **224**, 89–94.
- LEANDER, J.D. (1984). Effects of full and partial κ -agonists and μ -agonists on urine output of normally hydrated rats. *Neuropeptides*, **5**, 283–287.
- LEIGHTON, G.E., JOHNSON, M.A., MEECHAM, K.G., HILL, R.G. & HUGHES, J. (1987). Pharmacological profile of PD117302, a selective κ -opioid agonist. *Br. J. Pharmacol.*, **92**, 915–922.
- LORD, J.A.H., WATERFIELD, A.A., HUGHES, J. & KOSTERLITZ, H.W. (1977). Endogenous opioid peptides: multiple agonists and receptors. *Nature*, **267**, 495–499.
- MARTIN, W.R., EADES, C.G., THOMPSON, J.A., HUPPLER, R.E. & GILBERT, P.E. (1976). The effects of morphine and nalorphine like drugs in the nondependent and morphine-dependent chronic spinal dog. *J. Pharmacol. Exp. Ther.*, **197**, 517–532.
- McKNIGHT, A.T., CORBETT, A.D., MARCOLI, M. & KOSTERLITZ, H.W. (1985). The opioid receptors in the hamster vas deferens are of the δ -type. *Neuropharmacol.*, **24**, 1011–1017.
- MILLER, L. & SHAW, J.S. (1985). Characterisation of the δ -opioid receptor on the hamster vas deferens. *Neuropeptides*, **6**, 531–536.
- OISO, Y., IWASAKI, Y., KONDO, K., TAKATSUKI, K. & TOMITA, A. (1988). Effect of the opioid κ -receptor agonist U50488H on the secretion of arginine vasopressin. *Neuroendocrinology*, **48**, 658–662.
- OKA, T., NEGISHI, K., SUDA, M., MATUMIYA, T., INAZU, T. & UEKI, M. (1981). Rabbit vas deferens: a specific bioassay for opioid κ -receptor agonists. *Eur. J. Pharmacol.*, **73**, 235–236.
- PETERS, G.R., WARD, N.J., ANTAL, E.G., LAI, P.Y. & DEMAAR, E.W. (1987). Diuretic actions in man of a selective κ -opioid agonist: U-62,066E. *J. Pharmacol. Exp. Ther.*, **240**, 128–131.
- PORRECA, F., MOSBERG, H.I., HURST, R., HRUBY, V.J. & BURKS, T.F. (1984). Roles of μ , δ and κ opioid receptors in spinal and supraspinal mediation of gastrointestinal transit effects and hot-plate analgesia in the mouse. *J. Pharmacol. Exp. Ther.*, **230**, 341–348.
- SHEEHAN, M.J., HAYES, A.G. & TYERS, M.B. (1986). Pharmacology of δ -opioid receptors in the hamster vas deferens. *Eur. J. Pharmacol.*, **130**, 57–64.
- SHEEHAN, M.J., HAYES, A.G. & TYERS, M.B. (1988). Lack of evidence for ϵ -opioid receptors in the rat vas deferens. *Eur. J. Pharmacol.*, **154**, 237–245.
- SLIZGI, G.R., TAYLOR, C.J. & LUDENS, J.H. (1984). Effects of the highly selective κ -opioid, U-50,488, on renal function in the anaesthetized dog. *J. Pharmacol. Exp. Ther.*, **230**, 641–645.
- SMITH, C.F.C. & CARTER, A. (1986). Delta receptors in the rat vas deferens. *Arch. Int. Pharmacodyn.*, **284**, 181–192.
- TAVANI, A., GAMBINO, M.C. & PETRILLO, P. (1983). The opioid κ -selective compound U-50,488H does not inhibit intestinal propulsion in rats. *J. Pharm. Pharmacol.*, **36**, 343–344.
- VON VOIGTLANDER, P.F., LAHTI, R.A. & LUDENS, J.H. (1983). U-50, 488: a selective and structurally novel non- μ (κ) opioid agonist. *J. Pharmacol. Exp. Ther.*, **224**, 7–12.

(Received March 12, 1990)

Revised June 21, 1990

Accepted June 27, 1990)