Published in final edited form as:

Metabolism. 2007 February ; 56(2): 251–259.

Reverse cholesterol transport is regulated by varying fatty acyl chain saturation and sphingomyelin content in reconstituted high density lipoproteins

Philippe Marmillot, Sanket Patel, and M. Raj Lakshman

Lipid Research Laboratory, Veterans Affairs Medical Center, and The Departments of Medicine and Biochemistry and Molecular Biology, The George Washington University, Washington, DC

Abstract

Because phospholipid composition of HDL plays a vital role in its reverse cholesterol transport (RCT) function, we studied RCT in vitro (uptake and efflux) with reconstituted HDLs (rHDLs) containing phosphatidylcholine with fatty acids of increasing saturation levels (stearic, oleic, linoleic, linolenic), and without or with sphingomyelin. Uptake significantly increased from basal value when phosphatidylcholine component included up to 50% (%mol) of oleic or linolenic acid, but did not change with linoleic acid. Increasing oleic and linoleic acids to 100% (%mol) significantly decreased uptake, but increasing linolenic acid to the same value did not affect it. Sphingomyelin in rHDL significantly decreased uptake, but only with phosphatidylcholine containing unsaturated fatty acids, and not with saturated fatty acid. Efflux was not affected in a dose-dependent manner when oleic or linoleic acid content was increased, but was significantly increased with levels of linolenic acid up to 25% (%mol) in phosphatidylcholine, and was dramatically lowered with higher levels. Sphingomyelin in rHDL (phosphatidylcholine:sphingomyelin, 20:80, mol:mol) significantly increased efflux only with oleic or linoleic acid-containing rHDLs, compared to efflux without sphingomyelin. In conclusion, enrichment of phosphatidylcholine component up to 25% (%mol) as linolenic acid has a beneficial effect on RCT, while a higher percentage of it or other unsaturated fatty acids seems to be detrimental. Also, high sphingomyelin content decreases uptake with rHDL containing unsaturated fatty acids, whereas it increases efflux for rHDL containing oleic or linoleic acid. These results show for the first time the importance of sphingomyelin in RCT in a well-defined in vitro system.

Introduction

Excess cholesterol is removed from peripheral tissue by reverse cholesterol transport (RCT), a process in which cellular cholesterol is removed by high density lipoprotein (HDL) (i.e., cholesterol efflux) and transported to the liver (i.e., cholesterol uptake) for metabolic transformation into bile acid and steroid hormones (1–3). Two proteins have been shown to be involved in the cholesterol efflux process, the ATP binding cassette transporter 1 (ABCA1), and the scavenger receptor class BI (SR-BI). Both ABCA1 and SR-BI are expressed by macrophages and parenchymal liver cells. ABCA1 facilitates cellular cholesterol and phospholipids efflux, and SR-BI binds HDL with high affinity (4–7). Factors that affect RCT

Correspondence Dr. Raj Lakshman, Lipid Research Laboratory (151-T), Veterans Affairs Medical Center, 50 Irving Street NW, Washington, DC 20422, USA. Phone: 202-745-8462. Fax: 202-462-2006. E-mail: raj.lakshman@med.va.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

function of HDL are cholesterol and phospholipids content, lecithin cholesterol acyltransferase (LCAT) activity, and fatty acid saturation levels.

With regard to the latter factor, we showed in a previous investigation that the RCT function of HDL is affected by the saturation level of dietary fat intake. Our previous studies of oral feeding of alcohol diet for 8 weeks according to Lieber's method, and fish oil fat (only 2.8% of total dietary calories as omega-3 fatty acids (ω 3-FA); alpha-tocopherol 120 iu/L of diet) resulted in lower plasma and liver lipids (8). Furthermore, the same low level of dietary ω 3-FA restored the decreased apolipoprotein E (apoE) content in HDL due to ethanol. Also, we have shown that the presence of as low as 2.8% of total dietary calories as ω 3-FA in the alcohol diet can decrease the damage done by alcohol to the RCT mechanism, but also can enhance by several fold the ability of HDL to promote RCT, even in control rats (9). We also showed that there was no interaction between the effect of ω 3-FA and the effect of alcohol on hepatocytes in term of RCT function (9).

In addition to being modulated by HDL's fatty acid components, cholesterol efflux is positively correlated with sphingomyelin (SM) content in HDL in human with high serum HDL cholesterol (> 67.7 mg/dl) (10). Thus, factors that regulate local modification of HDL's phospholipid composition may have a large impact on RCT. It has been demonstrated that SR-BI-mediated free-cholesterol (FC) flux is very sensitive to the phosphatidylcholine (PC) and SM composition of HDL and that phospholipids composition of HDL alters the steady-state distribution of cholesterol between cells expressing SR-BI and HDL. Particularly, SM enrichment of HDL enhances the net efflux of cholesterol from SR-BI expressing COS-7 cells by decreasing influx of HDL cholesterol (11).

SM content of the cell membrane is thought to contribute to the maintenance of cellular cholesterol homeostasis because the cholesterol content of the cell membrane is positively correlated with SM content (12). The interaction between cholesterol and SM is the basis for the formation and maintenance of cholesterol/sphingolipid-enriched nano- and micro-domains (referred to as membrane "rafts") in the plane of plasma and other organelle (i.e., Golgi) membranes (13). SM is thought to bind cholesterol with high affinity and inhibit its efflux from the plasma membrane by preventing cholesterol desorption (14). In addition, SM prevents the exchange of cholesterol between the plasma membrane and intracellular pools (15). However, the role of SM in the function of circulating lipoproteins remains unclear. Numerous studies on reverse cholesterol transport reported how various well-defined plasma components, particularly reconstituted HDL particles with a variety of compositions, regulate cholesterol efflux from cells (16–18). However, there is no information on the combined role of the fatty acid saturation levels of the phospholipids, and the SM levels, when both are varying in individual HDL particles.

Thus, in the work presented herein, our objective was to investigate RCT function of HDL, namely cholesterol uptake and efflux, by varying both the saturation levels of the fatty acid chain of the PC component and the levels of SM in various reconstituted HDL species.

Methods

Materials

Materials were obtained from the following sources: Dulbecco's modified Eagle's medium (DMEM), glutamine, penicilline/streptomycine, Invitrogen (Carlsbad, CA); fetal calf serum (FCS), Life Technologies (Gaithersburg, MD); Bovine brain sphingomyelin, egg yolk phosphatidylcholine, Sigma-Aldrich (St. Louis, MO); [3 H]-cholesteryl oleate, Amersham Biosciences (Piscataway, NJ); Tissue culture plastic ware, Falcon (Lincoln, NJ); Wild type HepG2 cells and mouse macrophages J774, ATCC (Rockville, MD); Insulin, Eli Lilly

Company (Indianapolis, Inc); 1,2-Distearoyl-sn-Glycero-3-Phosphocholine (DSPC, 18:0), 1,2-Dioleoyl-sn-Glycero-3-Phosphocholine (DOPC, 18:1, n-9FA), 1,2-Dilinoleoyl-sn-Glycero-3-Phosphocholine (DLOPC, 18:2, n-6FA), 1,2-Dilinolenoyl-sn-Glycero-3-Phosphocholine (DLNPC, 18:3, n-3FA), Avanti Polar Lipids Inc. (Alabaster, AL).

Preparation of HDL protein fraction

HDL was isolated by sequential flotation from human pooled plasma (19) and its protein fraction was delipidated in 20 volumes of chloroform:methanol (1:1, vol./vol.) for 2 hours at 4°C, followed by centrifugation at 2,000 rpm (Beckman J6 centrifuge with 4.2 rotor). The HDL protein pellet was homogenized in 5 volumes of 25 mmol/l ammonium bicarbonate, 10 mmol/l sodium decyl sulfate, 0.1% (vol./vol.) β-mercaptoethanol and solubilized overnight at 4°C under gentle agitation. The HDL protein solution was then dialyzed in 25 mmol/l ammonium bicarbonate, 0.1% (vol./vol.) β-mercaptoethanol to remove sodium decyl sulfate, and its protein content measured using the Bradford's method with BSA as a standard (20). Apolipoprotein AI (ApoAI) content was determined by densitometry analysis of the HDL protein fraction separated by denaturing electrophoresis on a 4–20% (w/v) linear gradient acrylamide gel and stained with SYPRO® Ruby protein stain (Bio-Rad, Hercules, CA). Delipidated HDL protein fraction was used to reconstitute HDL as described below, taking into account its apoAI content and apoAI's molecular weight of 28,000 daltons.

Making of liposomes

The rHDL complexes were prepared by sodium cholate dispersion according to previously reported methods (21–24). All rHDL species were prepared with an initial PC(+/-SM):apoAI:CO ratio equals to 250:1.6:12.5. Various lipids (i.e., PC, SM, CO, [³H]CO) were mixed together from stock solutions in chloroform to obtain rHDL composition indicated in tables 1A and 1B, and then evaporated to dryness with a gentle flow of nitrogen or helium. Under these conditions PC:CO molar ratio was always kept at 20:1. Only those rHDL that would be used for cholesterol uptake assay with hepatocytes would contain radiolabeled cholesteryl oleate. Dispersion buffer (10 mmol/l Tris-HCl (pH 8.0), 154 mmol/l NaCl, 1 mmol/l NaN₃, 0.3 mmol/l EDTA) was then added to all lipid mixtures to obtain a final lipid concentration of 10 mg/ml. This was followed immediately by vortexing for 2 min, after which sodium cholate (30 mg/ml) was added to obtain a molar ratio lipids:sodium cholate of 1:1, followed by vortexing for 2 min and by 1 hour incubation at 4°C. Several cycles of 5-min sonication and incubation at 37°C were necessary to obtain clear liposome suspensions. ApoAI-containing protein fraction (5 – 10 mg/ml) was then added to the liposome suspension to obtain a molar ratio PC(+/-SM):apoAI of 156.25:1. Each sample mixture was then incubated for 1 hour at 4°C and dialyzed extensively against dispersion buffer after adjusting the volume to 0.5 ml with dispersion buffer. The homogeneity and size of the reconstituted lipoprotein complexes were assessed by size exclusion chromatography and native-PAGE, as described below, and by negative staining electron microscopy (25). Electron microscopy analysis of our rHDL preparation revealed characteristic disc shaped structures (average diameter: 15 nm), similar to those found in the literature for rHDL (22,26–28).

Analysis of rHDL

Chemical analysis was performed by measuring protein (20), triglycerides (29), phospholipid (30), sphingomyelin (31) and total cholesterol (32,33) content. Homogeneity and size analysis were performed by (a) size exclusion chromatography on a Biogel 0.5M column (27 × 1 cm) using an AKTA Purifier system (Pharmacia, Uppsala, Sweden) and (b) non-denaturing electrophoresis on a 8–25% gradient gel using PhastSystem (Pharmacia, Uppsala, Sweden) (34,35). Molecular weight estimation was performed by analyzing the gels with a FluorChem 8800 densitometer loaded with AlphaEaseFC software (Alpha Innotech Corporation, San

Leandro, CA). Rf values were exported into Excel spreadsheet software (Microsoft, Redmond, WA) and analyzed with the power or exponential regression trend line function. Values listed in the revised Tables 3A and 3B are those obtained with the best correlation coefficient.

Cell culture systems

For uptake experiments, HepG2 cells were maintained in DMEM supplemented with 2 mM L-glutamine, 1% pen/strep and 10% (vol./vol.) FCS, according to the instructions from the supplier. When needed, cells were subcultured and grown as monolayer in 6-well culture dishes by culturing 1×10^6 cells per well containing 4 ml culture medium, followed by incubation at 37°C in 5% CO₂ air humidified atmosphere. Confluence of the cell layers was observed after 36–48 hours. All experiments were performed in post confluent proliferating phase of cells.

For efflux experiments mouse macrophages J774, were maintained in DMEM supplemented with 2 mM L-glutamine, 1% pen/strep, 10% (vol./vol.) FCS, and 0.1 mU% insulin in 37°C in 5% CO₂ air humidified atmosphere. When needed, cells were split and grown as monolayer in 6-well culture dishes (1×10^6 cells per well in 4 ml culture medium). Confluence of the cell layers was observed after 24–36 hours. All experiments were performed in post confluent proliferating phase of cells.

Esterified cholesterol uptake assay from labeled rHDL

An adequate volume of dialyzed rHDL was mixed with DMEM (4 ml final volume) and added as a sterile solution (by filtration through 0.22 µm sterile filters) to each culture dish containing the confluent HepG2 cells. An aliquot (0.1 ml) of the incubation medium was analyzed for the total esterified cholesterol radioactivity added to each dish. The uptake of labeled esterified cholesterol by the cells was determined after 24 hours by counting the cell-associated radioactivity as follows: the cell medium was completely removed and, after thorough washing of the cells 3 times with PBS to remove the medium's radioactivity, the cells were dissolved in 1 ml PBS containing 0.1 % SDS and 1 % Triton X-100. The dissolved cells were mixed with 4 ml of Packard's Ultima Gold scintillation cocktail (Perkin Elmer, Wellesley, MA) and analyzed for radioactivity in a Beckman model LS-6500 liquid scintillation spectrometer.

Cholesterol Efflux Assay

Macrophages, confluent in 6-well plates, were first incubated with [³H]cholesteryl-oleate-labeled acetylated LDL (0.06 µCi and 115.5 µg of LDL protein per well) in 4 ml of culture medium for 24 h. After washing 3 times with PBS (pH 7.4), the macrophages were then incubated with 75 µg protein of indicated rHDL per well in culture medium without FCS for 6 hours. At the end of the incubation period, radioactivity was measured in the medium, as described above. Efflux was expressed as the percentage of the initial radioactivity loaded in the macrophages that is found in the medium after 6 hours. In these conditions, approximately 45% of labeled cholesteryl-oleate is hydrolyzed to free cholesterol, and more than 90% of the released cholesterol appeared as unesterified in the culture medium (36).

Statistical Analyses

The significance of the various effects was evaluated by a one-way analysis of variance (ANOVA) with post hoc Tukey's test.

Results

rHDL characterization

In order to carry out measurement of cholesterol transport with HDLs of various lipid compositions, we made reconstituted HDL particles (rHDL) from individual purified

components, namely, cholesteryl ester, phosphatidylcholine (PC), sphingomyelin (SM), and HDL protein fraction. Tables 1 and 2 summarize the composition of the various rHDL species used in this study.

For all rHDL species the molar ratio for total PC:apoAI:cholesteryl ester was 250:1.6:12.5. The protein fraction was obtained from delipidated human HDL that was isolated by sequential flotation, and apoAI content was determined by densitometry analysis of SDS-PAGE profile (results not shown). Analysis of the cholesterol to apoAI molar ratio showed that values differed from the calculated value of 7.8. As shown in Figure 1, increasing unsaturated PC content in rHDLs decreased their cholesterol content in a dose-dependent manner. It appears that adding linolenic-FA-containing PC (DLNPC, 18:3[n-3]) in rHDL minimized the loss of cholesterol, compared with linoleic-FA- (DLOPC, 18:2) or oleic-FA (DOPC, 18:1)-containing species. rHDL species with only saturated PC (DSPC, 18:0) had however a cholesterol:apoAI ratio higher than 7.8, namely 10.1. There was no correlation between cholesterol:apoAI ratio and particles' size.

In order to verify the homogeneity of freshly prepared rHDL species, their protein and cholesterol profiles were analyzed with species containing [³H]cholesteryl oleate. As shown in Figure 2A, a representative size exclusion chromatography (SEC) profile indicates that apoAI-containing rHDL elutes as a single peak that overlaps with [³H]cholesteryl oleate profile. The absorbance peak appearing in the early fractions (2 to 5) does not contain apoAI and does not appear to indicate the presence of a significant amount of protein, as shown by SDS-PAGE analysis of the eluted fractions. Also, cholesteryl ester is exclusively bound with apoAI-containing particles (Figure 2B).

After verifying the size homogeneity of each rHDL, we investigated size variation among the various rHDL species. Thus, size analysis was performed on various rHDL species by non-denaturing gel electrophoresis as described in the Experimental Procedures section. Figure 3A and Table 3A show that the rHDL species containing only saturated PC (i.e., DSPC) has the smallest size, namely 6.9 nm.

Analysis of the data listed in Table 3A indicates that higher levels of unsaturated FA slightly increase the size of rHDLs that contain DOPC (i.e., 18:1), but decrease the size of rHDLs that contain the highest level of DLOPC (i.e., 18:2), as well as the size of rHDLs that contain DLNPC (i.e., 18:3). Analysis of the data listed in Table 3B indicates that SM does not affect the size of rHDLs, except at its highest content (80%, mol%) with rHDL containing DLNPC.

Cholesterol uptake

As described in the following sections, we measured cholesterol uptake with rHDLs whose phospholipid composition had (a) PC with fatty acids of various levels of saturation, and (b) various levels of SM. Also, all rHDL species were radioactively labeled with [³H]cholesteryl oleate.

Effect of Fatty Acid Composition of PC—As shown in Figure 4, uptake of cholesterol in the hepatocyte system had a basal level when egg yolk phosphatidylcholine (EYPC) was used as sole source of phospholipid in rHDL. Uptake was the highest when rHDL PC component contained only saturated (18:0) fatty acid (DSPC only). When the proportions of DOPC (18:1) or DLNPC (18:3[n-3]) in rHDLs were increased up to 50% (molar %) of total PC content, uptake was significantly higher than basal value. However, the same increase in DLOPC (18:2) showed no change in cholesterol uptake. In contrast, uptake of cholesterol was significantly decreased below basal level when the proportion of DOPC or DLOPC in rHDL was increased to 100% (molar %) of total PC content. Interestingly, the same change in DLNPC did not change uptake level from basal value.

Effect of SM composition of PC—In this series of experiments, SM was added in increasing proportion into various rHDL species, while the ratio of saturated to unsaturated fatty acid was kept constant (namely, 3 to 1, as shown in Table 2). Results presented in Figure 5 show that increasing SM content dramatically decreased the uptake of rHDL species containing EYPC, or containing DOPC (18:1), DLOPC (18:2) and DLNPC (18:3[n-3]) ($P<0.01$, compare 20% and 80% SM), and to a lesser extent also decreased the uptake of rHDL containing DSPC (18:0) only ($P<0.05$, compare 20% and 40% SM). Also, uptake was the lowest when EYPC was the only PC component of rHDL.

Cholesterol efflux

Similar to the cholesterol uptake experiments described above, we measured cholesterol efflux with rHDLs whose PC composition had (a) fatty acids with various levels of saturation, and (b) various levels of SM, as described below. In this series of experiments, rHDL species were not radioactively labeled.

Effect of Fatty Acid Composition of PC—The most striking observation in the results presented in Figure 6 is the dramatic decrease of cholesterol efflux when rHDL DLNPC (18:3 [n-3]) content was increased from 50% to 100% ($P<0.01$). Importantly, when rHDL DLNPC (18:3[n-3]) content was only 25%, cholesterol efflux was significantly enhanced ($P<0.01$) when compared to basal value obtained with rHDL species whose sole source of phospholipids was EYPC. Other data from this set of experiments show that cholesterol efflux was not significantly affected, at least in a dose-dependent manner, whether DOPC (18:1) or DLOPC (18:2) was used to replace EYPC in rHDLs.

Effect of SM composition of PC—As described in the uptake experiment above, rHDL species with a constant ratio of saturated fatty acid (DSPC (18:0)) to unsaturated fatty acid (DOPC (18:1), DLOPC (18:2), or DLNPC (18:3[n-3])), namely 3 to 1, and an increasing proportion of SM were used to assay the efflux transport of cholesterol. Results are plotted in Figure 7 and show that SM does not significantly affect cholesterol efflux in the presence of rHDL species that contain EYPC, DSPC only, or DSPC/DLNPC (18:0/18:3[n-3]). However, SM increased significantly cholesterol efflux with rHDL species containing DSPC/DOPC (18:0/18:1), but only at the highest proportion of SM, namely 80% mol of PL. The most significant effect was obtained when efflux was carried out in the presence of rHDL containing DSPC/DLOPC (18:0/18:2), and there was a SM dose-dependent increase of cholesterol efflux. In summary, cholesterol efflux was significantly increased with DSPC/DOPC (18:0/18:1) or DSPC/DLOPC (18:0/18:2) in rHDL species whose PC:SM ratio was 20:80, compared to efflux with rHDLs without SM.

Discussion

Overall, these results show a dose dependent effect (positive and negative) of both unsaturated fatty acids and SM in the phospholipid component of rHDL on cholesterol uptake and cholesterol efflux. The effects of various HDL lipid components investigated in the present study on the RCT process are summarized in Figure 8.

Enrichment of rHDL phosphatidylcholine component with a limited amount of DLNPC (18:3 [n-3]), by up to 25% of the total phospholipids in molar terms, has a beneficial effect on cholesterol uptake (Figure 4). Any further increase in the DLNPC (18:3[n-3]) or other polyunsaturated fatty acids content seems to be detrimental for this function. These results are consistent with the observation that high levels of polyunsaturated fatty acids in the diet are associated with higher incidence of coronary artery disease (CAD), whereas lower amounts have cardioprotective effects, especially n-3FA.

Most studies on dietary oil supplement focus on the effect of the diet on lipid profiles. One work with monkeys described a direct correlation between HDL's acyl chain composition and the level of dietary oil saturation in various regimens (17). However, no subsequent effect on RCT was observed. In a recent study (37), HDL's composition (i.e., HDL-cholesterol) and RCT function were investigated in humans fed diets containing various oils defined by their level of saturated or unsaturated fatty acids. Results showed that a diet rich in n-3FA (i.e., 1.6% of total dietary calories as fish oil) improved RCT function. However, no analysis was made to determine HDL's FA composition. In the present study, HDL composition is well-defined and our results show that RCT is also increased in correlation with the addition of limited amounts of n-3FA in rHDL. Thus, we hypothesize that the improvement of RCT after dietary intake of fish oil is due to the enrichment of HDL with n-3FA. However, it is important to point out that the DLNPC used in the present study, although also containing n-3FA, is chemically different from EPA and EHA found in fish oil due to the difference in carbon chain length in (i.e., 18 carbons instead of 20 and 22, respectively).

When both RCT component are compared, our results show that cholesterol uptake by the liver is more sensitive to variation in the levels of unsaturated fatty acids in rHDL phospholipids composition than cholesterol efflux is. Indeed, cholesterol uptake is significantly increased from basal level by moderate enrichment of rHDL in DOPC (18:1) and DLNPC (18:3[n-3]), whereas efflux is significantly increased only with rHDL moderately enriched in DLNPC (18:3 [n-3]). Conversely, cholesterol uptake is significantly decreased from basal level when rHDL are highly enriched in DOPC (18:1) and DLOPC (18:2) (Figure 4), whereas efflux is significantly decreased only with rHDL highly enriched with DLNPC (18:3[n-3]) (Figure 6).

With regard to SM effect on RCT, we showed that cholesterol uptake is significantly decreased by high content of SM in rHDL. The reverse is true for cholesterol efflux, namely, high levels of SM in rHDL significantly increase it. These findings are in accordance with the known high affinity binding of SM with cholesterol, thus preventing its desorption from SM-rich rHDL in the uptake process, and promoting desorption from macrophage membrane in the efflux process (13).

Acknowledgements

This work was supported by a grant from NIAAA (R01 AA08149). The authors thank Dr. Robert Redman, Chief of Oral Pathology at the Veterans Administration Medical Center, Washington, DC, for his assistance with electron microscopy analysis of rHDL.

References

1. van der Velde AE, Groen AK. Shifting gears: liver SR-BI drives reverse cholesterol transport in macrophages. *J Clin Invest* 2005;115:2699–2701. [PubMed: 16200207]
2. Hill SA, McQueen MJ. Reverse cholesterol transport--a review of the process and its clinical implications. *Clin Biochem* 1997;30:517–525. [PubMed: 9399019]
3. Glomset JA. The plasma lecithin:cholesterol acyltransferase reaction. *J Lipid Res* 1968;9:155–167. [PubMed: 4868699]
4. Brousseau ME, Eberhart GP, Dupuis J, Asztalos BF, Goldkamp AL, Schaefer EJ, Freeman MW. Cellular cholesterol efflux in heterozygotes for tangier disease is markedly reduced and correlates with high density lipoprotein cholesterol concentration and particle size. *J Lipid Res* 2000;41:1125–1135. [PubMed: 10884295]
5. Lawn RM, Wade DP, Garvin MR, Wang X, Schwartz K, Porter JG, Seilhamer JJ, Vaughan AM, Oram JF. The Tangier disease gene product ABC1 controls the cellular apolipoprotein-mediated lipid removal pathway. *J Clin Invest* 1999;104:R25–31. [PubMed: 10525055]
6. Jian B, de la Llera-Moya M, Ji Y, Wang N, Phillips MC, Swaney JB, Tall AR, Rothblat GH. Scavenger receptor class B type I as a mediator of cellular cholesterol efflux to lipoproteins and phospholipid acceptors. *J Biol Chem* 1998;273:5599–5606. [PubMed: 9488688]

7. de la Llera-Moya M, Rothblat GH, Connelly MA, Kellner-Weibel G, Sakr SW, Phillips MC, Williams DL. Scavenger receptor BI (SR-BI) mediates free cholesterol flux independently of HDL tethering to the cell surface. *J Lipid Res* 1999;40:575–580. [PubMed: 10064746]
8. Lakshman MR, Chirtel SJ, Chambers LL. Roles of omega 3 fatty acids and chronic ethanol in the regulation of plasma and liver lipids and plasma apoproteins A1 and E in rats. *J Nutr* 1988;118:1299–1303. [PubMed: 3193248]
9. Marmillot P, Rao MN, Liu QH, Chirtel SJ, Lakshman MR. Effect of dietary omega-3 fatty acids and chronic ethanol consumption on reverse cholesterol transport in rats. *Metabolism* 2000;49:508–512. [PubMed: 10778877]
10. Fournier N, Paul JL, Atger V, Cogny A, Soni T, de la Llera-Moya M, Rothblat G, Moatti N. HDL phospholipid content and composition as a major factor determining cholesterol efflux capacity from Fu5AH cells to human serum. *Arterioscler Thromb Vasc Biol* 1997;17:2685–2691. [PubMed: 9409243]
11. Yancey PG, de la Llera-Moya M, Swarnakar S, Monzo P, Klein SM, Connelly MA, Johnson WJ, Williams DL, Rothblat GH. High density lipoprotein phospholipid composition is a major determinant of the bi-directional flux and net movement of cellular free cholesterol mediated by scavenger receptor BI. *J Biol Chem* 2000;275:36596–36604. [PubMed: 10964930]
12. Barenholz Y.; Gatt, S. Phospholipids. Hawthorne, JN.; Ansell, GB., editors. Elsevier Biomedical Press; Amsterdam: 1982 . p. 129-177.
13. Barenholz Y. Sphingomyelin and cholesterol: from membrane biophysics and rafts to potential medical applications. *Subcell Biochem* 2004;37:167–215. [PubMed: 15376621]
14. Lund-Katz S, Laboda HM, McLean LR, Phillips MC. Influence of molecular packing and phospholipid type on rates of cholesterol exchange. *Biochemistry* 1988;27:3416–3423. [PubMed: 3390441]
15. Slotte JP, Bierman EL. Depletion of plasma-membrane sphingomyelin rapidly alters the distribution of cholesterol between plasma membranes and intracellular cholesterol pools in cultured fibroblasts. *Biochem J* 1988;250:653–658. [PubMed: 3390137]
16. Davidson WS, Gillotte KL, Lund-Katz S, Johnson WJ, Rothblat GH, Phillips MC. The effect of high density lipoprotein phospholipid acyl chain composition on the efflux of cellular free cholesterol. *J Biol Chem* 1995;270:5882–5890. [PubMed: 7890719]
17. Gillotte KL, Lund-Katz S, de la Llera-Moya M, Parks JS, Rudel LL, Rothblat GH, Phillips MC. Dietary modification of high density lipoprotein phospholipid and influence on cellular cholesterol efflux. *J Lipid Res* 1998;39:2065–2075. [PubMed: 9788253]
18. Zhao Y, Sparks DL, Marcel YL. Effect of the apolipoprotein A-I and surface lipid composition of reconstituted discoidal HDL on cholesterol efflux from cultured fibroblasts. *Biochemistry* 1996;35:16510–16518. [PubMed: 8987984]
19. Havel RJ, Eder HA, Bragdon JH. The distribution and chemical composition of ultracentrifugally separated lipoproteins in human serum. *J Clin Invest* 1955;34:1345–1353. [PubMed: 13252080]
20. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248–254. [PubMed: 942051]
21. Jonas A. Reconstitution of high-density lipoproteins. *Meth Enzym* 1986;128:553–582. [PubMed: 3724523]
22. Matz CE, Jonas A. Micellar complexes of human apolipoprotein A-I with phosphatidylcholines and cholesterol prepared from cholate-lipid dispersions. *J Biol Chem* 1982;257:4535–4550. [PubMed: 6802835]
23. Chen GC, Albers JJ. Characterization of proteoliposomes containing apoprotein A-I: a new substrate for the measurement of lecithin:cholesterol acyltransferase activity. *J Lipid Res* 1982;680–691. [PubMed: 6811681]
24. Laccotripe M, Makrides SC, Jonas A, Zannis VI. The carboxyl-terminal hydrophobic residues of apolipoprotein A-I affect its rate of phospholipid binding and its association with high density lipoprotein. *J Biol Chem* 1997;272:17511–17522. [PubMed: 9211897]
25. Forte TM, Nordhausen RW. Electron microscopy of negatively stained lipoproteins. *Meth Enzym* 1986;128:442–457. [PubMed: 2425222]

26. Lerch PG, Förtsch V, Hodler G, Bolli R. Production and characterization of a reconstituted high density lipoprotein for therapeutic applications. *Vox Sang* 1996;71:155–164. [PubMed: 8912458]
27. Nichols AV, Gong EL, Blanche PJ, Forte T. Characterization of discoidal complexes of phosphatidylcholine, apolipoprotein AI and cholesterol by gradient gel electrophoresis. *Biochim Biophys Acta* 1983;750:353–364. [PubMed: 6407531]
28. Gillotte KL, Davidson WS, Lund-Katz S, Rothblat GH, Phillips MC. Apolipoprotein A-I structural modification and the functionality of reconstituted high density lipoprotein particles in cellular cholesterol efflux. *J Biol Chem* 1996;271:23792–23798. [PubMed: 8798607]
29. McGowan MW, Artiss JD, Strandbergh DR, Zak B. A peroxidase-coupled method for the colorimetric determination of serum triglycerides. *Clin Chem* 1983;29:538–542. [PubMed: 6825269]
30. Bartlett GR. Phosphorus assay in column chromatography. *J Biol Chem* 1959;234:466–468. [PubMed: 13641241]
31. McGowan MW, Artiss JD, Zak B. Enzymatic colorimetry of lecithin and sphingomyelin in aqueous solution. *Clin Chem* 1983;29:1513–1517. [PubMed: 6872211]
32. Allain CC, Poon LS, Chan CS, Richmond W, Fu PC. Enzymatic determination of total serum cholesterol. *Clin Chem* 1974;20:470–475. [PubMed: 4818200]
33. Amundson DM, Zhou M. Fluorometric method for the enzymatic determination of cholesterol. *J Biochem Biophys Methods* 1999;38:43–52. [PubMed: 10078872]
34. Calabresi L, Vecchio G, Frigerio F, Vavassori L, Sirtori CR, Franceschini G. Reconstituted high-density lipoproteins with a disulfide-linked apolipoprotein A-I dimer: evidence for restricted particle size heterogeneity. *Biochemistry* 1997;36:12428–12433. [PubMed: 9376346]
35. Nichols AV, Krauss RM, Musliner TA. Nondenaturing polyacrylamide gradient gel electrophoresis. *Meth Enzym* 1986;128:417–431. [PubMed: 3724517]
36. Ho YK, Brown MS, Goldstein JL. Hydrolysis and excretion of cytoplasmic cholesteryl esters by macrophages: stimulation by high density lipoprotein and other agents. *J Lipid Res* 1980;21:391–398. [PubMed: 7381331]
37. Montoya MT, Porres A, Serrano S, Fruchart JC, Mata P, Gerique JA, Castro GR. Fatty acid saturation of the diet and plasma lipid concentrations, lipoprotein particle concentrations, and cholesterol efflux capacity. *Am J Clin Nutr* 2002;75:484–491. [PubMed: 11864853]

Abbreviations

CAD	Coronary Artery Disease
apoAI	apolipoprotein AI
FA	fatty acid
SM	sphingomyelin
PC	phosphatidylcholine
RCT	reverse cholesterol transport
VLDL	very low density lipoprotein
LDL	low density lipoprotein

HDL	high density lipoprotein
SR-BI	scavenger receptor class-B type I
EPA	eicosapentaenoic acid
EHA	docosahexaenoic acid
n-3FA	omega 3 fatty acid

Figure 1. Molar ratio of various rHDL species

At the end of the procedure for rHDL making, described in the Methods section, each species was analyzed for its total cholesterol and protein content. Values are displayed on top of the corresponding bars for each rHDL species. rHDL species composition is expressed as molar percentage of each PC. The horizontal dotted line represents the calculated ratio, based on the initial amount of cholesterol and protein added during rHDL making procedure.

Figure 2. Incorporation of cholesterol into apoAI-containing rHDL

A representative set of data for rHDL species containing 50% DLNPC (see Table 1) and labeled with [3H]cholesteryl oleate is shown. A. Size Exclusion Chromatography profile on BioGel 0.5M of 100 μ g protein of rHDL. Protein was monitored continuously spectrophotometrically at 280 nm (solid line), and radioactivity was measured by liquid scintillation on a 0.1-ml aliquot of each 2.5-ml fraction (thin line with open circles). B. An aliquot (12 μ l) of each fraction 4 to 10 was analyzed by 4–15% gradient SDS-PAGE, followed by protein staining with SYPRO® Ruby (BioRad Laboratories, Hercules, CA).

Figure 3. Size analysis of rHDL. A. rHDL without SM. B. rHDL with SM

Each indicated rHDL species (approximately 1 μ g protein) was subjected to electrophoresis on a 8–25% polyacrylamide native gel. Protein bands were revealed by coomassie staining. Particle size markers were determined by densitometry analysis using a Fluorchem8800 gel analyzer (Alpha Innotech Corp. San Leandro, CA) loaded with AlphaEasyFC software for calculations. A. Representative gel for rHDL species containing DOPC. In order to minimize the graph's complexity, rHDL composition is displayed as molar percentage of the varying PC species. Thus, 100% DSPC, DSPC only; 25% DOPC, DSPC:DOPC (187.5:62.5); 50% DOPC, DSPC:DOPC (75:75); and 75% DOPC, DSPC:DOPC (62.5:187.5). B: Representative gel for rHDL species containing DLOPC and SM. rHDL species' composition is displayed according to SM content only. Thus, 100:0 corresponds to DSPC:DOPC:SM (187.5:52.5:0), 80:20 to DSPC:DOPC:SM (150:50:50), 60:40 to DSPC:DOPC:SM (112.5:37.5:100), and 20:80 to DSPC:DOPC:SM (37.5:12.5:200). Data from densitometry measurements of the rHDL species' particle size are reported in Tables 3A and 3B. Particles size markers were bovine serum albumin (7.1 nm), lactate dehydrogenase (8.2 nm), thyroglobulin (9.2 nm), catalase (12.2 nm), and apoferitin (17.0).

Figure 4. Cholesterol uptake function of rHDL species that contain various amount of PC with increasing saturation levels in the FA chain

There are 5 major groups of rHDL, as indicated in the abscissa of graph. rHDLs belonging to “EYPC only” and “18:0 only” groups have only one source of PC, namely EYPC and DSPC (or 18:0 FA- containing PC), respectively. Other rHDL groups contain two sources of PC, namely DSPC in all of them and an “unsaturated PC”, i.e., DOPC (18:1), DLOPC (18:2), or DLNPC (18:3). These groups are divided into four rHDL subgroups, each with various molar ratios of DSPC and “unsaturated PC”, as indicated in the frame of the graph. Detailed rHDL compositions are given in Table 1. Data are the means $\pm$ SD of triplicate measurements. For all rHDL groups, bars with a different superscript letter are significantly different, $P < 0.05$ (ANOVA with post hoc Tukey’s test).

Figure 5. Cholesterol uptake function of rHDL species that contain various amounts SM and various saturation levels in the PC FA chain

There are 5 major groups of rHDL, as indicated in the abscissa of graph. rHDLs belonging to “EYPC only” and “18:0 only” groups have only one source of PC, namely EYPC and DSPC (or 18:0 FA- containing PC), respectively. Other rHDL groups contain two sources of PC, namely DSPC in all of them and an “unsaturated PC”, i.e., DOPC, DLOPC, or DLNPC, at a constant molar ratio. All groups are divided into four rHDL subgroups, each with various molar ratios of PC and SM, as indicated in the frame of the graph. SM’s content is expressed as a molar percentage of total phospholipids (PL). Detailed rHDL composition of rHDL is given in Table 2. Data are the means $\pm$ SD of triplicate measurements. Bars within each rHDL groups that do not have an identical superscript letter are significantly different, $P < 0.05$ (ANOVA with post hoc Tukey’s test).

Figure 6. Cholesterol efflux function of rHDL species that contain various amount of PC with increasing saturation levels in the FA chain
Cholesterol efflux methodology is described in details in the Materials and Methods section. Data are the means $\pm$ SD of triplicate measurements. Bars within each rHDL groups that do not have an identical superscript letter are significantly different, $P < 0.05$ (ANOVA with post hoc Tukey's test). Legend is otherwise identical to the legend of Figure 4.

Figure 7. Cholesterol efflux function of rHDL species that contain various amounts of SM and various saturation levels in the PC FA chain
 Cholesterol efflux methodology is described in details in the Materials and Methods section. Data are the means $\pm$ SD of triplicate measurements. Bars within each rHDL groups that do not have an identical superscript letter are significantly different, $P < 0.05$ (ANOVA with post hoc Tukey's test). Legend is otherwise identical to the legend of Figure 4.

Figure 8. Schematic representation of the interaction of various HDL phospholipids components with the RCT process

Liver is on the left side of the schematics, indicating cholesterol uptake, while a macrophage is represented on the right side, indicating cholesterol efflux. RCT is the transport of cholesterol by HDL from peripheral tissues (e.g., macrophages) to the liver. Fatty acid content of various PCs: DOPC, oleic acid (18:2); DLOPC, linoleic acid (18:2); DLNPC, linolenic acid (18:3 [n-3]). SM, sphingomyelin.

Table 1**Composition of rHDL species (without SM)**

Chemical composition of various phospholipids DSPC, DOPC, DLOPC, and DLNPC is given in the Materials and Methods section. EYPC: egg yolk phosphatidylcholine.

rHDL Species	Molar ratio
EYPC	EYPC only
DSPC (18:0)	DSPC only
DSPC:DOPC (18:0/18:1)	75:25
	50:50
	25:75
	0:100
DSPC:DLOPC (18:0/18:2)	75:25
	50:50
	25:75
	0:100
DSPC:DLNPC (18:0/18:3)	75:25
	50:50
	25:75
	0:100

Table 2**Composition of rHDL species (with SM)**

Chemical composition of various phospholipids DSPC, DOPC, DLOPC, and DLNPC is given in the Materials and Methods section.

rHDL Species	molar ratio	PC:SM molar ratio	DS:(DO, DLO, or DLN) molar ratio
DSPC:SM (18:0/SM)	PC only 200:50 150:100 50:200	PC only 80:20 60:40 20:80	DSPC only
DSPC:DOPC:SM (18:0/18:1/SM)	PC only 150:50:50 112.5:37.5:100 37.5:12.5:200	PC only 80:20 60:40 20:80	75:25
DSPC:DLOPC:SM (18:0/18:2/SM)	PC only 150:50:50 112.5:37.5:100 37.5:12.5:200	PC only 80:20 60:40 20:80	75:25
DSPC:DLNPC:SM (18:0/18:3/SM)	PC only 150:50:50 112.5:37.5:100 37.5:12.5:200	PC only 80:20 60:40 20:80	75:25

Tables 3A and 3B

Particle size of rHDL species

A. rHDL without SM. B. rHDL with SM. rHDLs were subjected to native gel electrophoresis, and sizes were measured by densitometry, as described in the Materials and Methods section. rHDL species in 3A and 3B, whose sizes are listed in grayed areas, have their corresponding electrophoretic profile displayed in Figures 3A and 3B, respectively.

A			B		
Size distribution of rHDL without SM			Size distribution of rHDL with SM		
rHDL species	% PC	nm	rHDL species	PC:SM ratio	nm
DSPC only	100% DSPC	6.9	DSPC:SM		
DSPC:DOPC			250:0	DSPC only	6.9
			200:50	80:20	6.9
			150:100	60:40	6.8
			50:250	20:80	6.9, 7.4
187.5:62.5	25% DOPC	9.1, 8.1	DSPC:DOPC:SM		
75:75	50% DOPC	9.3	187.5:62.5:0	100:0	6.9, 7.7
			150:50:50	80:20	6.9, 7.8
62.5:187.5	75% DOPC	9.3	112.5:37.5:100	60:40	7.7
0:250	100% DOPC	9.4	37.5:12.5:200	20:80	7.8
DSPC:DLOPC			DSPC:DLOPC:SM		
187.5:62.5	25% DLOPC	8.8	187.5:62.5:0	100:0	7.6
			150:50:50	80:20	7.6
75:75	50% DLOPC	9.1	112.5:37.5:100	60:40	7.6
62.5:187.5	75% DLOPC	9.0	37.5:12.5:200	20:80	7.8, 8.6, 8.9
0:250	100% DLOPC	7.5	DSPC:DLNPC:SM		
DSPC:DLNPC			187.5:62.5:0	100:0	7.4
187.5:62.5	25% DLNPC	8.7, 7.5	150:50:50	80:20	7.4
75:75	50% DLNPC	7.7, 6.3	112.5:37.5:100	60:40	7.4
			37.5:12.5:200	20:80	9.4
62.5:187.5	75% DLNPC	7.8, 6.4	EYPC:SM		
0:250	100% DLNPC	8.0, 6.4	250:0	EYPC only	8.9, 7.7
			200:50	80:20	8.9, 7.7
			150:100	60:40	9.0, 7.8
			50:250	20:80	8.9, 7.8