with typical large eosinophilic cytoplasmic inclusions, while the intervening placental tissue was normal.

Scrapings from one of the skin lesions, when cultured on the allantoic membrane of developing chick embryos, grew vaccinia virus.

COMMENT

The appearance of the foetus and its weight suggest that death had probably occurred rather later than the date of cessation of movements would indicate.

While nobody would dispute the overriding need for protection against smallpox in all contacts, this case demonstrates that the foetus is at risk from maternal vaccinia in mid-term. It is not an isolated example, for similar cases have been reported by Wielenga *et al.* (1961), Lynch (1932), MacDonald and MacArthur (1955), and Wiersum (1956). There are probably others that have not been recorded.

Thanks are due to Professor A. S. Duncan for permission to publish this case, to Mr. T. J. Cooke for the illustrations, and to Dr. A. Evans for the isolation of the vaccinia virus.

D. M. ENTWISTLE, M.B., B.S.

P. T. BRAY, M.R.C.P., D.C.H., B.Sc.

K. M. LAURENCE, M.A., M.B., Ch.B.

Departments of Gynaecology and of Child Health
of the Welsh National School of Medicine
at Llandough Hospital, Penarth, Glamorgan.

REFERENCES

- Dixon, C. W. (1962). *Brit. med. J.*, **1**, 1262.
Lynch, F. W. (1932). *Arch. Derm. Syph. (Chic.)*, **26**, 997.
MacDonald, A. M., and MacArthur, P. (1955). *Arch. Dis. Childh.*, **28**, 311.
Wielenga, G., van Tongeren, H. A. E., Ferguson, A. H., and van Rijssel, Th. G. (1961). *Lancet*, **1**, 258.
Wiersum, A. K. (1956). *Ned. T. Geneesk.*, **100**, 971.

"Painless" Intussusception

Although the diagnosis of intussusception in infants is frequently straightforward, cases do occur which present considerable difficulty. The serious results of delay in starting treatment are universally appreciated, hence early correct diagnosis is of particular importance. Virtually all authorities quote dramatic abdominal pain as the outstanding symptom of the disease, and the absence of this may well lead to dangerous temporizing unless such a possibility is kept in mind. Two such cases have recently been encountered.

CASE 1

A male infant aged 5 months was admitted to hospital at midday on February 14, 1961. Two days previously he had appeared listless and had a slight cough. He had refused his feeds and had passed a loose stool, neither blood nor mucus being obvious in it. About 5.30 p.m. on the day before admission he had produced a projectile vomit amounting to approximately a cupful, and he continued to vomit smaller amounts every half-hour up to 10 p.m. During this period he had one further loose motion, again without blood or mucus. Throughout the night he regurgitated mouthfuls of fluid at two-hourly intervals. Shortly after 9 p.m. there was further projectile vomit consisting of 2-3 oz. (60-90 ml.) of bilious fluid. There was nothing to suggest that he was suffering any pain apart from a tendency to whimper when he was being sick.

On examination he was fretful and somewhat dehydrated. The abdomen was normal in appearance, but a definite tumour was palpable in the right hypochondrium and the caecum could not be felt in the right iliac fossa. There was no guarding or suggestion of tenderness. Rectal examination revealed frank blood and mucus on the examining finger. A diagnosis of intussusception was made

and operation immediately undertaken. This confirmed the presence of an ileo-colic intussusception with the head extending to the middle of the transverse colon. It was reduced fairly easily, but gross thickening of the terminal 2 in. (5 cm.) of the ileum was noted. The post-operative convalescence was uneventful and he went home one week after operation.

CASE 2

A female infant just under 8 weeks old was admitted to hospital on May 7, 1961, after examination at home. She had begun to vomit the previous evening, and on the day of admission had brought up all feeds. Nevertheless she had appeared cheerful, and nothing was volunteered by her mother to suggest that she had been in obvious pain. Four stools of the typical "red-currant jelly" type were passed during the course of the illness.

On examination the child appeared rather pale, but there was no evidence of dehydration. The right iliac fossa was "empty" on palpation and in the right hypochondrium there was a vague sensation of a mass; the child seemed to resent pressure in this area. Blood and mucus were noted on the finger after rectal examination. An intussusception was diagnosed, and this was confirmed at operation. As in Case 1 this was of ileo-colic type, the intussusception having advanced as far as the hepatic flexure. Reduction proved extremely difficult, the serous coat of the ascending colon splitting at two or three points during this procedure. After reduction was completed these areas were repaired and the abdomen was closed. The post-operative course was free of complications and the child was discharged on May 14.

COMMENT

The traditional teaching of the symptoms of intussusception has always stressed the dramatic onset, featured chiefly by colicky abdominal pain resulting in the child crying and drawing up its knees at regular intervals (Ladd and Gross, 1941; Eckhoff, 1955). It is obvious from the description of the above cases that reliance on signs of acute pain can be very misleading and might well lead to serious delay in diagnosis. Eckhoff mentions the fact that vomiting is usually one of the later symptoms of the disease, but Smith (1954) holds that early vomiting is extremely common. In both Cases 1 and 2 vomiting was a predominating feature, but, apart from this, neither child appeared unduly upset at any stage of the illness. In Case 2 the passage of blood and mucus in the stool was suggestive of the nature of the condition, but diagnosis from gastro-enteritis was rendered more complicated by the unusually early age at which symptoms occurred: Ladd and Gross quoted 484 cases in which only one child was under 2 months old. It might be postulated that the degree of pain is proportional to the tightness of the intussusception. This, however, does not seem to be the case, as the impaction in Case 2 was sufficient to make reduction extremely difficult and hazardous. It is obvious, therefore, that the practitioner or surgeon must consider the diagnosis of intussusception in all cases of vomiting in infancy, irrespective of whether pain is evident: such children merit particularly careful assessment of the abdominal signs and should always be submitted to rectal examination.

RONALD P. CUMMING, M.B., F.R.C.S.Ed.,
Consultant Surgeon, Shetland Hospitals.

REFERENCES

- Eckhoff, N. L. (1955). *British Encyclopaedia of Medical Practice*, 2nd ed., **7**, 183. Butterworth, London.
Ladd, W. E., and Gross, R. E. (1941). *Abdominal Surgery in Infancy and Childhood*, p. 107. Saunders, Philadelphia and London.
Smith, R. (1954). *Progress in Clinical Surgery*, Series I, p. 101. Churchill, London.