IS BEHAVIOR IN TEMPORAL LOBE EPILEPSY DIFFERENT THAN IN OTHER EPILEPSIES? THE JURY IS OUT

Interictal Depression, Anxiety, Personality Traits, and Psychological Dissociation in Patients with Temporal Lobe Epilepsy (TLE) and Extra-TLE. Swinkels WA, van Emde Boas W, Kuyk J, van Dyck R, Spinhoven P. *Epilepsia* 2006;47(12): 2092–2103. **PURPOSE:** This study was performed to investigate the relation between symptoms of interictal depression, anxiety, personality traits, and psychological dissociation with the localization and lateralization of the epileptogenic zone in patients with partial epilepsy. **METHODS:** All patients were diagnosed according to the localization-related concept of the 1989 International League Against Epilepsy (ILAE) Classification of Epilepsies and Epileptic Syndromes, and the localization and lateralization of the epileptogenic zone was established by using the clinical criteria for noninvasive presurgical evaluation. This resulted in 67 patients with temporal lobe epilepsy (TLE) and 64 patients with extra-TLE. All patients were assessed on the various aspects of psychopathology by using a comprehensive battery of standardized diagnostic instruments. **RESULTS:** We did not find the hypothesized excess of psychiatric symptoms in patients with (mesial) TLE in comparison with patients with extra-TLE. We also found no differences between patients with the lateralization of epilepsy in the left versus the right hemisphere. **CONCLUSIONS:** TLE per se cannot be considered a risk factor in developing more or more severe symptoms of psychopathology in patients with partial epilepsy. Concomitant factors, such as the duration of epilepsy, seizure frequency, and frontal lobe dysfunction may play an additional role. Our findings support the hypothesis of a multifactorial explanation for the psychiatric symptoms in patients with epilepsy.

COMMENTARY

The report by Swinkels and colleagues provides further evidence that behavioral changes are not limited to patients with partial epilepsy whose seizure foci arise in the temporal lobe but also occur in patients with extratemporal foci. Nearly 90% of the extratemporal foci were in the frontal lobes. The study was moderately powered yet was not sufficient in size to identify small differences between mesial and laterobasal temporal lobe epilepsy (TLE) groups. Swinkels et al. found no differences between patients with mesial and laterobasal TLE, with TLE compared to extra-TLE, or with left versus right foci, across a variety of standard psychiatric inventories for depression, anxiety, and personality. Are patients with TLE different?

Controversy has endured regarding the specificity of behavioral changes in TLE as well as to the relative risk of such changes in patients with TLE as compared to patients with other epilepsy syndromes. The temporal lobes contain the amygdala and other limbic areas that modulate emotion, memory, and assign emotional valence to environmental stimuli. Acute and subacute temporal lobe disorders such as encephalitis, stroke, and tumor cause a wide spectrum of behavioral pathology, including irritability, aggression, mania, depression, paranoia, and psychosis. As Geschwind suggested, it would be expected that

seizure foci arising in limbic portions of the temporal lobe also could cause behavioral changes (1). And indeed, they can. Yet the frontal lobes also contain large tracks of limbic cortex (i.e., anterior cingulate and orbitofrontal) as well as the prefrontal regions that make critical contributions to personality and to social and executive functions. Seizure foci arising in frontal regions similarly would be expected to alter behavioral function and result in mental pathology in some patients. And indeed, they do.

Swinkels and colleagues' study adds to a growing body of data that neither lateralization nor localization predicts risk of depression or anxiety in patients with partial epilepsy (2,3). A prospective, multicenter, epilepsy surgery study of 360 patients found no differences in the rates of depression or anxiety based on the side or lobe of seizure focus at baseline or the side or lobe of surgery (2). Similar to the Swinkels et al. investigation, this study found that higher baseline seizure frequency predicted an increased risk of psychiatric morbidity.

Studies published in the past few decades provide mounting evidence that a variety of partial as well as generalized epilepsies are associated with higher rates of psychiatric disease and adverse psychosocial outcomes when compared to patients with various medical conditions or other types of neurological disorders (4,5,6). It appears that in some patients, almost all epilepsies can alter interictal behavior. The accumulating data seem to move further away from the original concept that TLE is a risk factor for psychopathology and behavioral change. Or do

they? Given the role of the frontal lobes in behavior, changes in mood, personality, as well as in social and executive functions would be expected in some patients with frontal lobe epilepsy.

Geschwind's main goal was to cull out specific features and clustering of behavioral changes in TLE. Thus, traits such as hypergraphia, increased emotionality and religious interests, or hyposexuality and viscosity, were considered relatively specific for TLE, and the concurrence of three or four of these traits in a patient suggested TLE. However, subsequent studies using the Bear-Fedio Inventory led to conflicting results, as some investigators found differences in particular features between TLE and other epilepsy syndromes, while many others did not (7). Yet, certain elements, such as hypergraphia, religiosity, and hyposexuality may have anatomic correlates, for example, to hippocampal atrophy (8,9).

More than 30 years after Waxman and Geschwind proposed an interictal behavioral syndrome in TLE, the jury is still out (10). Much of the controversy is perceptual. The purported syndrome was not defined by depression, anxiety, or other traditional psychiatric classifications. Indeed, that was the point! This interictal behavioral syndrome of TLE is not seen in patients with primary psychiatric disorders. Thus, the finding that the frequency of depression and anxiety disorders is similar among groups with TLE and other epilepsies is consistent with Waxman and Geschwind's original thesis. However, the surface has barely been scratched on the issue of social and behavioral changes in people with epilepsy. And until additional research brings more clarity, it remains uncertain if some localized or generalized epilepsy syndromes are associated with a characteristic spectrum or cluster of behavioral changes.

by Orrin Devinsky, MD

References

1. Geschwind N. Behavioural changes in temporal lobe epilepsy. *Psychol Med* 1979;9:217–219.
2. Devinsky O, Barr WB, Vickrey BG, Berg AT, Bazil CW, Pacia SV, Langfitt JT, Walczak TS, Sperling MR, Shinnar S, Spencer SS. Changes in depression and anxiety after resective surgery for epilepsy. *Neurology* 2005;65:1744–1749.
3. Helmstaedter C, Sonntag-Dillender M, Hoppe C, Elger CE. Depressed mood and memory impairment in temporal lobe epilepsy as a function of focus lateralization and localization. *Epilepsy Behav* 2005;5:696–701.
4. Cutting S, Lauchheimer A, Barr W, Devinsky O. Adult-onset idiopathic generalized epilepsy: clinical and behavioral features. *Epilepsia* 2001;42:1395–1398.
5. Weiser HG. Selective amygdalohippocampectomy: indications, investigative technique and results. *Adv Tech Stand Neurosurg* 1986;13:39–133.
6. Wirrell EC, Camfield CS, Camfield PR, Dooley JM, Gordon KE, Smith B. Long-term psychosocial outcome in typical absence epilepsy. Sometimes a wolf in sheep's clothing. *Arch Pediatr Adolesc Med* 1997;151a:152–158.
7. Devinsky O, Najjar S. Evidence against the existence of a temporal lobe epilepsy personality syndrome. *Neurology* 1999;53(5 suppl 2):S13–25.
8. Wuerfel J, Krishnamoorthy ES, Brown RJ, Lemieux L, Koeppe M, Tebartz van Elst L, Trimble MR. Religiosity is associated with hippocampal but not amygdala volumes in patients with refractory epilepsy. *J Neurol Neurosurg Psychiatry* 2004;75:640–642.
9. van Elst LT, Krishnamoorthy ES, Baumer D, Selai C, von Gunten A, Gene-Cos N, Ebert D, Trimble MR. Psychopathological profile in patients with severe bilateral hippocampal atrophy and temporal lobe epilepsy: evidence in support of the Geschwind syndrome? *Epilepsy Behav* 2003;4:291–297.
10. Waxman SG, Geschwind N. The interictal behavior syndrome of temporal lobe epilepsy. *Arch Gen Psychiatry* 1975;32:1580–1586.

THE STATUS OF INTRAVENOUS VALPROATE FOR STATUS

Sodium Valproate vs Phenytoin in Status Epilepticus: A Pilot Study. Misra UK, Kalita J, Patel R. *Neurology* 2006;67(2):340–342. Sixty-eight patients with convulsive status epilepticus (SE) were randomly assigned to two groups to study the efficacy of sodium valproate (VPA) and phenytoin (PHT). Seizures were aborted in 66% in the VPA group and 42% in the PHT group. As a second choice in refractory patients, VPA was effective in 79% and PHT was effective in 25%. The side effects in the two groups did not differ. Sodium valproate may be preferred in convulsive SE because of its higher efficacy.

COMMENTARY

First line treatment for status epilepticus in the United States consists of intravenous (IV) benzodiazepines, usually lorazepam; the treatment protocol is based primarily on the results of the landmark Veterans Affairs Cooperative Status

Epilepticus trial results (1). In that trial, lorazepam was most effective (65% of patients), followed by phenobarbital (58%), diazepam plus phenytoin (56%), and then phenytoin alone (44%; statistically inferior to lorazepam). Another important observation from the trial was that when first line treatment was ineffective (which occurred in 45% of cases), the second agent was rarely effective—only 7% of patients responded to the second agent overall. This finding has led some experts to suggest moving directly to the highly effective and rapidly acting, anesthesia-inducing, continuous infusions (e.g., midazolam or propofol) when lorazepam fails. These continuous infusions can be initiated instead of or simultaneous with loading of the next agent, which is most commonly phenytoin or fosphenytoin (though phenobarbital still remains a reasonable option).

Misra et al. now have performed a useful prospective, randomized, head-to-head clinical trial of IV valproate versus IV phenytoin as first line treatments for convulsive SE in 68 patients of any age (the majority were adults). The study was performed in India and was not blinded. Status epilepticus was appropriately defined as >10 min of seizure activity (continuous or intermittent without return to baseline). Why benzodiazepines were not given first is unclear, but the study design makes the results easier to interpret, even if a bit less applicable to clinical care in the United States. As the initial treatment, valproate (30 mg/kg over 15 min) was superior to phenytoin (18 mg/kg at 50 mg/min), aborting status epilepticus in two-thirds of patients versus 42%, a difference barely reaching statistical significance. If the first medication failed, patients were crossed over to the other. As a second line agent, valproate was markedly superior to phenytoin ($p = 0.004$). About half the patients had recurrence of seizures within 24 h, with seizure freedom nonsignificantly more likely in those who received valproate alone (57%) than phenytoin alone (43%). Adverse effects were similar, but phenytoin was associated with hypotension in two patients (compared to none with valproate) and respiratory depression in two individuals (compared to one on valproate).

Appropriately, an EEG was performed if the patient did not wake up within 1 h after status epilepticus, as these patients are at high risk for nonconvulsive seizures. Information regarding the results of these EEGs or even how many were performed is not provided. However, prior studies have addressed this issue. For example, DeLorenzo et al. studied EEG after convulsive status epilepticus in 164 patients (2). After seemingly successful treatment of convulsive movements, 14% of the patients were in nonconvulsive status epilepticus on EEG and 48% had nonconvulsive seizures on prolonged monitoring—the vast majority with no overt clinical signs. Intermittent and continuous seizures were independently associated with worse outcome, with evidence of a “dose effect”: mortality was 13% if no seizures were found, 32% with intermittent seizures, and 51% when continuous seizures were seen.

The study by Misra et al. is not the first to show the utility of IV valproate for status epilepticus, but it is the first to demonstrate superiority over another commonly used IV antiepileptic drug. A prior retrospective study of IV valproate use in 102 patients in seizure emergencies, many of whom were in status epilepticus, showed success in 86% of patients (defined as cessation of seizures within 15 min and no return over 12 h) (3). Success in the subgroup of 35 patients with status epilepticus was 77%; 29 of those 35 patients had received benzodiazepines prior to valproate. There were no serious side effects reported. Although it is only approved up to a rate of 3 mg/kg/min, IV valproate has been shown to be safe and well tolerated at rapid infusion rates of up to 6 mg/kg/min (allowing a full load in approximately 5 min) (4). It has shown efficacy in multiple types of status epilepticus, in medically unstable elderly patients (5), and in all age groups, including children (6).

As mentioned, in the Veterans Affairs Cooperative Study, IV phenytoin by itself had the lowest success rate. However, it is unknown how valproate would have fared compared to the other study arms (including to the gold standard of lorazepam), because they were not part of the trial. Benzodiazepines should remain the first line treatment for status epilepticus—either IV, nasally, buccally, or rectally. However, when IV benzodiazepines fail, there are many reasonable options: phenytoin, fosphenytoin, midazolam, propofol, phenobarbital, and now IV valproate. It remains to be seen if IV valproate is effective when IV lorazepam has failed, though it appears to be based on the series from Peters and Pohlmann-Eden (3). If a patient is still convulsing after 0.1 mg/kg of lorazepam, proceeding to midazolam or propofol and simultaneously loading a longer-term antiepileptic drug, such as IV phenytoin/fosphenytoin or IV valproate, is a reasonable approach as well. However, a randomized trial of potential treatment options after lorazepam failure is needed. It would have been interesting if Misra and colleagues had included a third arm that evaluated lorazepam as well, though doing so would have required a larger, more costly trial.

Valproate is not without its downside. It can cause coagulopathy as a result of thrombocytopenia (usually dose-related and benign), platelet dysfunction, and hypofibrinogenemia as well as from deficiencies of multiple coagulation factors (e.g., von Willebrand factor, vitamin K-dependent coagulation factors, and factor XIII) (7). It also is associated with hyperammonemic encephalopathy, parkinsonism, and rarely pancreatitis or hepatic failure. Nonetheless, it is quite well tolerated as a rapid IV infusion and is effective for all types of status epilepticus. Now that there is evidence that valproate may be more effective than IV phenytoin for convulsive status epilepticus, it should be confirmed in a larger, double-blinded study, including an evaluation of its efficacy following benzodiazepines.

The study by Misra et al. and other recent studies clearly establish IV valproate as one of the prime players in the treatment of status epilepticus. Time will tell if recently released IV levetiracetam will be similarly effective and well tolerated and where exactly all these agents fit into current treatment algorithms. It certainly is good news to have more IV agents available for status epilepticus and more prospective randomized trials such as this one.

by Lawrence J. Hirsch, MD

References

1. Treiman DM, Meyers PD, Walton NY, Collins JF, Colling C, Rowan AJ, Handforth A, Faught E, Calabrese VP, Uthman BM, Ramsay RE, Mamdani MB. A comparison of four treatments for generalized convulsive status epilepticus. Veterans Affairs Status Epilepticus Cooperative Study Group. *N Engl J Med* 1998;339:792–798.
2. DeLorenzo RJ, Waterhouse EJ, Towne AR, Boggs JG, Ko D, DeLorenzo GA, Brown A, Garnett L. Persistent nonconvulsive status epilepticus after the control of convulsive status epilepticus. *Epilepsia* 1998;39:833–840.
3. Peters CN, Pohlmann-Eden B. Intravenous valproate as an innovative therapy in seizure emergency situations including status epilepticus—experience in 102 adult patients. *Seizure* 2005;14:164–169.
4. Limdi NA, Shimpi AV, Faught E, Gomez CR, Burneo JG. Efficacy of rapid IV administration of valproic acid for status epilepticus. *Neurology* 2005;64:353–355.
5. Sinha S, Naritoku DK. Intravenous valproate is well tolerated in unstable patients with status epilepticus. *Neurology* 2000;55:722–724.
6. Yu KT, Mills S, Thompson N, Cunanan C. Safety and efficacy of intravenous valproate in pediatric status epilepticus and acute repetitive seizures. *Epilepsia* 2003;44:724–726.
7. Gerstner T, Teich M, Bell N, Longin E, Dempfle CE, Brand J, König S. Valproate-associated coagulopathies are frequent and variable in children. *Epilepsia* 2006;47:1136–1143.

EPILEPSY SURGERY IN THE FRONTAL LOBE: *TERRA INCOGNITA* OR NEW FRONTIER?

Surgical Outcome and Prognostic Factors of Frontal Lobe Epilepsy Surgery. Jeha LE, Najm I, Bingaman W, Dinner D, Widdess-Walsh P, Luders H. *Brain* 2007;130(Pt 2):574–584. Epub 2007 Jan 5. Frontal lobe epilepsy (FLE) surgery is the second most common surgery performed to treat pharmacoresistant epilepsy. Yet, little is known about long-term seizure outcome following frontal lobectomy. The aim of this study is to investigate the trends in longitudinal outcome and identify potential prognostic indicators in a cohort of FLE patients investigated using modern diagnostic techniques. We reviewed 70 patients who underwent a frontal lobectomy between 1995 and 2003 (mean follow-up 4.1 ± 3 years). Data were analysed using survival analysis and multivariate regression with Cox proportional hazard models. A favourable outcome was defined as complete seizure-freedom, allowing for auras and seizures restricted to the first post-operative week. The estimated probability of complete seizure-freedom was 55.7% [95% confidence interval (CI) = 50–62] at 1 post-operative year, 45.1% (95% CI = 39–51) at 3 years, and 30.1% (95% CI = 21–39) at 5 years. Eighty per cent of seizure recurrences occurred within the first 6 post-operative months. Late remissions and relapses occurred, but were rare. After multivariate analysis, the following variables retained their significance as independent predictors of seizure recurrence: MRI-negative malformation of cortical development as disease aetiology [risk ratio (RR) = 2.22, 95% CI = 1.40–3.47], any extrafrontal MRI abnormality (RR = 1.75, 95% CI = 1.12–2.69), generalized/non-localized ictal EEG patterns (RR = 1.83, 95% CI = 1.15–2.87), occurrence of acute post-operative seizures (RR = 2.17, 95% CI = 1.50–3.14) and incomplete surgical resection (RR = 2.56, 95% CI = 1.66–4.05) (log likelihood-ratio test P -value < 0.0001). More than half of patients in favourable prognostic categories were seizure-free at 3 years, and up to 40% were seizure-free at 5 years, compared to <15% in those with unfavourable outcome predictors. These data underscore the importance of appropriate selection of potential surgical candidates.

COMMENTARY

The temporal lobectomy has been the most common neurosurgical treatment for medically intractable focal epilepsy. Its effectiveness has been well demonstrated, most

recently by a randomized controlled trial (1). Nonetheless, a substantial portion of intractable epilepsy originates outside the mesial temporal region, and as epilepsy surgery has become more widely available, large academic epilepsy centers are evaluating greater numbers of extratemporal epilepsy patients for neurosurgical treatment. The overall chance of seizure freedom from resections in extratemporal regions is acknowledged to be lower than that of temporal resections, however, past literature shows quite variable outcomes and prognostic indicators for

frontal lobe epilepsy surgery (2–4). When should neurosurgical treatment be considered for frontal lobe epilepsy?

In this regard, the report by Jeha et al. is welcome because it presents a carefully analyzed series of 70 frontal epilepsy resections over a 9-year period, excluding only patients with prior neurosurgery, resections extending outside the frontal lobe, or with less than 1 year of follow-up. This series represents outcomes of patients for whom the best diagnostic and surgical techniques available were used, representing a “snapshot” of the best therapeutic approaches of the recent past. It demonstrates a rate of seizure-freedom that justifies neurosurgical treatment for many subgroups of frontal epilepsy patients.

The most significant finding of the study concerns the strong effect of particular factors on patient outcome. The preoperative clinical characteristics of patients, such as gender, age at the time of seizure onset or operation, family history, and seizure types and frequency, did not significantly affect prognosis. Multivariate analysis demonstrated that a seizure-free outcome depended on accurately locating the seizure focus, defining its full extent, and removing all of it, since the significant predictive factors were MRI findings, interictal and ictal EEG findings, and completeness of resection. After surgery, occurrence of auras or immediate postoperative seizures was a negative predictor. In the subset of patients with favorable prognostic factors on multivariate analysis, seizure-free outcome approached that seen for temporal lobectomy, with a 50% to 60% seizure-free rate at 3 years, while the rate was only 0% to 14% for those with unfavorable factors. There were 12 patients who had all favorable prognostic factors, with localized and completely resected MRI lesions, localizing ictal EEG, and no acute postoperative seizures; 11 of these 12 were completely seizure free at the last follow-up.

A strength of this report is the longitudinal analysis, since surgical outcome is very much a function of duration of follow-up. The analysis reveals a significant decrease in complete seizure freedom over the 5-year period following surgery—much like what has been described after temporal lobectomy (5–7). However, recurrence tended to happen earlier in this frontal lobe series, with 80% occurring in the first 6 months. Of concern, recurrence also was more persistent, with 60% of the failing patients not seizure-free at the last follow-up. However, seizure-freedom at 3 and 5 years was significantly higher in patients with favorable prognostic factors.

Univariate analysis of the effect of etiology on patient outcome was informative. Twenty-one patients had a well-defined etiology of neoplasm, vascular malformation, or encephalomalacia, with 12 individuals achieving seizure freedom. Seven were cryptogenic, with normal MRI and pathology, and only two of

these seven became seizure free. The largest group was made up of 41 patients with cortical developmental malformations. It is this group that poses the greatest challenge for presurgical diagnostic evaluation. Fifteen of 29 patients with MRI-positive malformations were completely seizure-free. In contrast, 11 of 12 with MRI-negative, but pathologically demonstrated cortical developmental malformations had seizure recurrence. This finding implies that correct localization of a seizure focus, as documented by neuropathology, is not sufficient to produce a good outcome if the full extent of the pathology cannot be outlined for complete surgical removal. When fully characterized by neuroimaging, cortical dysplasia, and other developmental malformations are amenable to successful surgical treatment.

This study confirms that treatment of medically intractable cases with well-delineated and totally removable seizure foci in the frontal lobe is familiar territory and that these patients most definitely are surgical candidates. In contrast, the evaluation of those with poorly defined dysplasia or normal MRI represents *terra nova*—a new frontier, in which diagnostic innovations are needed to improve outcomes and extend the benefits of frontal lobe epilepsy surgery to more individuals.

by John W. Miller, MD, PhD

References

1. Wiebe S, Blume WT, Girvin JP, Eliasziw M. A randomized, controlled trial of surgery for temporal lobe epilepsy. *N Engl J Med* 2001;345:311–318.
2. Fish DR, Smith SJ, Quesney LF, Andermann F, Rasmussen T. Surgical treatment of children with medically intractable frontal or temporal lobe epilepsy: results, highlights of 40 years' experience. *Epilepsia* 1993;34:244–247.
3. Mosewich RK, So EL, O'Brien TJ, Cascino GD, Sharbrough FW, Marsh WR, Meyer FB, Jack CR, O'Brien PC. Factors predictive of the outcome of frontal lobe epilepsy surgery. *Epilepsia* 2000;41:843–849.
4. Zaatreh MM, Spencer DD, Thompson JL, Blumenfeld H, Novotny EJ, Mattson RH, Spencer SS. Frontal lobe tumoral epilepsy: clinical, neurophysiologic features and predictors of surgical outcome. *Epilepsia* 2002;43:727–733.
5. Jeha LE, Najm IM, Bingaman WE, Khandwala F, Widdess-Walsh P, Morris HH. Predictors of outcome after temporal lobectomy for the treatment of intractable epilepsy. *Neurology* 2006;66:1938–1940.
6. McIntosh AM, Kalnins RM, Mitchell LA, Fabinyi GC, Briellmann RS, Berkovic SF. Temporal lobectomy: long-term seizure outcome, late recurrence and risks for seizure recurrence. *Brain* 2004;127:2018–2030.
7. Yoon HH, Kwon HL, Mattson RH, Spencer DD, Spencer SS. Long-term seizure outcome in patients initially seizure-free after resective epilepsy surgery. *Neurology* 2003;61:445–450.

THE GENETICS OF TEMPORAL LOBE EPILEPSY AND IMPLICATIONS FOR TREATMENT

Temporal Lobe Epilepsy and GEFS+ Phenotypes Associated with SCN1B Mutations. Scheffer IE, Harkin LA, Grinton BE, Dibbens LM, Turner SJ, Zielinski MA, Xu R, Jackson G, Adams J, Connellan M, Petrou S, Wellard RM, Briellmann RS, Wallace RH, Mulley JC, Berkovic SF. *Brain* 2007;130(Pt 1):100–109. *SCN1B*, the gene encoding the sodium channel $\beta 1$ subunit, was the first gene identified for generalized epilepsy with febrile seizures plus (GEFS+). Only three families have been published with *SCN1B* mutations. Here, we present four new families with *SCN1B* mutations and characterize the associated phenotypes. Analysis of *SCN1B* was performed on 402 individuals with various epilepsy syndromes. Four probands with missense mutations were identified. Detailed electroclinical phenotyping was performed on all available affected family members including quantitative MR imaging in those with temporal lobe epilepsy (TLE). Two new families with the original C121W *SCN1B* mutation were identified; novel mutations R85C and R85H were each found in one family. The following phenotypes occurred in the six families with *SCN1B* missense mutations: 22 febrile seizures, 20 febrile seizures plus, five TLE, three other GEFS+ phenotypes, two unclassified and ten unaffected individuals. All individuals with confirmed TLE had the C121W mutation; two underwent temporal lobectomy (one with hippocampal sclerosis and one without) and both are seizure free. We confirm the role of *SCN1B* in GEFS+ and show that the GEFS+ spectrum may include TLE alone. TLE with an *SCN1B* mutation is not a contraindication to epilepsy surgery.

COMMENTARY

Temporal lobe epilepsy is the most common partial epilepsy. It was once considered to be predominantly an acquired form of epilepsy, and no temporal lobe epilepsy syndrome was listed among the presumably genetic idiopathic epilepsies, even though a genetic element was often recognized (1). However, in the past decade a number of genetic temporal lobe epilepsy syndromes have been described (2–4). First reported was a benign familial temporal lobe epilepsy syndrome identified in twins, with clinical evidence of mesial temporal lobe origin (2). The syndrome was not associated with hippocampal sclerosis. Pedigree analysis was consistent with an autosomal dominant inheritance with incomplete penetrance. Extrapolation from its incidence in monozygotic twins suggested that it was a common epileptic syndrome, perhaps not previously recognized because of its excellent response to medical treatment and its frequently very mild manifestations. It soon became clear that this was not the only genetic temporal lobe epilepsy syndrome and that familial temporal lobe epilepsy is clinically heterogeneous (3,5). Some patients have a history of antecedent febrile seizures or hippocampal sclerosis while others do not; some patients have easily controlled seizures while others have refractory epilepsy, leading to presurgical evaluation and epilepsy surgery. While most familial temporal lobe epilepsy is of mesial temporal origin, familial lateral temporal lobe syndromes with an identified genetic mutation have been recognized. Autosomal dominant partial epilepsy, with auditory

(4) or aphasic (6) manifestations, is related to mutations in the leucine-rich, glioma-inactivated 1 (LGI1) gene.

Mesial temporal lobe epilepsy is much more common than lateral temporal lobe epilepsy and is frequently associated with antecedent childhood complex febrile convulsions, which are considered to have a strong genetic component (7,8). Some evidence suggests that in families with febrile seizures temporal lobe epilepsy may be in part a consequence of febrile status epilepticus (9). One genetic epilepsy syndrome, which includes febrile seizures, is generalized epilepsy with febrile seizures plus (GEFS+) (10,11). GEFS+ most often is due to a mutation in a sodium channel gene and manifests with variable phenotypes, including the typical febrile convulsion syndrome, febrile seizures persisting beyond age 6 years, or coexistent with afebrile generalized tonic–clonic seizures (called febrile seizures plus), as well as other generalized seizure types (8,9). Temporal lobe epilepsy is an occasional phenotype with GEFS+, but published cases thus far had suggested that temporal lobe epilepsy with this condition may be a consequence of febrile status epilepticus, rather than a direct expression of GEFS+ (11,12). In their report, Scheffer and colleagues screened a large number of individuals for mutations in one of the sodium channel genes implicated in GEFS+. In the affected families, they found several individuals with temporal lobe epilepsy that occurred only with one particular mutation and was not necessarily preceded by febrile seizures or associated with hippocampal sclerosis. Scheffer and colleagues concluded that: “GEFS+ phenotypes may include temporal lobe epilepsy in its own right.”

One important impetus for the identification of genetic causes of epilepsy is the expectation that more specific and effective therapy may follow. However, there has not been a clear impact of genetic discoveries on the treatment of temporal lobe

epilepsy. One specific question in the management of epilepsy is whether a genetic basis of epilepsy precludes epilepsy surgery. Intuitively, focal resection does not seem appropriate in genetic epilepsy because the genetic mutation is expected to affect the brain diffusely. However, the data do not support this intuitive conclusion. In a report of epilepsy surgery in 20 patients with familial temporal lobe epilepsy of unspecified genetic basis, surgical outcome was excellent in 85% (13). The three patients in the study with less than excellent outcomes had either absent or symmetrical hippocampal atrophy, while all others had unilateral or asymmetric hippocampal atrophy. Scheffer and colleagues report excellent surgical outcome in two individuals with GEFS+ and temporal lobe epilepsy, including one without hippocampal sclerosis. They confirm that a GEFS+ mutation is specifically not a contraindication to epilepsy surgery when the epileptogenic zone is well localized.

by Bassel W. Abou-Khalil, MD

References

1. Commission on Classification and Terminology of the International League Against Epilepsy. Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsia* 1989;30:389–399.
2. Berkovic SF, McIntosh A, Howell RA, Mitchell A, Sheffield LJ, Hopper JL. Familial temporal lobe epilepsy: a common disorder identified in twins. *Ann Neurol* 1996;40:227–235.
3. Cendes F, Lopes-Cendes I, Andermann E, Andermann F. Familial temporal lobe epilepsy: a clinically heterogeneous syndrome. *Neurology* 1998;50:554–557.
4. Winawer MR, Ottman R, Hauser WA, Pedley TA. Autosomal dominant partial epilepsy with auditory features: defining the phenotype. *Neurology* 2000;54:2173–2176.
5. Kobayashi E, Lopes-Cendes I, Guerreiro CA, Sousa SC, Guerreiro MM, Cendes F. Seizure outcome and hippocampal atrophy in familial mesial temporal lobe epilepsy. *Neurology* 2001;56:166–172.
6. Gu W, Brodtkorb E, Steinlein OK. LGI1 is mutated in familial temporal lobe epilepsy characterized by aphasic seizures. *Ann Neurol* 2002;52:364–367.
7. French JA, Williamson PD, Thadani VM, Darcey TM, Mattson RH, Spencer SS, Spencer DD. Characteristics of medial temporal lobe epilepsy: I. Results of history and physical examination. *Ann Neurol* 1993;34:774–780.
8. Hamati-Haddad A, Abou-Khalil B. Epilepsy diagnosis and localization in patients with antecedent childhood febrile convulsions. *Neurology* 1998;50:917–922.
9. Maher J, McLachlan RS. Febrile convulsions. Is seizure duration the most important predictor of temporal lobe epilepsy? *Brain* 1995;118(Pt 6):1521–1528.
10. Scheffer IE, Berkovic SF. Generalized epilepsy with febrile seizures plus. A genetic disorder with heterogeneous clinical phenotypes. *Brain* 1997;120(Pt 3):479–490.
11. Singh R, Scheffer IE, Crossland K, Berkovic SF. Generalized epilepsy with febrile seizures plus: a common childhood-onset genetic epilepsy syndrome. *Ann Neurol* 1999;45:75–81.
12. Abou-Khalil B, Ge Q, Desai R, Ryther R, Bazyk A, Bailey R, Haines JL, Sutcliffe JS, George AL Jr. Partial and generalized epilepsy with febrile seizures plus and a novel SCN1A mutation. *Neurology* 2001;57:2265–2272.
13. Kobayashi E, D'Agostino MD, Lopes-Cendes I, Andermann E, Dubeau F, Guerreiro CA, Schenka AA, Queiroz LS, Olivier A, Cendes F, Andermann F. Outcome of surgical treatment in familial mesial temporal lobe epilepsy. *Epilepsia* 2003;44:1080–1084.

HELPFUL DATA, BUT LESS CERTAINTY

Childhood Mesial Temporal Sclerosis. Ng YT, McGregor AL, Duane DC, Jahnke HK, Bird CR, Wheless JW. *J Child Neurol* 2006;21(6):512–517. The prevalence and clinical characteristics of mesial temporal sclerosis have not been well studied in children. All brain magnetic resonance imaging (MRI) reports of children less than 14 years of age were reviewed from two tertiary institutions. A 52-month period from one institution and a 37-month period from the other were reviewed. All reports of definite or possible mesial temporal sclerosis were noted. These patients' MRIs were then reviewed to confirm the MRI diagnostic criteria of mesial temporal sclerosis. The charts of the patients who satisfied these criteria were reviewed in detail. Three thousand one hundred brain MRI reports were reviewed. Twenty-six reports of mesial temporal sclerosis were found. Twenty-four of the 26 films satisfied the criteria of mesial temporal sclerosis by MRI after the films were reviewed. The prevalence among all pediatric brain MRI studies was 0.77%. All patients had presented with seizures, that is, there were no "incidental" findings of mesial temporal sclerosis. Four patients had a history of febrile seizures. Mesial temporal sclerosis is an uncommon finding in children, but when it occurs, it is always associated with epilepsy. Asymptomatic mesial temporal sclerosis or mesial temporal sclerosis not associated with a seizure disorder did not occur in our series. Febrile seizures can occur in association with mesial temporal sclerosis presenting in childhood.

COMMENTARY

This careful, retrospective study by Ng and colleagues of a patient MRI database serves as a reminder that mesial temporal sclerosis (MTS) may already be present in childhood. The mean age of children whose MRI disclosed abnormalities consistent with MTS was 6.9 years (range: 0.7–12.7 years). Two factors likely contributed to detection of MTS in 24 (3.9%) of the 623 pediatric epilepsy patients: (a) use of a specific epilepsy protocol with fine-cut coronal T₂-weighted sequences perpendicular to the hippocampi (supplemented by coronal fluid-attenuated inversion recovery (FLAIR) images from one institution), and (b) the severity of seizure disorders of these 24 patients. Among 22 patients for whom therapeutic data were available, 11 (50%) were daily taking more than one antiepileptic drug. Moreover, 13 of the 24 patients had a potentially epileptogenic MRI lesion in addition to MTS, a significant neurological antecedent (e.g., encephalitis, trauma), or both. Curiously, febrile seizures occurred in only 4 of the 24 children, 2 of whom had potentially epileptogenic factors. These data support the concept that MTS may result from one or more of several pathophysiological factors (1).

Although MTS appeared only among epilepsy patients and not among the 2,477 children with other conditions, the implied high specificity is negated by their employment of the MTS-revealing epilepsy protocol only for seizure disorder patients. Subsequent application of the epilepsy protocol to a set of sequential patients referred for MRI for any reason may more confidently establish specificity. Results may relate to findings by Benbadis et al., which identified both hippocampal atrophy and T2 signal abnormalities in 8 (4%) of 204 patients without a seizure history who underwent MRI (2). Thus, the Ng et al. study (pediatric epilepsy patients) and the Benbadis study (nonepilepsy patients) disclosed abnormalities compatible with MTS in 3.9% and 4%, respectively!

Some epileptologists have proposed that two conditions must coexist for MTS to develop: (a) an initial precipitating injury (e.g., prolonged febrile seizure, CNS infection, head trauma) and (b) any factor that increases vulnerability to neuronal injury. Accumulating evidence supports various candidates for this second mechanism—among them, hippocampal dysgenesis and a genetic component (3,4). As only 4 of the

24 MTS patients in the Ng et al. study had febrile seizures, other tandems (such as trauma and hippocampal dysgenesis) may suffice.

Despite the presence of MTS, clinical descriptions and video-telemetry disclosed complex partial seizures in only 16 of 22 patients for whom data were available in the Ng et al. study. It is known that temporal lobe seizures in young children (< 6 years) may manifest as symmetrical tonic or clonic events of the limbs (5). In fact, EEG-documented seizures originating in a temporal lobe that contained MTS were identified in only 8 of 18 patients with an identified epileptogenic focus. This finding may reflect the dual pathology of MTS and a separate epileptogenic area. (6).

Advances in neuroimaging disclose abnormalities, such as MTS, that may have escaped earlier techniques. Data of this study underline the need for clinicians to realize that lesion-based temporal lobe epilepsy may begin in childhood. However, establishment of specificity, that is, clinical significance of any abnormality revealed in a laboratory test, remains a challenge.

by Warren T. Blume, MD

References

1. Najm IM, Babb TL, Mohamed A, Diehl B, Ng TC, Bingaman WE, Luders HO. Mesial temporal sclerosis. In: Luders HO, Comair YG, eds. *Epilepsy Surgery*, 2nd edition. Philadelphia: Lippincott Williams and Wilkins, 2001:95–103.
2. Benbadis SR, Wallace J, Reed Murtagh F. MRI evidence of mesial temporal sclerosis in subjects without seizures. *Seizure* 2002;11:340–343.
3. Lewis DV. Losing neurons: selective vulnerability and mesial temporal sclerosis. *Epilepsia* 2005;46(suppl. 7):39–44.
4. Kobayashi E, D'Agostino MD, Lopes-Cendes I, Berkovic SF, Li ML, Andermann E, Andermann F, Cendes F. Hippocampal atrophy and T2-weighted signal changes in familial mesial temporal epilepsy. *Neurology* 2003; 60: 405–409.
5. Brockhaus A, Elger CE. Complex partial seizures of temporal lobe origin in children of different age groups. *Epilepsia* 1995; 36: 1173–1181.
6. Lawn N, Londono A, Sawrie S, Morawetz R, Martin R, Gilliam F, Faught E, Kuzniecky R. Occipitoparietal epilepsy, hippocampal atrophy, and congenital developmental abnormalities. *Epilepsia* 2000;41:1546–1553.