Published in final edited form as:

Horm Behav. 2007 June ; 52(1): 45–55.

SEXUAL BEHAVIOR IN MALE RODENTS

Elaine M. Hull and Juan M. Dominguez

Department of Psychology and Program in Neuroscience, Florida State University, Tallahassee, FL 32306-1270

Abstract

The hormonal factors and neural circuitry that control copulation are similar across rodent species, although there are differences in specific behavior patterns. Both estradiol (E) and dihydrotestosterone (DHT) contribute to the activation of mating, although E is more important for copulation and DHT, for genital reflexes. Hormonal activation of the medial preoptic area (MPOA) is most effective, although implants in the medial amygdala (MeA) can also stimulate mounting in castrates. Chemosensory inputs from the main and accessory olfactory systems are the most important stimuli for mating in rodents, especially in hamsters, although genitosensory input also contributes. Dopamine agonists facilitate sexual behavior, and serotonin (5-HT) is generally inhibitory, though certain 5-HT receptor subtypes facilitate erection or ejaculation. Norepinephrine agonists and opiates have dose-dependent effects, with low doses facilitating and high doses inhibiting behavior.

Keywords

Rats; mice; hamsters; guinea pigs; estradiol; dihydrotestosterone; testosterone; medial preoptic area; medial amygdala; genital reflexes

Introduction

Reproductive behaviors and their neural and hormonal regulation vary widely across species. Yet much research has focused on relatively few animals. We describe the behaviors of male rodents and their neural, hormonal, and experiential regulation. We begin with rats, the most common subjects of laboratory research. We then describe the behaviors of male mice, hamsters, and guinea pigs, noting similarities and differences among species. Sexual behavior is highly interactive; here we concentrate on the male, keeping in mind that the contributions of the female are equally important. Because of the vast amount of research on rodents, and the page limits for this manuscript, we can cite only a small portion of it. For additional details, please consult Hull et al. (2006) or Hull et al. (2002).

Description of male rat copulatory behaviors and *ex copula* reflexes

Male rats usually begin a sexual encounter by investigating the female's face and anogenital region. Both partners may emit mutually arousing 50 kHz ultrasonic vocalizations. The male approaches from the female's rear, mounts, and gives several rapid shallow thrusts (19–23 Hz) with his pelvis; if he detects the female's vagina, he gives a deeper thrust, inserting his penis into her vagina for 200–300 msec (Beyer et al., 1981). He then springs backward rapidly and grooms his genitals. After 7 to 10 intromissions, 1 to 2 minutes apart, he will ejaculate.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Ejaculation is characterized by a longer, deeper thrust (750–2000 msec) and much slower dismount (Beyer et al., 1981). It is accompanied by rhythmic contractions of the bulbospongiosus and ischiocavernosus muscles at the base of the penis, and of anal sphincter and skeletal muscles (Holmes et al., 1991). After ejaculation, he grooms himself and then rests during the postejaculatory interval (PEI), which may last for 6 to 10 minutes before resuming mating. During the first 50 – 75% of the PEI, the male will not copulate again and emits 22 kHz ultrasonic vocalizations. During the latter 25%, he may resume copulation if presented with a novel female or a mildly painful stimulus. After 7–8 ejaculations males reach satiety and usually will not copulate again for 1 to 3 days. Previous sexual experience confers greater copulatory “efficiency” and increased resistance to the effects of various lesions, castration, and stress (reviewed in Hull et al., 2006).

Copulatory ability is acquired between 45 and 75 days of age (reviewed in Meisel and Sachs, 1994). Prepubertal castration prevented the onset of mating behavior, and exogenous testosterone (T) or estradiol (E₂) hastened its development. Aging male rats lose the ability to ejaculate, which is not restored by exogenous T (Chambers et al., 1991). A decline in estrogen receptors (ER) (Roselli et al., 1993), but not androgen receptors (AR) (Chambers et al., 1991), may underlie the deficit in old males.

Ex copula reflexes can be observed in several contexts. Spontaneous or drug-induced erections occur in the home cage or neutral arena. Volatile odors from an estrous female elicit noncontact erections, which may be a model for psychogenic erections in humans. In rats “touch-based” erections can be elicited by restraining the male on his back and retracting the penile sheath. These erections result from engorgement of the corpus spongiosum, which produces tumescence of the glans penis (reviewed in Hull et al., 2006; Meisel and Sachs, 1994). Anteroflexions also occur; these result from contractions of the ischiocavernosus muscle and erection of the corpus cavernosum, causing the penis to rise from its normal posteroflexed position. Occasionally, seminal emission occurs in this context. The continuing pressure of the retracted sheath around the base of the penis provides the stimulus for these touch-based reflexes. Finally, the urethrogenital reflex has been studied in anesthetized male and female rats as a model of orgasm in humans (McKenna et al., 1991). It is elicited by urethral distension, followed by release; it consists of clonic contractions of the perineal muscles.

Hormonal factors in the activation of male rat mating behavior

Male sexual behavior in virtually all vertebrate species is dependent on T, secreted by the Leydig cells of the testes and metabolized in target cells to either E₂ (by aromatization) or dihydrotestosterone (DHT, by 5 α -reduction). Plasma T is undetectable within 24 hours of castration (Krey and McGinnis 1990); however, copulatory ability decreases gradually over days or weeks. Five to 10 days of T are usually required to reinstate mating (McGinnis et al., 1989). However, E₂ increased chemo-investigation and mounting by castrates within 35 min (Cross and Roselli 1999). Therefore, rapid, probably membrane-based, hormonal effects may contribute to sexual motivation, but longer-term genomic effects are required for full restoration of mating.

The major hormone to activate sexual behavior in male rats is E₂, as proposed by the “aromatization hypothesis” (reviewed in Hull et al., 2006). DHT, which is nonaromatizable and has greater affinity for ARs than does T, is ineffective when administered alone. However, E₂ does not fully maintain male rat sexual behavior (McGinnis and Dreifuss, 1989; Putnam et al., 2003) or partner preference (Vagell and McGinnis, 1997). Thus, androgens contribute to motivation and performance and are also necessary and sufficient to maintain *ex copula* genital reflexes (Cooke et al., 2003; Manzo et al., 1999; Meisel et al., 1984). Although E₂ was ineffective in maintaining *ex copula* reflexes, it did maintain vaginal intromissions *in copula*.

(O'Hanlon, 1981). Sachs (1983) suggested that E activates a "behavioral cascade" that can elicit genital reflexes *in copula*, but cannot disinhibit them *ex copula*.

Effects of systemically administered drugs on male rat sexual behavior

Transmitters often act synergistically in multiple sites, and the site of action often is not known *a priori*. Therefore, systemic drug administration can be useful. Table 1 summarizes the effects on male rat sexual behavior of drugs and treatments that affect neurotransmitter function in more than one brain area.

Brain areas that regulate male rat sexual behavior

Chemosensory input from the main and vomeronasal systems is probably the most important stimulus for male rodent sexual behavior. Bilateral olfactory bulbectomy, which removes both the main and vomeronasal pathways, produced variable impairment of copulation and noncontact erections, with sexually naïve males being more susceptible to impairment (reviewed in Hull et al., 2006). Information from the main and accessory olfactory systems is processed in the medial amygdala (MeA), along with somatosensory input from the genitals, relayed through the parvocellular portion of the subparafascicular nucleus (SPFp), which is also part of an ejaculation circuit in several species (reviewed in Hull et al., 2006). Input from the MeA, both directly and via the bed nucleus of the stria terminalis (BNST), to the medial preoptic area (MPOA) is critical for copulation in male rats (Kondo and Arai, 1995).

The MPOA is arguably the most critical site for orchestrating male sexual behavior. It receives sensory input indirectly from all sensory systems and sends reciprocal connections back to those sources, thereby enabling the MPOA to influence the input that it receives (Simerly and Swanson, 1986). It also sends output to hypothalamic, midbrain, and brain stem nuclei that regulate autonomic and somatomotor patterns and motivational states (Simerly and Swanson, 1988). Many studies have reported severe and long-lasting impairment of copulation following lesions of the MPOA (reviewed in Hull et al., 2006). However, male rats with MPOA lesions continued to show noncontact erections (Liu et al., 1997) and bar-press for a light that had been paired with access to a female (Everitt, 1990). Everitt (1990) suggested that the MPOA is important only for copulation, and not sexual motivation. However, MPOA lesions impaired sexual motivation in other contexts, including preference for a female partner (Edwards and Einhorn, 1986; Paredes et al., 1998) and pursuit of a female (Paredes et al., 1993).

Conversely, stimulation of the MPOA facilitated copulation, but did not elicit mating in sated males (Rodriguez-Manzo et al., 2000). Stimulation also increased intracavernosal pressure in anesthetized males (Giuliano et al., 1996) and elicited the urethrogenital reflex without urethral stimulation (Marson and McKenna, 1994). The MPOA does not project directly to the lower spinal cord, where erection and seminal emission are controlled; thus, it must activate other areas that, in turn, elicit those reflexes.

The MPOA is the most effective site for hormonal stimulation of mating in castrated rats; however, T or E₂ implants in the MPOA did not fully restore copulation, and DHT implants were ineffective (reviewed in Hull et al., 2006). Therefore, both ER and AR in the MPOA contribute to copulatory ability of male rats; however, hormonal effects elsewhere are required for full activation of behavior.

MPOA microinjections of the classic dopamine (DA) agonist apomorphine facilitated copulation in gonadally intact and castrated rats and increased touch-based reflexes (reviewed in Dominguez & Hull, 2005; Hull et al., 2006). MPOA apomorphine also restored copulation in males with large amygdala lesions (Dominguez et al., 2001). Conversely, a DA antagonist inhibited copulation and touch-based reflexes and decreased sexual motivation without

affecting motoric function (reviewed in Dominguez and Hull, 2005; Hull et al., 2006). These effects were anatomically and behaviorally specific.

DA is released in the MPOA before and during copulation (Hull et al., 1995; Sato et al., 1995). Again, there was both behavioral and anatomical specificity. Recent, but not concurrent, T was necessary for the DA increase and copulation (Hull et al., 1995). A major factor promoting MPOA DA release is nitric oxide (NO), in both basal and female-stimulated conditions (reviewed in Dominguez and Hull, 2005; Hull et al., 2006). NO synthase immunoreactivity (NOS-ir) is positively regulated by both T and E₂ (Du and Hull, 1999; Putnam et al., 2005). NO is also important for copulatory performance, as a NOS inhibitor (L-NAME) in the MPOA blocked copulation in naïve males, impaired mating in experienced males, and prevented the facilitation produced in saline-treated males by 7 pre-exposures to an estrous female (Lagoda et al., 2004). Input from the MeA is required for the DA response to a female, but not for basal DA levels (Dominguez et al., 2001). Chemical stimulation of the MeA resulted in increases in extracellular DA in the MPOA comparable to those produced by a female (Dominguez and Hull, 2001). There are no DA-containing neurons in the amygdala of male rats; however, some efferents from the MeA to the MPOA, and even more from the BNST, appeared to be glutamatergic (Dominguez et al., 2003). Reverse-dialysis of glutamate into the MPOA increased DA release, an effect blocked by a NOS inhibitor (Dominguez et al., 2004). In addition, extracellular glutamate increased during copulation and rose to 300% of basal levels in the two-minute sample collected during ejaculation; reverse dialysis of glutamate reuptake inhibitors facilitated several measures of copulation (Dominguez et al., 2006). Similarly, glutamate microinjected into the MPOA increased intracavernous pressure (Giuliano et al., 1996) and the urethrogenital reflex (Marson and McKenna, 1994) in anesthetized rats. Therefore, a consistent picture emerges, in which glutamate, at least in part from the MeA and BNST, facilitates copulation and genital reflexes, both directly and via NO-mediated increases in DA, which also contributes to the initiation and progress of copulation. Other neurotransmitters in the MPOA that may facilitate male rat sexual behavior are norepinephrine, acetylcholine, prostaglandin E₂, and hypocretin/orexin (hcrt/orx), whereas GABA and 5-HT may be inhibitory. Low levels of opioids may facilitate, and higher doses inhibit copulation (reviewed in Hull et al., 2006).

Electrophysiological recordings revealed that different MPOA neurons contribute to sexual motivation and copulatory performance (Shimura et al., 1994). Mating increases Fos-ir in the MPOA (reviewed in Hull et al., 2006), with greater increases in sexually experienced males, compared to naïve ones, even though the experienced males had fewer intromissions preceding ejaculation (Lumley and Hull, 1999). Therefore, sexual experience may enhance the processing of sexually relevant stimuli.

The mesocorticolimbic DA tract, ascending from the ventral tegmental area (VTA) to the nucleus accumbens (NAc) and prefrontal cortex, is important for reinforcement and appetitive behaviors. It receives input from the MPOA (Simerly and Swanson, 1988) and numerous other sources. VTA or NAc lesions increased PEIs and decreased noncontact erections, but did not affect copulation (reviewed in Hull et al., 2006). Conversely, electrical stimulation of the VTA facilitated copulation (Markowski and Hull, 1995). Applications of drugs to the VTA or NAc primarily affected general activation, rather than specifically sexual behavior (reviewed in Hull et al., 2006). Mating activated Fos-ir in the NAc and VTA, and an estrous female-stimulated increase was enhanced by prior sexual experience (Lopez and Ettenberg, 2002a). Copulation and/or exposure to the odor of an estrous female increased DA release in the NAc (reviewed in Hull et al., 2006). Reverse dialysis of 5-HT into the anterior lateral hypothalamic area (LHA) decreased basal DA in the NAc and prevented the rise that otherwise occurred with the introduction of a female (Lorrain et al., 1999). Because 5-HT is increased in the LHA at

the time of ejaculation (Lorrain et al., 1997), the resulting decrease in NAc DA may contribute to the PEI.

The paraventricular nucleus (PVN) of the hypothalamus comprises a magnocellular division, which releases oxytocin and vasopressin into the circulation from the posterior pituitary, and a parvocellular division, which projects to several brain areas and the spinal cord. Excitotoxic lesions of the parvocellular portion decreased noncontact erections but did not impair copulation (Liu et al., 1997). Similar lesions decreased the amount of semen ejaculated and the number of oxytocin-containing fibers in the spinal cord, but again did not affect copulation (Ackerman et al., 1997). Lesions that encompassed both divisions did impair copulation, as well as touch-based and noncontact erections (Liu et al., 1997). Argiolas and Melis have provided an elegant picture in which DA, oxytocin, and glutamate (Melis et al., 2004) increase production of NO in oxytocinergic cells in the PVN, which then release oxytocin in the hippocampus (Melis et al., 1992), spinal cord (Ackerman et al., 1997), and elsewhere, thereby increasing erection and seminal emission and possibly enhancing copulation (reviewed in Argiolas and Melis, 2004). GABA and opioids inhibit these processes. This lab has also shown that DA (Melis et al., 2003), glutamate, (Melis et al., 2004), and NO (Melis et al., 1998) are released in the PVN during copulation.

Several additional brain areas influence male rat sexual behavior. 5-HT is released in the LHA at the time of ejaculation, as noted above, and microinjection of an SSRI into the LHA inhibited copulation (Lorrain et al., 1997). Therefore, this may be one site at which SSRI antidepressants act to inhibit sexual function. In addition, hypocretin/orexin (hcrt/orx) neurons reside in the LHA and are activated (Fos-ir) following copulation, and the numbers of hcrt/orx neurons decreased after castration (Muschamp et al., submitted). Furthermore, 5-HT inhibits hcrt/orx neurons in the LHA (Li et al., 2002). Therefore, a possible way in which LHA 5-HT inhibits sexual behavior is by inhibiting hcrt/orx neurons, which would remove their facilitative effect on VTA DA cell firing (Muschamp et al., submitted).

The nucleus paragigantocellularis (nPGi) of the medulla is a major source of inhibition of male rat sexual behavior. Lesions facilitated copulation and delayed sexual satiety (Yells et al., 1992). Similar lesions facilitated touch-based reflexes (Holmes et al., 2002; Marson et al., 1992) and allowed the urethrogenital reflex to be elicited without spinal transection (Marson and McKenna, 1990). Most of the axons projecting from the nPGi to the lumbosacral spinal cord contain 5-HT (Marson and McKenna, 1992). A 5-HT neurotoxin decreased the descending inhibition of the urethrogenital reflex, and application of 5-HT to the spinal cord suppressed that reflex in spinal-transected rats (Marson and McKenna, 1994). Thus, 5-HT from the nPGi is a major inhibitor of genital reflexes.

An ejaculation generator in the lumbar spinal cord comprises galanin- and cholecystokinin (CCK)-containing neurons, which showed Fos-ir only after ejaculating (Truitt and Coolen, 2002; Truitt et al., 2003). Lesions of these neurons severely impaired ejaculation; therefore, they not only carry ejaculation-specific sensory input to the brain, but also elicit ejaculation (Truitt and Coolen, 2003).

Description of male mouse copulatory behavior and penile reflexes

The mouse has become popular for behavioral studies, largely because of our ability to generate transgenics, knockouts, and knockdowns (see Burns-Cusato et al., 2004, for an excellent review). The male mouse begins an encounter by investigating the female's anogenital region, often lifting or pushing her with his nose. The male then presses his forepaws against the female's flanks and makes rapid, shallow pelvic thrusts. When his penis enters the female's vagina, his repeated thrusting becomes slower and deeper. After numerous intromissions, the male ejaculates, during which he may freeze for 25 seconds before dismounting or falling off

of the female. There are many strain differences in mouse mating. For example, ejaculation latencies ranged from 594 to 6943 seconds, and the numbers of intromissions preceding ejaculation ranged from 5 to 142. PEIs ranged from 17 to 60 minutes, although introduction of a novel female decreased the PEI, with some males ejaculating on the first intromission with the new female (Mosig and Dewsbury, 1976). In place preference tests both intromissions and ejaculations were shown to be rewarding (Kudwa et al., 2005).

Touch-based reflexes have also been observed in mice. Unlike rats, intact male mice did not show spontaneous reflexes while restrained with their penile sheath retracted; however, abdominal pressure did elicit erections, but not anteroflexions (Sachs, 1980). The bulbospongiosus muscle contributes to erections during intromission and especially to cups (intense erections that hold semen against the female's cervix), which are important for impregnating a female (Elmore and Sachs, 1988).

Hormonal factors in the activation of male mouse mating behavior

T is more effective than either DHT or E₂ in restoring precopulatory and copulatory behaviors in castrated mice, with sensitivity to DHT and E₂ varying widely among strains (reviewed in Burns-Cusato et al., 2004). T can also have rapid effects, as it facilitated mounting within 60 minutes in castrates (James and Nyby, 2002). Synthetic androgens (5 α -androstenediols) that can be aromatized to E, but not 5 α -reduced to DHT, were even more effective than T in restoring sexual behavior (Ogawa et al., 1996). One strain, the B6D2F1 hybrid, recovered the ability to copulate about three weeks after castration without exogenous hormones (McGill and Manning, 1976). These "continuer" males depend on E₂; although the source of the E₂ is not clear, it may be produced in the brain (Sinchak et al., 1996).

Roles of hormones in specific brain areas of male mice

Implantation of T into the MPOA completely restored ultrasonic vocalization, partially restored urine marking, and had little effect on mounting or urine preference (Sipos and Nyby, 1996). However, additional implants of T in the VTA, which were ineffective alone, produced synergistic effects on mounting and urine preference. E₂ implants in the MPOA were as effective as T (Nyby et al., 1992).

Steroid receptor mutants

The testicular feminization (*Tfm*, or androgen insensitivity) mutation in mice, as well as other animals, results from deletion of a single base in the AR gene (reviewed in Burns-Cusato et al., 2004). *Tfm* males appear phenotypically female, are infertile, and engage in no sexual behavior if tested without exogenous hormones. Small testes secrete low levels of T and DHT. However, if these males are castrated and treated with daily injections of DHT, T, E, or E + DHT, they begin to show variable amounts of sexual behavior, including occasional ejaculations (Olsen, 1992). Mice lacking the ER α (ER α KO) show little sexual behavior, even when castrated and replaced with T (Rissman et al., 1999; Wersinger and Rissman, 2000a). This is not due to a lack of hormones, as ER α KO males secrete more T than do wild-type mice, due to diminished ER-mediated negative feedback (Wersinger et al., 1997). Castration of ER α KO males and replacement with normal levels of T (Wersinger et al., 1997) or higher than normal levels of DHT (Ogawa et al., 1998) increased mounting, but did not restore ejaculation. Systemic injections of the DA agonist apomorphine restored mating and partner preference of ER α KO males to normal (Wersinger and Rissman, 2000b). However, apomorphine icv restored only mounts and intromissions (described in Burns-Cusato et al., 2004). Pubertal males lacking ER β (ER β KO) acquired the ability to ejaculate later than did WT males, but were otherwise normal (Temple et al., 2003). Males lacking both ERs did not copulate at all when gonadally intact (Ogawa et al., 2000). However, apomorphine was able to stimulate mounting in most

animals and intromitting in half; none ejaculated (described in Burns-Cusato et al., 2004). Genetic males lacking both the AR and ER α did not copulate, even after castration and replacement with T; however, the combination of E₂ replacement and systemic apomorphine did stimulate mounting in some animals (described in Burns-Cusato et al., 2004). Males lacking aromatase (ArKO) are unable to synthesize E but have normal receptors. Fewer ArKO males mounted, intromitted, and ejaculated, and had longer latencies when they did; however, about one-third of them were able to sire litters when placed with a female for a prolonged time (Bakker et al., 2002; Matsumoto et al., 2003).

Effects of systemically administered drugs on male mouse sexual behavior

Please see Table 2 for a summary of systemic drug effects on male mice and hamsters.

The roles of various brain areas in male mouse sexual behavior

Chemosensory cues are extremely important for sexual behavior in male mice (reviewed in Hull et al., 2006). However, the vomeronasal system may have an important, but not critical, role in mating. MPOA lesions severely impaired copulation in male mice, as in other species (reviewed in Hull et al., 2006). ER α KO had less nNOS-ir in the MPOA than WT or *Tfm* mice; therefore, E up-regulates nNOS-ir in mice (Scordalakes et al., 2002) as well as in rats.

Description of male hamster copulatory behavior

Mating behavior of hamsters differs in numerous ways from that of rats and mice (reviewed in Dewsbury, 1979). The female Syrian golden hamster remains in a lordosis posture continuously through successive copulations. Mating progresses more rapidly than in rats, with inter-intromission intervals of only 10 seconds and PEIs increasing from ~35 sec after the first ejaculation to ~90 sec after the ninth. Intromissions and ejaculations are longer, ~2.4 and 3.4 sec, respectively. Hamsters also have more ejaculations than rats, often 9 or 10, followed by a series of “long intromissions,” with intravaginal thrusting and no sperm transfer, prior to satiety. Detailed analysis of the hamster mating pattern, using accelerometric and polygraphic technique, revealed that trains of pelvic thrusting averaged about 1 sec, though trains associated with mounts were longer than those with intromissions and ejaculations (Arteaga & Moralí, 1997). The frequencies of pelvic thrusting averaged 15 thrusts per sec, although trains during mounts were slower. During intromissions there was a period without any thrusting, whereas during ejaculation, thrusting was of higher frequency (16.4/sec) and less vigor. Long intromissions were characterized by ~ 6 to 25 sec of slow intravaginal thrusting (1 to 2 per sec). The duration of penile insertion was longer in ejaculations than in intromissions, but was shorter in than in long intromissions.

Hormones

Lack of T during puberty impaired copulation after T replacement in adulthood, compared with castrates with T replacement during puberty (Schultz et al., 2004). Repeated sexual experiences did not compensate for these deficits. The odor of a receptive female activated Fos-ir in the MPOA even before puberty (Romeo et al., 1998), but did not increase the DA metabolite DOPAC (a measure of DA activity) until after puberty (Schultz et al., 2003). Therefore, puberty may be a second organizational period in which gonadal hormones permanently alter neural processing in areas that regulate sexual behavior (Romeo et al., 2002; Schultz et al., 2004).

Effects of systemically administered drugs in male hamsters

Please see Table 2 for a summary of systemic drug effects in mice and hamsters.

The roles of various brain areas in male hamster sexual behavior

Bilateral olfactory bulbectomy or combined deafferentation of the main and accessory olfactory systems permanently abolished sexual behavior (reviewed in Hull et al., 2006). Deafferentation of the accessory olfactory system had variable effects, with experienced males being less affected (Meredith, 1986). Mating-induced increases in Fos-ir in the main and accessory olfactory bulbs were specific to chemosensory stimuli, rather than to mating (reviewed in Hull et al., 2006).

Either T or E, but not DHT, implants into the MeA restored copulatory behavior in castrated male hamsters (Wood, 1996). Thus, hormonal activation of the MeA is sufficient for expression of sexual behavior in male hamsters. Projections from the MeA travel via the stria terminalis and ventral amygdalofugal pathway to the BNST, MPOA, and other areas. Cutting the stria terminalis delayed and slowed copulation, and combined cuts of both pathways eliminated copulation (Lehman et al., 1983).

As with many other species, the MPOA is critical for sexual behavior in male hamsters. However, steroid implants in castrates have variable effects and are not sufficient to restore behavior fully (Wood and Newman, 1995). Chemosensory cues activated Fos in the MPOA of male hamsters (Kollack-Walker and Newman, 1997). nNOS-ir is co-localized with gonadal steroid receptors in the MPOA, and castration decreased nNOS-ir (Hadeishi and Wood, 1996). As in rats, extracellular DA levels rose in the MPOA of male hamsters presented with an estrous female; this increase was blocked by bilateral or ipsilateral, but not contralateral or sham, bulbectomy (Triemstra et al., 2005).

Description of male guinea pig copulatory behavior

Male guinea pigs engage in several species-typical precopulatory behaviors, including nibbling the female's fur on her head and neck, sniffing her anogenital region, and making guttural sounds while either circling the female or shifting his weight on his two rear feet while keeping his forepaws stationary (Thornton et al., 1991). The male then approaches the female from the rear, places his chest over the female's back while clasping her sides, and begins pelvic thrusting, which usually results in a vaginal intromission (Valenstein et al., 1954). Males can intromit at a rate of approximately 1 per minute (Thornton et al., 1991), and 80% can ejaculate in a 15-min test (Butera & Czaja, 1985.) Although a male that ejaculates with a single female usually does not reinitiate copulation within the next hour, he may copulate with a different female (Grunt & Young, 1952).

Hormones

Unlike in male rats, systemically administered DHT can fully restore copulation in castrated male guinea pigs (Butera & Czaja, 1985). Furthermore, DHT implants into the MPOA were also sufficient to activate copulation in castrates (Butera and Czaja, 1989).

Summary and unanswered questions

Although there are differences in the copulatory elements among rodents, the hormonal factors and neural circuitry that control those elements are similar. Both E and DHT contribute to the activation of mating, although E is more important for copulation and DHT, for genital reflexes of rats, mice and hamsters. Hormonal activation of the MPOA is most effective, although implants in the MeA can also stimulate mounting in castrates. Chemosensory inputs from the main and accessory olfactory systems are the most important stimuli for mating, especially in hamsters, although genitosensory input via the SPFP also contributes. DA agonists facilitate sexual behavior when injected either systemically or into the MPOA or PVN. 5-HT agonists,

especially 5-HT_{1B}, tend to inhibit behavior, although 5-HT_{2C} agonists facilitate erection and 5-HT_{1A} agonists facilitate ejaculation (except in mice). norepinephrine agonists and opiates have dose-dependent effects, with low doses facilitating and high doses inhibiting behavior.

Acknowledgements

Preparation of this manuscript was supported by NIMH grants R01 MH 40826 and K02 MH 001714 to EMH.

References

- Ackerman AE, Lange GM, Clemens LG. Effects of paraventricular lesions on sex behavior and seminal emissions in male rats. *Physiol Behav* 1997;63:49–53. [PubMed: 9402614]
- Ågmo A, Paredes R. Opioids and sexual behavior in the male rat. *Pharmacol Biochem Behav* 1988;30:1021–1034. [PubMed: 3227027]
- Ågmo A, Picker Z. Catecholamines and the initiation of sexual behavior in male rats without sexual experience. *Pharmacol Biochem Behav* 1990;35:327–334. [PubMed: 2320640]
- Ahlenius S, Larsson K. Effects of selective D₁ and D₂ antagonists on male rat sexual behavior. *Experientia* 1990;46:1026–1028.
- Ahlenius S, Larsson K. Opposite effects of 5-methoxy-N, N-di-methyl-tryptamine and 5-hydroxytryptophan on male rat sexual behavior. *Pharmacol Biochem Behav* 1991;38:201–205. [PubMed: 1673250]
- Ahlenius S, Larsson K. Evidence for an involvement of 5-HT_{1B} receptors in the inhibition of male rat ejaculatory behavior produced by 5-HTP. *Psychopharmacology* 1998;137:374–382. [PubMed: 9676898]
- Ahlenius S, Larsson K, Arvidsson LE. Effects of stereoselective 5-HT_{1A} agonists on male rat sexual behavior. *Pharmacol Biochem Behav* 1989;33:691–695. [PubMed: 2587610]
- Argiolas A. Neuropeptides and sexual behaviour. *Neurosci Biobehav Rev* 1999;23:1127–1142. [PubMed: 10643821]
- Argiolas A, Melis MR. The role of oxytocin and the paraventricular nucleus in the sexual behavior of male mammals. *Physiol Behav* 2004;83:309–317. [PubMed: 15488547]
- Arteaga M, Morali G. Characteristics of the motor and genital copulatory responses of the male hamster. *J Physiol Paris* 1997;91:311–316. [PubMed: 9457664]
- Arteaga M, Motte-Lara J, Velazquez-Moctezuma J. Effects of yohimbine and apomorphine on the male sexual behaviour pattern of the golden hamster (*Mesocricetus auratus*). *Eur Neuropsychopharmacol* 2002;12:39–45. [PubMed: 11788239]
- Bakker J, Honda S, Harada N, Balthazart J. Sexual partner preference requires a functional aromatase (cyp19) gene in male mice. *Horm Behav* 2002;42:158–171. [PubMed: 12367569]
- Benelli A, Bertolini A, Poggioli R, Cavazzuti E, Calza L, Giardino L, Arletti R. Nitric oxide is involved in male sexual behavior of rats. *Eur J Pharmacol* 1995;294:505–510. [PubMed: 8750712]
- Beyer C, Contreras G, Morali G, Larsson K. Effects of castration and sex steroid treatment on the motor copulatory pattern in the rat. *Physiol Behav* 1981;27:727–730. [PubMed: 7323176]
- Bialy M, Beck J, Abramczyk P, Trzebski A, Przybylski J. Sexual behavior in male rats after nitric oxide synthesis inhibition. *Physiol Behav* 1996;60:139–143. [PubMed: 8804654]
- Boscarino BT, Parfitt DB. Chronic oral administration of clomipramine decreases sexual behavior in the male Syrian hamster (*Mesocricetus auratus*). *Physiol Behav* 2002;75:361–366. [PubMed: 11897263]
- Burnett AL, Nelson RJ, Calvin DC, Liu JX, Demas GE, Klein SL, Kriegsfeld LJ, Dawson TM, Snyder SH. Nitric oxide-dependent penile erection in mice lacking neuronal nitric oxide synthase. *Mol Med* 1996;2:288–296. [PubMed: 8784782]
- Burns-Cusato M, Scordalakes EM, Rissman EF. Of mice and missing data: what we know (and need to learn) about male sexual behavior. *Physiol Behav* 2004;83:217–232. [PubMed: 15488541]
- Butera PC, Czaja JA. Maintenance of target tissue and sexual behavior with dihydrotestosterone in male rats and guinea pigs. *Physiol Behav* 1985;34:319–321. [PubMed: 4001193]
- Butera PC, Czaja JA. Effects of intracranial implants of dihydrotestosterone in the reproductive physiology and behavior of male guinea pigs. *Horm Behav* 1989;23:424–431. [PubMed: 2793082]

- Cantor JM, Binik YM, Pfaus JG. Chronic fluoxetine inhibits sexual behavior in the male rat: reversal with oxytocin. *Psychopharmacology* 1999;144:355–362. [PubMed: 10435408]
- Chambers KC, Thornton JE, Roselli CE. Age-related deficits in brain androgen binding and metabolism, testosterone, and sexual behavior of male rats. *Neurobiol Aging* 1991;12:123–130. [PubMed: 2052127]
- Clark, JT. Sexual arousal and performance are modulated by adrenergic-neuropeptide-steroid interactions. In: Bancroft, J., editor. *The Pharmacology of Sexual Function and Dysfunction: Proceedings of the Esteve Foundation Symposium VI*; Son Vida, Mallorca. 9–12 October 1994; Amsterdam: Excerpta Medica; 1995. p. 55–68.
- Clark JT, Smith ER. Clonidine suppresses copulatory behavior and erectile reflexes in male rats: Lack of effect of naloxone pretreatment. *Neuroendocrinology* 1990;51:357–364.
- Clark JT, Smith ER, Davidson JM. Enhancement of sexual motivation in male rats by yohimbine. *Science* 1984;225:847–849. [PubMed: 6474156]
- Clark JT, Smith ER, Davidson JM. Evidence for the modulation of sexual behavior by α -adrenoceptors in male rats. *Neuroendocrinology* 1985;41:36–43. [PubMed: 2991794]
- Cooke BM, Breedlove SM, Jordan C. Both estrogen receptors and androgen receptors contribute to testosterone-induced changes in the morphology of the medial amygdala and sexual arousal in male rats. *Horm Behav* 2003;43:335–346.
- Cross E, Roselli CE. 17 β -estradiol rapidly facilitates chemoinvestigation and mounting in castrated male rats. *Am J Physiol* 1999;276(5 Pt 2):R1346–1350. [PubMed: 10233026]
- Dewsbury, DA. Description of sexual behavior in research on hormone-behavior interactions. In: Beyer, C., editor. *Endocrine Control of Sexual Behavior*. Raven Press; NY: 1979. p. 3–32.
- Dominguez JM, Balfour ME, Coolen LM. Copulation-induced activation of NMDA receptor containing neurons in the medial preoptic nucleus. *Abst Soc Behav Neuroendocrinol Horm Behav* 2003;44:46.
- Dominguez JM, Gil M, Hull EM. Preoptic glutamate facilitates male sexual behavior. *J Neurosci* 2006;26:1699–1703. [PubMed: 16467517]
- Dominguez JM, Hull EM. Stimulation of the medial amygdala enhances medial preoptic dopamine release: implications for male rat sexual behavior. *Brain Res* 2001;917:225–229. [PubMed: 11640908]
- Dominguez JM, Hull EM. Dopamine, the medial preoptic area, and male sexual behavior. *Physiol Behav* 2005;86:356–68. [PubMed: 16135375]
- Dominguez JM, Muschamp JW, Schmich JM, Hull EM. Nitric oxide mediates glutamate-evoked dopamine release in the medial preoptic area. *Neuroscience* 2004;125:203–210. [PubMed: 15051159]
- Dominguez J, Riolo JV, Xu Z, Hull EM. Regulation by the medial amygdala of copulation and medial preoptic dopamine release. *J Neurosci* 2001;21:349–355. [PubMed: 11150352]
- Du J, Hull EM. Effects of testosterone on neuronal nitric oxide synthase and tyrosine hydroxylase. *Brain Res* 1999;836:90–98. [PubMed: 10415408]
- Edwards DA, Einhorn LC. Preoptic and midbrain control of sexual motivation. *Physiol Behav* 1986;37:329–335. [PubMed: 3488558]
- Elmore LA, Sachs BD. Role of the bulbospongiosus muscles in sexual behavior and fertility in the house mouse. *Physiol Behav* 1988;44:125–129. [PubMed: 3237806]
- Everitt BJ. Sexual motivation: A neural and behavioral analysis of the mechanisms underlying appetitive and copulatory responses of male rats. *Neurosci Biobehav Rev* 1990;14:217–232. [PubMed: 2190121]
- Fernandez-Fewell GD, Meredith M. Facilitation of mating behavior in male hamsters by LHRH and AcLHRH5–10: interaction with the vomeronasal system. *Physiol Behav* 1995;57:213–21. [PubMed: 7716195]
- Ferrari F, Ottani A, Giuliani D. Influence of sildenafil on central dopamine-mediated behaviour in male rats. *Life Sci* 2002;70:1501–1508. [PubMed: 11895101]
- Frank JL, Hendricks SE, Olson CH. Multiple ejaculations and chronic fluoxetine: effects on male rat copulatory behavior. *Pharmacol Biochem Behav* 2000;66:337–342. [PubMed: 10880687]

- Giuliano F. Control of penile erection by the melanocortinergic system: experimental evidences and therapeutic perspectives. *J Androl* 2004;25:683–694. [PubMed: 15292097]
- Giuliano F, Bernabe J, Alexandre L, Niewoehner U, Haning H, Bischoff E. Pro-erectile effect of vardenafil: in vitro experiments in rabbits and in vivo comparison with sildenafil in rats. *Eur Urol* 2003;44:731–736. [PubMed: 14644128]
- Giuliano F, Rampin O, Brown K, Courtois F, Benoit G, Jardin A. Stimulation of the medial preoptic area of the hypothalamus in the rat elicits increases in intracavernous pressure. *Neurosci Lett* 1996;209:1–4. [PubMed: 8734895]
- Grunt JA, Young WC. Psychological modification of fatigue following orgasm (ejaculation) in the male guinea pig. *J Comp Physiol Psychol* 1952;45:508–510. [PubMed: 13000021]
- Hadeishi Y, Wood RI. Nitric oxide synthase in mating behavior circuitry of male Syrian hamster brain. *J Neurobiol* 1996;30:480–492. [PubMed: 8844512]
- Holmes GM, Chapple WD, Leipheimer RE, Sachs BD. Electromyographic analysis of male rat perineal muscles during copulation and reflexive erections. *Physiol Behav* 1991;49:1235–1246. [PubMed: 1896506]
- Holmes GM, Hermann GE, Rogers RC, Bresnahan JC, Beattie MS. Dissociation of the effects of nucleus raphe obscurus or rostral ventrolateral medulla lesions on eliminatory and sexual reflexes. *Physiol Behav* 2002;75:49–55. [PubMed: 11890952]
- Hull EM, Du J, Lorrain DS, Matuszewich L. Extracellular dopamine in the medial preoptic area: implications for sexual motivation and hormonal control of copulation. *J Neurosci* 1995;15:7465–7471. [PubMed: 7472498]
- Hull EM, Lumley LA, Matuszewich L, Dominguez J, Moses J, Lorrain DS. The roles of nitric oxide in sexual function of male rats. *Neuropharmacology* 1994;33:1499–1504. [PubMed: 7532834]
- Hull, EM.; Meisel, RL.; Sachs, BD. Male sexual behavior. In: Pfaff, DW.; Arnold, AP.; Etgen, AM.; Fahrbach, SE.; Rubin, RT., editors. *Hormones, Brain and Behavior*. Academic Press; 2002. p. 3-137.
- Hull, EM.; Wood, RI.; McKenna, KE. The neurobiology of male sexual behavior. In: Neill, J.; Donald, Pfaff, editors. *The Physiology of Reproduction*. 3. Elsevier Press; 2006. p. 1729-1824.
- James PJ, Nyby JG. Testosterone rapidly affects the expression of copulatory behavior in house mice (*Mus musculus*). *Physiol Behav* 2002;75:287–294. [PubMed: 11897254]
- Kollack-Walker S, Newman SW. Mating-induced expression of c-fos in the male Syrian hamster brain: Role of experience, pheromones, and ejaculations. *J Neurobiol* 1997;32:481–501. [PubMed: 9110260]
- Kondo Y, Arai Y. Functional association between the medial amygdala and the medial preoptic area in regulation of mating behavior in the male rat. *Physiol Behav* 1995;57:69–73. [PubMed: 7878127]
- Krey LC, McGinnis MY. Time-courses of the appearance/disappearance of nuclear androgen + receptor complexes in the brain and adenohypophysis following testosterone administration/withdrawal to castrated male rats: Relationships with gonadotropin secretion. *J Steroid Biochem* 1990;35:403–408. [PubMed: 2109153]
- Kriegsfeld LJ, Demas GE, Huang PL, Burnett AL, Nelson RJ. Ejaculatory abnormalities in mice lacking the gene for endothelial nitric oxide synthase (eNOS^{-/-}). *Physiol Behav* 1999;67:561–566. [PubMed: 10549894]
- Kudwa AE, Dominguez-Salazar E, Cabrera DM, Sibley DR, Rissman EF. Dopamine D5 receptor modulates male and female sexual behavior in mice. *Psychopharmacology* 2005;180:206–14. [PubMed: 15696326]
- Lagoda G, Muschamp JM, Vigdorichik A, Hull EM. A nitric oxide synthase inhibitor in the medial preoptic area inhibits copulation and stimulus sensitization in male rats. *Behav Neurosci* 2004;118:1317–1323. [PubMed: 15598140]
- Lehman MN, Powers JB, Winans SS. Stria terminalis lesions alter the temporal pattern of copulatory behavior in the male golden hamster. *Behav Brain Res* 1983;8:109–128. [PubMed: 6849676]
- Leipheimer RE, Sachs BD. GABAergic regulation of penile reflexes and copulation in rats. *Physiol Behav* 1988;42:351–357. [PubMed: 2838857]
- Leyton M, Stewart J. Acute and repeated activation of male sexual behavior by tail pinch: opioid and dopaminergic mechanisms. *Physiol Behav* 1996;60:77–85. [PubMed: 8804646]

- Liu YC, Salamone JD, Sachs BD. Impaired sexual response after lesions of the paraventricular nucleus of the hypothalamus in male rats. *Behav Neurosci* 1997a;111:1361–1367. [PubMed: 9438804]
- Liu YC, Salamone JD, Sachs BD. Lesions in medial preoptic area and bed nucleus of stria terminalis: Differential effects on copulatory behavior and noncontact erection in male rats. *J Neurosci* 1997b; 17:5245–5253. [PubMed: 9185562]
- Lopez HH, Ettenberg A. Haloperidol challenge during copulation prevents subsequent increase in male sexual motivation. *Pharmacol Biochem Behav* 2000;67:387–393. [PubMed: 11124405]
- Lopez HH, Ettenberg A. Dopamine antagonism attenuates the unconditioned incentive value of estrous female cues. *Pharmacol Biochem Behav* 2001;68:411–416. [PubMed: 11325393]
- Lopez HH, Ettenberg A. Exposure to female rats produces differences in c-fos induction between sexually-naïve and experienced male rats. *Brain Res* 2002a;947:57–66. [PubMed: 12144853]
- Lopez HH, Ettenberg A. Sexually conditioned incentives: Attenuation of motivational impact during dopamine receptor antagonism. *Pharmacol Biochem Behav* 2002b;72:65–72. [PubMed: 11900770]
- Lorrain DS, Matuszewich L, Friedman RD, Hull EM. Extracellular serotonin in the lateral hypothalamic area is increased during postejaculatory interval and impairs copulation in male rats. *J Neurosci* 1997;17:9361–9366. [PubMed: 9364081]
- Lorrain DS, Riolo JV, Matuszewich L, Hull EM. Lateral hypothalamic serotonin inhibits nucleus accumbens dopamine: implications for sexual refractoriness. *J Neurosci* 1999;19:7648–7652. [PubMed: 10460270]
- Lumley LA, Hull EM. Effects of a D₁ antagonist and of sexual experience on copulation-induced Fos-like immunoreactivity in the medial preoptic nucleus. *Brain Res* 1999;829:55–68. [PubMed: 10350530]
- Maeda N, Matsuoka N, Yamaguchi I. Septohippocampal cholinergic pathway and penile erections induced by dopaminergic and cholinergic stimulants. *Brain Res* 1990;537:163–168. [PubMed: 1982238]
- Maillard CA, Edwards DA. Excitotoxin lesions of the zone incerta/lateral tegementum continuum: Effects on male sexual behavior in rats. *Behav Brain Res* 1991;46:143–149. [PubMed: 1664729]
- Malmnas CO. The significance of dopamine, versus other catecholamines, for L-dopa induced facilitation of sexual behavior in the castrated male rat. *Pharmacol Biochem Behav* 1976;4:521–526. [PubMed: 951430]
- Manzo J, Cruz MR, Hernandez ME, Pacheco P, Sachs BD. Regulation of noncontact erection in rats by gonadal steroids. *Horm Behav* 1999;35:264–270. [PubMed: 10373338]
- Markowski VP, Hull EM. Cholecystokinin modulates mesolimbic dopaminergic influences on male rat copulatory behavior. *Brain Res* 1995;699:266–274. [PubMed: 8616630]
- Marson L, McKenna KE. The identification of a brainstem site controlling spinal sexual reflexes in male rats. *Brain Res* 1990;515:303–308. [PubMed: 2357567]
- Marson L, McKenna KE. A role for 5-hydroxytryptamine in descending inhibition of spinal sexual reflexes. *Exp Brain Res* 1992;88:313–320. [PubMed: 1577105]
- Marson L, McKenna KE. Serotonergic neurotoxic lesions facilitate male sexual reflexes. *Pharmacol Biochem Behav* 1994a;47:883–888. [PubMed: 7518085]
- Marson L, McKenna KE. Stimulation of the hypothalamus initiates the urethrogenital reflex in male rats. *Brain Res* 1994b;638:103–108. [PubMed: 8199850]
- Marson L, List MS, McKenna KE. Lesions of the nucleus paragigantocellularis alter ex copula penile reflexes. *Brain Res* 1992;592:187–192. [PubMed: 1450909]
- Mas M, Fumero B, Perez-Rodriguez I. Induction of mating behavior by apomorphine in sexually sated rats. *Euro J Pharmacol* 1995;280:331–334.
- Matsumoto T, Honda S, Harada N. Alteration in sex-specific behaviors in male mice lacking the aromatase gene. *Neuroendocrinology* 2003;77:416–424. [PubMed: 12845227]
- McGill TE, Manning A. Genotype and retention of the ejaculatory reflex in castrated male mice. *Anim Behav* 1976;24:507–518. [PubMed: 962190]
- McGinnis MY, Dreifuss RM. Evidence for a role of testosterone-androgen receptor interactions in mediating masculine sexual behavior in male rats. *Endocrinology* 1989;124:618–626. [PubMed: 2912689]

- McGinnis MY, Mirth MC, Zebrowski AF, Dreifuss RM. Critical exposure time for androgen activation of male sexual behavior in rats. *Physiol Behav* 1989;46:159–165. [PubMed: 2602453]
- Meisel, RL.; Sachs, BD. The physiology of male sexual behavior. In: Knobil, E.; Neill, JD., editors. *Physiology of Reproduction*. 2. Raven Press; New York: 1994. p. 3-106.
- Meisel RL, O'Hanlon JK, Sachs BD. Differential maintenance of penile responses and copulatory behavior by gonadal hormones in castrated male rats. *Horm Behav* 1984;18:56–64. [PubMed: 6706319]
- Melis MR, Stancampiano R, Argiolas A. Hippocampal oxytocin mediates apomorphine-induced penile erection and yawning. *Pharm Biochem Behav* 1992;42:61–66.
- Melis MR, Succu S, Mascia MS, Cortis L, Argiolas A. Extra-cellular dopamine increases in the paraventricular nucleus of the hypothalamus: correlation with penile erection and yawning. *Eur J Neurosci* 2003;17:1266–1272. [PubMed: 12670314]
- Melis MR, Succu S, Mascia MS, Cortis L, Argiolas A. Extracellular excitatory amino acids increase in the paraventricular nucleus of male rats during sexual activity: main role of N-methyl-D-aspartic acid receptors in erectile function. *Eur J Neurosci* 2004;19:2569–2575. [PubMed: 15128410]
- Melis MR, Succu S, Mauri A, Argiolas A. Nitric oxide production is increased in the paraventricular nucleus of the hypothalamus of male rats during non-contact penile erections and copulation. *Eur J Neurosci* 1998;10:1968–1974. [PubMed: 9753084]
- Meredith M. Vomeronasal organ removal before sexual experience impairs male hamster mating behavior. *Physiol Behav* 1986;36:737–743. [PubMed: 3714848]
- Moses J, Hull EM. A nitric oxide synthesis inhibitor administered into the medial preoptic area increases seminal emissions in an *ex copula* reflex test. *Pharmacol Biochem Behav* 1999;63:345–348. [PubMed: 10418773]
- Mosig DW, Dewsbury DA. Studies of the copulatory behavior of house mice (*Mus musculus*). *Behav Biol* 1976;16:463–473. [PubMed: 989316]
- Nyby J, Matochik JA, Barfield RJ. Intracranial androgenic and estrogenic stimulation of male-typical behaviors in house mice, *Mus domesticus*. *Horm Behav* 1992;26:24–45. [PubMed: 1563726]
- Ogawa S, Chester AE, Hewitt SC, Walker VR, Gustafsson JA, Smithies O. Abolition of male sexual behaviors in mice lacking estrogen receptors alpha and beta (alpha beta ERKO). *Proc Natl Acad Sci U S A* 2000;97:14737–14741. [PubMed: 11114183]
- Ogawa S, Robbins A, Kumar N, Pfaff DW, Sundaram K, Bardin CW. Effects of testosterone and 7 alpha-methyl-19-nortestosterone (MENT) on sexual and aggressive behaviors in two inbred strains of male mice. *Horm Behav* 1996;30:74–84. [PubMed: 8724182]
- Ogawa S, Washburn TF, Taylor J, Lubahn DB, Korach KS, Pfaff DW. Modification of testosterone-dependent behaviors by estrogen receptor-alpha gene disruption in male mice. *Endocrinology* 1998;139:5058–5069. [PubMed: 9832445]
- O'Hanlon JK, Meisel RL, Sachs BD. Estradiol maintains castrated male rats' sexual reflexes in copula, but not ex copula. *Behav Neur Biol* 1981;32:269–273.
- Olsen, KL. Genetic influence on sexual behavior differentiation. In: Gerall, AA.; Moltz, H.; Ward, IL., editors. *Sexual differentiation, handbook of behavioral neurobiology*. Plenum Press; New York: 1992. p. 1-40.
- Paredes RG, Highland L, Karam P. Socio-sexual behavior in male rats after lesions of the medial preoptic area: evidence for reduced sexual motivation. *Brain Res* 1993;618:271–276. [PubMed: 8374757]
- Paredes RG, Tzschenke T, Nakach N. Lesions of the medial preoptic area/anterior hypothalamus (MPOA/AH) modify partner preference in male rats. *Brain Res* 1998;813:81–83.
- Pehek EA, Thompson JT, Hull EM. The effects of intrathecal administration of the dopamine agonist apomorphine on penile reflexes and copulation in the male rat. *Psychopharmacology* 1989b; 99:304–308. [PubMed: 2574477]
- Pfaus JG, Wilkins MF. A novel environment disrupts copulation in sexually naïve but not experienced male rats: reversal with naloxone. *Physiol Behav* 1995;57:1045–1049. [PubMed: 7652023]
- Popova NK, Amstislavskaya TG. Involvement of the 5-HT(1A) and 5-HT(1B) serotonergic receptor subtypes in sexual arousal in male mice. *Psychoneuroendocrinol* 2002;27:609–618.
- Putnam SK, Sato S, Hull EM. Hormonal maintenance of copulation in castrates: association with extracellular dopamine in MPOA. *Horm Behav* 2003;44:419–426. [PubMed: 14644636]

- Putnam SK, Sato S, Hull EM. Effects of testosterone metabolites on copulation, medial preoptic dopamine content, and nitric oxide synthase. *Horm Behav* 2005;47:513–522. [PubMed: 15811352]
- Rampin O, Jerome N, Suaudeau C. Proerectile effects of apomorphine in mice. *Life Sci* 2003;72:2329–2336. [PubMed: 12639699]
- Rissman EF, Wersinger SR, Fugger HN, Foster TC. Sex with knockout models: behavioral studies of estrogen receptor alpha. *Brain Res* 1999;835:80–90. [PubMed: 10448199]
- Rodriguez-Manzo G. Yohimbine interacts with the dopaminergic system to reverse sexual satiation: further evidence for a role of sexual motivation in sexual exhaustion. *Eur J Pharmacol* 1999;372:1–8. [PubMed: 10374708]
- Rodriguez-Manzo G, Fernandez-Guasti A. Participation of the central noradrenergic system in the reestablishment of copulatory behavior of sexually exhausted rats by yohimbine, naloxone, and 8-OH-DPAT. *Brain Res Bull* 1995;38:399–404. [PubMed: 8535863]
- Rodriguez-Manzo G, Lopez-Rubalcava C, Hen R, Fernandez-Guasti A. Participation of 5-HT(1B) receptors in the inhibitory actions of serotonin on masculine sexual behaviour of mice: pharmacological analysis in 5-HT(1B) receptor knock-out mice. *Brit J Pharmacol* 2002;136:1127–1134. [PubMed: 12163345]
- Rodríguez-Manzo G, Pellicer F, Larsson K, Fernandez-Guasti A. Stimulation of the medial preoptic area facilitates sexual behavior but does not reverse sexual satiation. *Behav Neurosci* 2000;114:553–560. [PubMed: 10883805]
- Romeo RD, Parfitt DB, Richardson HN, Sisk CL. Pheromones elicit equivalent levels of Fos-immunoreactivity in prepubertal and adult male Syrian hamsters. *Horm Behav* 1998;34:48–55. [PubMed: 9735228]
- Romeo RD, Richardson HN, Sisk CL. Puberty and the maturation of the male brain and sexual behavior: recasting a behavioral potential. *Neurosci Biobehav Rev* 2002;26:381–391. [PubMed: 12034137]
- Roselli CE, Thornton JE, Chambers KC. Age-related deficits in brain estrogen receptors and sexual behavior of male rats. *Behav Neurosci* 1993;107:202–209. [PubMed: 8447953]
- Sachs BD. Sexual reflexes of spinal male house mice. *Physiol Behav* 1980;24:489–492. [PubMed: 7375569]
- Sachs, BD. Potency and fertility: hormonal and mechanical causes and effects of penile actions in rats. In: Balthazart, J.; Pröve, E.; Gilles, R., editors. *Hormones and Behaviour in Higher Vertebrates*. Springer-Verlag; Berlin: 1983. p. 86–110.
- Sachs BD, Bitran D. Spinal block reveals roles for brain and spinal cord in the mediation of reflexive erection in rats. *Brain Res* 1990;528:99–108. [PubMed: 2245341]
- Sachs BD, Valcourt RJ, Flagg HC. Copulatory behavior and sexual reflexes of male rats treated with naloxone. *Pharmacol Biochem Behav* 1981;14:251–253. [PubMed: 7208564]
- Sala M, Braida D, Leone MP, Calcaterra P, Monti S, Gori E. Central effect of yohimbine on sexual behavior in the rat. *Physiol Behav* 1990;47:165–173. [PubMed: 2326333]
- Sato Y, Wada H, Horita H, Suzuki N, Shibuya A, Adachi H, Kato R, Tsukamoto T, Kumamoto Y. Dopamine release in the medial preoptic area during male copulatory behavior in rats. *Brain Res* 1995;692:66–70. [PubMed: 8548321]
- Scaletta LL, Hull EM. Systemic or intracranial apomorphine increases copulation in long-term castrated male rats. *Pharm Biochem Behav* 1990;37:471–475.
- Schnur SL, Smith ER, Lee RL, Mas M, Davidson JM. A component analysis of the effects of DPAT on male rat sexual behavior. *Physiol Behav* 1989;45:897–901. [PubMed: 2528754]
- Schultz KM, Richardson HN, Romeo RD, Morris JA, Lookingland KJ, Sisk CL. Medial preoptic area dopaminergic responses to female pheromones develop during puberty in the male Syrian hamster. *Brain Res* 2003;988:139–145. [PubMed: 14519535]
- Schultz KM, Richardson HN, Zehr JL, Osetek AJ, Menard TA, Sisk CL. Gonadal hormones masculinize and defeminize reproductive behaviors during puberty in the male Syrian hamster. *Horm Behav* 2004;45:242–249. [PubMed: 15053940]
- Scordalakes EM, Shetty SJ, Rissman EF. Roles of estrogen receptor alpha and androgen receptor in the regulation of neuronal nitric oxide synthase. *J Comp Neurol* 2002;453:336–344. [PubMed: 12389206]

- Shimura T, Yamamoto T, Shimokochi M. The medial preoptic area is involved in both sexual arousal and performance in male rats: re-evaluation of neuron activity in freely moving animals. *Brain Res* 1994;640:215–222. [PubMed: 8004448]
- Simerly RB, Swanson LW. The organization of neural inputs to the medial preoptic nucleus of the rat. *J Comp Neurol* 1986;246:312–342. [PubMed: 3517086]
- Simerly RB, Swanson LW. Projections of the medial preoptic nucleus: a Phaseolis vulgaris leucoagglutinin anterograde tract-tracing study in the rat. *J Comp Neurol* 1988;270:209–242. [PubMed: 3259955]
- Sinchak K, Roselli CE, Clemens LG. Levels of serum steroids, aromatase activity, and estrogen receptors in preoptic area, hypothalamus, and amygdala of B6D2F1 male house mice that differ in the display of copulatory behavior after castration. *Behav Neurosci* 1996;110:593–602. [PubMed: 8889004]
- Sipos ML, Nyby JG. Concurrent androgenic stimulation of the ventral tegmental area and medial preoptic area: synergistic effects on male-typical reproductive behaviors in house mice. *Brain Res* 1996;729:29–44. [PubMed: 8874874]
- Smith ER, Lee RL, Schnur SL, Davidson JM. Alpha 2-adrenoceptor antagonists and male sexual behavior: I. Mating behavior. *Physiol Behav* 1987a;41:7–14. [PubMed: 2891151]
- Steers WD, de Groat WC. Effects of m-chlorophenylpiperazine on penile and bladder function in rats. *Am J Physiol* 1989;257:R1441–1449. [PubMed: 2604003]
- Sugiura K, Yoshimura H, Yokoyama M. An animal model of copulatory disorder induced by social stress in male mice: effects of apomorphine and L-dopa. *Psychopharmacology* 1997;133:249–255. [PubMed: 9361330]
- Szczycka MS, Zhou QY, Palmiter RD. Dopamine-stimulated sexual behavior is testosterone dependent in mice. *Behav Neurosci* 1998;112:1229–1235. [PubMed: 9829800]
- Tallentire D, McRae G, Spedding R, Clark R, Vickery B. Modulation of sexual behaviour in the rat by a potent and selective 2-adrenoceptor antagonist, delequamine, (RS-15385–197). *Brit J Pharm* 1996;118:63–72.
- Temple JL, Scordalakes EM, Bodo C, Gustafsson JA, Rissman EF. Lack of functional estrogen receptor beta gene disrupts pubertal male sexual behavior. *Horm Behav* 2003;44:427–434. [PubMed: 14644637]
- Thornton JE, Irving S, Goy RW. Effects of prenatal antiandrogen treatment on masculinization and defeminization of guinea pigs. *Physiol Behav* 1991;50:471–475. [PubMed: 1800997]
- Valenstein ES, Riss W, Young WC. Sex drive in genetically heterogeneous and highly inbred strains of male guinea pigs. *J Comp Physiol Psychol* 1954;47:162–165. [PubMed: 13152234]
- Triemstra JL, Nagatani S, Wood RI. Chemosensory cues are essential for mating-induced dopamine release in MPOA of male Syrian hamsters. *Neuropsychopharmacology* 2005;30:1436–1442. [PubMed: 15702137]
- Truitt WA, Coolen LM. Identification of a potential ejaculation generator in the spinal cord. *Science* 2002;297:1566–1569. [PubMed: 12202834]
- Truitt WA, Shipley MT, Veening JG, Coolen LM. Activation of a subset of lumbar spinothalamic neurons after copulatory behavior in male but not female rats. *J Neurosci* 2003;23:325–31. [PubMed: 12514231]
- Vagell ME, McGinnis MY. The role of aromatization in the restoration of male rat reproductive behavior. *J Neuroendocrinol* 1997;9:415–421. [PubMed: 9229352]
- van Furth WR, van Ree JM. Endogenous opioids and sexual motivation and performance during the light phase of the diurnal cycle. *Brain Res* 1994;636:175–179. [PubMed: 8156406]
- Vega Matuszyk J, Larsson K, Eriksson E. The selective serotonin reuptake inhibitor fluoxetine reduces sexual motivation in male rats. *Pharmacol Biochem Behav* 1998;60:527–532. [PubMed: 9632236]
- Wersinger SR, Rissman EF. Oestrogen receptor alpha is essential for female-directed chemo-investigatory behaviour but is not required for the pheromone-induced luteinizing hormone surge in male mice. *J Neuroendocrinol* 2000a;12:103–110. [PubMed: 10718905]
- Wersinger SR, Rissman EF. Dopamine activates masculine sexual behavior independent of the estrogen receptor alpha. *J Neurosci* 2000b;20:4248–4254. [PubMed: 10818161]

- Wersinger SR, Sannen K, Villalba C, Lubahn DB, Rissman EF, De Vries GJ. Masculine sexual behavior is disrupted in male and female mice lacking a functional estrogen receptor alpha gene. *Horm Behav* 1997;32:176–183. [PubMed: 9454668]
- Westberry J, Meredith M. The influence of chemosensory input and gonadotropin releasing hormone on mating behavior circuits in male hamsters. *Brain Res* 2003;974:1–16. [PubMed: 12742619]
- Witt DM, Insel TR. Increased Fos expression in oxytocin neurons following masculine sexual behavior. *J Neuroendocrinol* 1994;6:13–18. [PubMed: 8025564]
- Wood RI. Estradiol, but not dihydrotestosterone, in the medial amygdala facilitates male hamster sex behavior. *Physiol Behav* 1996;59:833–841. [PubMed: 8778874]
- Wood RI, Newman SW. Integration of chemosensory and hormonal cues is essential for mating in the male Syrian hamster. *J Neurosci* 1995;15:7261–7269. [PubMed: 7472480]
- Yamada K, Emson P, Hokfelt T. Immunohistochemical mapping of nitric oxide synthase in the rat hypothalamus and colocalization with neuropeptides. *J Chem Neuroanat* 1996;10:295–316. [PubMed: 8811420]
- Yells DP, Hendricks SE, Prendergast MA. Lesions of the nucleus paragigantocellularis effects on mating behavior in male rats. *Brain Res* 1992;596:73–79. [PubMed: 1468004]
- Zarrindast MR, Mamanpush SM, Rashidy-Pour A. Morphine inhibits dopaminergic and cholinergic induced ejaculation in rats. *Gen Pharmacol* 1994;25:803–808. [PubMed: 7958745]

copulation in short-term (Malmnas, 1976) and long-term (Scaletta & Hull, 1990) castrated rats.
rats appear to facilitative male sexual behavior, high levels of peripheral NE activity inhibits erections (Stefanick et al, 1983) by vasoconstricting penile arteries.
promote ejaculation, but inhibit erections. facilitate erections, but inhibit ejaculations. inhibit ejaculation
muscarinic agonists may facilitate copulation and erections. although contradictory results have been reported. These effects may be mediated, in part, by peripheral influences on penile muscles and/or the parasympathetic nervous system.
these apparently contradictory effects is that low to moderate levels of endogenous opioids may facilitate sexual motivation and genital reflexes; however, exogenous opiates or high levels of endogenous opioids may inhibit those same functions.
literator of parasympathetic mediated erections, it may inhibit sympathetically mediated seminal emission and ejaculation (Moses & Hull, 1999)

behavior

	REMARKS
umpin et al., 2003; Niiikura et al., 2002; Sczypka, 1998) 2b)	DA-deficient mice were activated by L-DOPA and were more sensitive to L-DOPA and T (Sczypka et al., 1998).
)	Increasing NE release with an α_2 autoreceptor agonist facilitated copulation.
; Popova and Amstislavskaya, 2002) 2002)	5-HT _{1B} knock-out mice were less impaired by serotonergic drugs, but needed more stimulation to ejaculate, possibly because of compensatory mechanisms (Rodriguez-Manzo, 2002).
egsfeld et al., 1999)	nNOS knockouts had normal erections, due to increased eNOS, but ejaculated with fewer intromissions; thus NO may help prevent "premature ejaculation."
nd Meredith, 1995)	GnRH is released in response to female hamster odor.

Horn Behav. Author manuscript; available in PMC 2007 August 27.