was designed to answer two questions: (1) Is methaqualone a reliable hypnotic? (2) Is the action of this drug comparable with that of the usual dose of cyclobarbitone? Other aspects of the pharmacological action of methaqualone were not studied. It is emphasized that the conclusions reached depend entirely on patients' opinions; but the reliability of their assessments was checked by the technique of investigation, and this included the use of a placebo.

FIG. 3.—Trial 3, showing 300 mg. of methaqualone to be superior to 200 mg. of cyclobarbitone.

The main conclusions are that methaqualone in a single dose of 150 mg. is a reliable hypnotic; and no important difference can be detected between the hypnotic action of 150 mg. of methaqualone and 200 mg. of cyclobarbitone.

We are grateful to Professor S. Alstead for permission to study the patients under his care. We are pleased to acknowledge assistance from Mr. P. S. Waldie, who prepared the figures. The tablets used in the trials were kindly supplied by Boots Pure Drug Company Limited.

REFERENCES

- Bross, I. (1952). *Biometrics*, **8**, 188.
Ravina, A. (1959). *Presse méd.*, **67**, 891.
Thomson, T. J. (1958). *Brit. med. J.*, **2**, 1140.

Applications of radioisotopes in the study of tropical diseases were reviewed at an international symposium held jointly by the International Atomic Energy Agency and W.H.O. in Bangkok from December 12 to 16, 1960. The meeting showed that the nature and causes of certain types of anaemia widely prevalent in tropical areas are now being investigated with radioactive isotopes of iron and chromium, while radioactive iodine has become indispensable in efforts to determine the causative factors of endemic goitre. As for diseases caused by insects such as malaria or those caused by parasites such as hookworm, radioisotopes help the study of the physiology, biochemistry, and behaviour pattern of insects and parasites concerned and in devising effective methods of attacking them. The application of isotopes in the investigation of disorders due to nutritional deficiencies was also discussed at the meeting.

FURTHER EXPERIENCE WITH METHOTREXATE IN CHORIONCARCINOMA

BY

A. E. R. BUCKLE, M.B., B.S., F.R.C.S., M.R.C.O.G.

Senior Registrar in Obstetrics and Gynaecology, King's College Hospital, London, and the Woolwich Group of Hospitals

The demonstration of the inhibition of oestrogen-induced growth of the female genital tract in the experimental animal by folic-acid antagonists (Hertz and Sebrell, 1944; Hertz, 1948a, 1948b; Hertz and Tullner, 1949) and the effect of aminopterin in inducing foetal death *in utero* (Thiersch, 1952) was followed by the use of the folic-acid antagonist methotrexate (4-amino-N-10-methylpteroylglutamic acid), in the treatment of chorioncarcinoma (Li *et al.*, 1956, 1958). Since that time, reports have come from America on the use of this drug in a larger series of cases (Hertz *et al.*, 1958, 1959), and a small number of cases have been recorded from this country (Doniach *et al.*, 1958; Buckle, 1959; Bagshawe and Brooks, 1959; Bagshawe and McDonald, 1960).

In a previous paper (Buckle, 1959) a case was recorded in which an initial response to the drug was followed by drug resistance and death, the plea being made at that time for the collection of such cases in suitable centres in this country in order that this form of therapy could be assessed. A similar suggestion has also recently been made (*Brit. med. J.*, 1960). Although the value of recording single cases of tumour chemotherapy is generally of little value, exception may be made in the case of chorioncarcinoma in view of its rarity, and a further case is recorded.

Case Report

A gravida-2, aged 20, was admitted to hospital on December 17, 1959, at 30 weeks pregnancy, with painless, bright-red vaginal bleeding. She had had an idiopathic thoracic scoliosis at the age of 14 which had not progressed, although she had backache intermittently, especially during pregnancy. She had also been under investigation in 1958 for headaches and disturbances of consciousness, at which time she had been found to have bilateral rotatory nystagmus and diminished sensation to pin-prick on the left half of the body. Investigations had been inconclusive and there had been no change in the physical findings since that time. Her previous pregnancy, in 1958, was normal with the delivery of a live male infant, weighing 6½ lb. (2,950 g.).

Her last menstrual period began on May 10, 1959, and she was admitted to hospital at 16 weeks with vaginal bleeding. No abnormal findings were recorded and she was discharged nine days later. Pregnancy continued uneventfully until December 17, when she was admitted to hospital with a painless ante-partum haemorrhage. Apart from the vaginal bleeding, no abnormal findings were recorded and expectant treatment was adopted. Despite rest and sedation, the slight vaginal loss continued and she went into labour, delivering herself of a living premature female infant, weight 3 lb. (1,360 g.), seven days after admission. Examination of the placenta revealed no abnormality.

On the fifth day after birth the infant developed severe cyanotic attacks and died, necropsy showing minimal intracranial haemorrhage. Death was recorded as due to prematurity.

The patient had an uneventful puerperium and was discharged from hospital eight days after delivery, only to be readmitted 12 days later with a history of lower abdominal pain and heavy vaginal blood loss for six days. On examination she was very pale (haemoglobin 45%), the blood-pressure was 100/50 mm. Hg, and pulse rate of 96 a minute. There was lower abdominal tenderness, and vaginal examination revealed the cervix open, with blood-clot protruding.

After preliminary transfusion, evacuation of the uterus was carried out; profuse bleeding occurred at this time, necessitating packing the uterus. The pack was removed after 24 hours and the bleeding did not recur.

48 hours after operation she complained of chest pain and brought up blood-stained sputum. She had a low pyrexia, and a chest x-ray examination revealed diffuse shadowing in both lung fields, of uncertain origin. *Direct questioning of the patient at this time elicited the information that she had had occasional small haemoptyses since the 20th week of pregnancy.* The temperature was fluctuating between 99 and 101° F. (37.2 and 38.3° C.), and a further chest film taken two days later revealed scattered opacities in both upper lobes. The question of tuberculosis infection was raised, sputa were sent for examination, and P.A.S. and isoniazid were given. No tubercle bacilli were identified, and by the 11th day after evacuation of the uterus the temperature had risen to 104° F. (40° C.).

The uterus was palpable in the abdomen, with bilateral adnexal masses present on pelvic examination. The report on the curettings obtained at the time of evacuation of the uterus stated: "The specimen consisted of blood clot and trophoblast. The nuclei are irregular, but could be within normal limits. It is suggested that a repeat curetting be performed if necessary and that a male-toad test be also performed for chorionic gonadotrophin."

Three days later, with a persistently high temperature, vaginal examination revealed a dark-purple nodule present on the lower anterior vaginal wall; the male-toad test was positive and a diagnosis of chorioncarcinoma was made. After preliminary blood transfusion, laparotomy was

performed on January 28, 1960, five weeks after delivery and 16 days after readmission.

At operation the uterus was found to be infiltrated by growth which extended laterally into the broad ligaments and anteriorly into the peritoneum of the utero-vesical reflection. Multiple metastases were present in the liver. Total hysterectomy and bilateral salpingo-oophorectomy was performed, operation incisions being carried out through involved tissue. The post-operative course was

FIG. 1.—Photograph of the uterus and adnexae showing widespread invasion of the uterus by chorioncarcinoma.

followed by a fall in the temperature and pulse rate, although the patient remained very ill. In view of the widespread metastases, she was transferred for chemotherapy seven days after operation.

Section report on the uterus stated: "The specimen consisted of uterus, tubes, and ovaries. The organ is 11 cm. long, 8 cm. wide, and 5 cm. thick. Several haemorrhagic tumours are present in the wall of the uterus and cervix, section showing the presence of a chorioncarcinoma invading the muscular wall of the uterus" (Fig. 1).

Condition of Patient on Transfer

She complained of pain down the right thigh and on the outer side of the right leg, pain in the lower abdomen, shortness of breath, and tiredness. She appeared pale and ill, and was coughing up much blood-stained sputum. Chest examination confirmed the marked thoracic scoliosis, but no adventitious sounds were heard. Abdominal examination revealed tenderness under the right costal margin, and pelvic examination confirmed the presence of a metastatic nodule on the lower anterior vaginal wall. Bimanual examination was limited on account of the recent operation wound, but a mass was present to the right, bulging into the vagina.

Investigations.—Chest x-ray examination: "There are a number of soft indefinite patches of consolidation in both lungs; the appearances are consistent with metastases of chorioncarcinoma." Sputum: "One large atypical hyperchromatic cell of malignant type is present; this is similar to cells seen in section of the uterus." Vaginal smear: "One or two clumps of cells are present, the appearances of which are compatible with chorioncarcinoma." Blood: These results, together with succeeding estimations, are shown in Fig. 2. Marrow biopsy: This revealed normal cellularity of the marrow.

Subsequent Treatment and Progress

Three days after admission the patient appeared worse, with high pyrexia and tachycardia. Methotrexate and 6-mercaptopurine were started on February 8, 100 mg. of the former and 1,500 mg. of the latter being given over a five-day period. There was the usual systemic toxic response at the end of the course, but a dramatic improvement occurred in the patient's condition, the temperature settling to normal, the pulse rate falling, and the blood-stained sputum disappearing. A further chest x-ray examination on February 17 showed some nodularity of the lungs, but the films failed to show the opacities previously present, particularly those of the left lung field.

Despite the general improvement in condition, the patient continued to complain of severe pain in the right leg; neurological opinion confirmed an L5-S1 lesion on the right side, probably due to secondary involvement of roots or lumbosacral plexus. Further chemotherapy was instituted, 100 mg. of methotrexate and 500 mg. of 6-mercaptopurine being given over a period of five days. Agranulocytosis followed, the neutrophils falling to 7% of a total white count of 700 c.mm. Treatment with penicillin was begun and barrier nursing instituted, the white cell count gradually returning to normal. It was not found necessary to give folic acid, but the 6-mercaptopurine was thereafter omitted.

A further course of 100 mg. of methotrexate was given over five days, beginning on March 14, when she had increasing pain in both legs, for which very large doses of narcotics were required. It was thought that the pain was due to an extension of the growth around or along roots into the spinal canal, and phenol or alcohol block was considered in an attempt to relieve the pain. The patient, however, became more ill and was unfit for transfer to the neurosurgical unit; within seven days the pain lessened, probably as a result of the destruction of the affected fibres.

Vaginal examination in early April showed that the metastasis in the lower anterior vaginal wall had disappeared although a considerable mass was still present on the right side of the pelvis. After discussion, deep x-ray therapy was begun on April 6 and completed on May 18, the dosage being: anterior oblique right pelvis, 3,100 r; anterior oblique left pelvis, 3,100 r; posterior oblique right pelvis, 2,950 r; posterior oblique left pelvis, 2,950 r; and lumbar spine, 3,000 r.

During the course of deep x-ray therapy there was a recurrence of severe pain in the legs and haemoptyses, and

the male-toad test again became positive in high dilution. A further chest x-ray film on April 25 showed some vague patches in the right upper lobe, which were regarded as further metastases. Despite the danger of marrow depression, it was decided to give further methotrexate, and 100 mg. was administered over a four-day period. After this course of treatment, together with the deep x-ray therapy to the pelvis, there was a slow subsidence of the leg pain and the haemoptyses ceased. Despite neurological involvement, the patient was able to walk and was therefore discharged from hospital on May 20 for out-patient surveillance—methandrostenolone, 10 mg. daily, being given in an attempt to increase the patient's weight.

Methotrexate, 2.5 mg. daily, was begun on May 26 on account of a continuing positive male toad test, but was discontinued after three weeks because of anaemia and a falling platelet count. As the anaemia was of severe degree (haemoglobin 42%), the patient was admitted for blood transfusion. At this time there was no evidence of residual disease in the pelvis, a chest x-ray film was clear, and the male toad test was weakly positive on undiluted urine. After transfusion the patient was discharged home. She remained well, but methotrexate, 100 mg. over a four-day period, was recommenced on July 16 on account of a dilution positive male toad test. There was the usual systemic toxic response, and the patient was admitted to hospital for five days. She was again sent home, and a further course of methotrexate was begun on August 26 on account of a continuing positive male toad test, 100 mg. again being given over a five-day period. The patient remains under surveillance.

Discussion

The active form of folic acid is folinic acid, a coenzyme in nucleic acid synthesis, so that serious upset in the latter will follow the administration of any drug which interferes with the normal utilization of folic acid in nucleic acid metabolism and cytopoiesis. In common with other antimetabolites, methotrexate has no specificity of action when used in the treatment of chorioncarcinoma, and therapy will be associated with systemic toxic effects, which have been detailed elsewhere (Hertz *et al.*, 1958; *Brit. med. J.*, 1960).

Combined therapy with 6-mercaptopurine, an anti-metabolite of adenine and hypoxanthine, has been employed in treatment (Li *et al.*, 1958; Buckle, 1959; Bagshawe and Brooks, 1959; Bagshawe and McDonald, 1960), but severe marrow depression may follow the use of 6-mercaptopurine, and there is no certain evidence that the combination of drugs is better than methotrexate alone.

Other forms of therapy have been employed, including nitrogen mustard (Anderson *et al.*, 1954) and 6-diazo-5-oxo-1-norleucine (D.O.N.) (Li, 1959). Neither of these drugs produced remission in those cases of Hertz *et al.* (1958) which were failing to respond to methotrexate. Nevertheless, the use of D.O.N. was reported upon by Li (1959) as being of value in so far as the drug might be considered as an alternative to

FIG. 2.—Treatment and progress of case.

methotrexate, for the former does not rely entirely on renal or hepatic function for excretion or detoxification.

It is of interest that the diazo-amino acids D.O.N. and azaserine (*o*-diazo-acetyl-L-serine), which were discovered during the search for anti-tumour antibiotics, were found to inhibit growth and induce abnormalities in embryonic development, producing characteristic syndromes of abnormalities in the developing chick embryo when injected into the yolk sac (Dagg *et al.*, 1956). It was further observed that resorption of fetuses of pregnant rats occurred if azaserine was administered (Murphy and Karnofsky, 1956). Interference with embryonic development appears to be a general property of the diazo-amino acids, and it is disappointing that the use of these substances in the treatment of chorioncarcinoma has not been as promising as that of methotrexate.

While it is reasonable to expect response from chemotherapy in the treatment of scattered small metastases, blood-borne agents are less likely to produce an effect against large tumour masses. This appears to be borne out by published cases, where, after hysterectomy and bilateral salpingo-oophorectomy, satisfactory results have followed the use of cytotoxic drugs in the treatment of small isolated or multiple secondary deposits. It must also be remembered that deep x-ray therapy is also extremely effective in the treatment of metastases, and this fact must be borne in mind in attempting to assess the results of treatment by chemotherapy, especially in a disease such as chorioncarcinoma, which may run a variable course, with or without treatment. In advanced cases chemotherapy should form part of an integrated scheme of treatment in conjunction with surgery and deep x-ray therapy, as in the case detailed above.

Summary

A case of widespread chorioncarcinoma is described. The diagnosis was made subsequent to a premature delivery, although its onset may have been as early as the middle trimester of pregnancy. Surgical treatment was followed by chemotherapy and deep x-ray therapy, the patient continuing under out-patient surveillance.

I thank Lady Burt for referring the patient, and the staff of all involved departments at King's College Hospital for their interest and continuing assistance in the management of the patient.

REFERENCES

- Anderson, H. E., Bisgard, J. D., and Greene, A. M. (1954). *A.M.A. Arch. Surg.*, **68**, 829.
 Bagshawe, K. D., and Brooks, W. D. W. (1959). *Lancet*, **1**, 653.
 — and McDonald, J. M. (1960). *Brit. med. J.*, **2**, 426.
Brit. med. J., **1960**, **1**, 1718.
 Buckle, A. E. R. (1959). *Brit. med. J.*, **2**, 1210.
 Dagg, C. P., Karnofsky, D. A., Lacon, C., and Roddy, J. (1956). *Proc. Amer. Ass. Cancer Res.*, **2**, 101.
 Doniach, I., Crookston, J. H., and Cope, T. I. (1958). *J. Obstet. Gynaec. Brit. Emp.*, **65**, 553.
 Hertz, R. (1948a). *Science*, **107**, 300.
 — (1948b). *Proc. Soc. exp. Biol. (N.Y.)*, **67**, 113.
 —, Bergenstal, D. M., Lipsett, M. B., Price, E. B., and Hilbish, T. F. (1958). *J. Amer. med. Ass.*, **168**, 845.
 —, —, — (1959). *Ann. N.Y. Acad. Sci.*, **80**, 262.
 — and Sebrell, W. H. (1944). *Science*, **100**, 293.
 — and Tullner, W. W. (1949). *Endocrinology*, **44**, 278.
 Li, M. C. (1959). *Ann. N.Y. Acad. Sci.*, **80**, 280.
 —, Hertz, R., and Bergenstal, D. M. (1958). *New Engl. J. Med.*, **259**, 66.
 — and Spencer, D. B. (1956). *Proc. Soc. exp. Biol. (N.Y.)*, **93**, 361.
 Murphy, M. L., and Karnofsky, D. (1956). *Cancer*, **9**, 955.
 Thiersch, J. B. (1952). *Amer. J. Obstet. Gynec.*, **63**, 1298.

FURTHER OBSERVATIONS ON THE PERSISTENCE OF ANTIBODY IN ADOLESCENTS AFTER A THIRD DOSE OF KILLED POLIOMYELITIS VACCINE

BY

E. J. C. KENDALL, M.B., M.R.C.P.

Medical Officer, Epsom College, Epsom, Surrey

ANN MILLER, B.Sc.

AND

J. O'H. TOBIN, B.M., B.Ch., Dip.Bact.

Biological Standards Control Laboratory, Medical Research Council Laboratories, London

In a recent publication (Kendall *et al.*, 1960) it was shown that 18 to 24 months after a third, booster, dose of poliomyelitis vaccine the antibody titre had declined to about a fifth of the immediate post-inoculation level. It was pointed out, however, that the results obtained gave no indication of the rate of subsequent antibody decline. The observations presented here may help in clarifying this point.

Methods

Thirty public-school boys, aged 15 to 18 years, who were initially "triple negative," took part in the study. They had been given two doses of British poliomyelitis vaccine four weeks apart, followed by a third dose 6 to 12 months later, and all had vaccine-induced antibody to each poliovirus type when bled 18 to 24 months after this third dose. More blood was obtained nine months later, and the two serum samples taken 18 to 24 months and 27 to 33 months after the third dose were titrated in parallel for poliomyelitis-neutralizing antibody by the galactose colour test, using four tubes per threefold serum dilution and 30–100 TCID₅₀ of the three types of poliovirus. The virus strains used were Brunenders (type 1), MEF-1 (type 2), and Saukett (type 3). Seven subjects from the same group of public-school boys, who had no antibody when bled at 18 to 24 months and were given a fourth dose of vaccine at that time, were also bled again and their serum was tested.

Results

A comparison between the antibody levels obtained in each of the 30 subjects is given in Figs. 1 to 3. The fall in the geometric mean titres between 18 to 24 and 27 to 33 months after the third dose was 22% for type 1, 23% for type 2, and 15% for type 3. Only one boy had no detectable type 1 antibody at the time of this bleed, and his titres were excluded from the figure for this type. The difference in titre for each of the seven boys given a fourth dose is noted in the accompanying Table, and in their case the fall in mean titre was 72% for type 1, 41% for type 2, and 40% for type 3.

Discussion

The decline in poliomyelitis antibody titre found between 18 to 24 and 27 to 33 months after a third, booster, dose of vaccine was slower than that found during the first 18 to 24 months. The rate of decline had dropped from 50% per year to under 30%. This finding agrees with those of Salk (1958) and Brown (1960), who found that there was a decrease in the rate of antibody decline in children after the first one or two years of receiving a third dose.