past decade. The countries of origin of these patients, their manner of presentation, and some diagnostic difficulties are described.

At some time while these patients were under observation in this country, acid-fast bacilli were found in the skin of at least one-third, at which stage they were potentially infective; but fortunately the resistance of the adult population to the disease is high. Intimate contact is necessary before a susceptible person can acquire the disease, although such contact need only be brief. The public health aspect of this disease in some respects is similar, on an extremely limited scale, to that of pulmonary tuberculosis, in which only a proportion of the 345,000 cases at present on the chest-clinic registers in England and Wales are "open"—that is, potentially infective.

Although some cases are self-healing, others progress to more serious and permanent complications. An awareness that the disease exists and has increased in this country will lead to earlier diagnosis. The following quotation from the Royal College of Physicians Report (1867) is of historical interest and applies in some measure to the situation in the present century.

"A few rare cases of indigenous origin have been met with in the British Isles during the present century; but by far the majority of the examples of disease which have been recognized in this country have occurred in persons who either have been born in one of our tropical possessions, or have been resident there. Such instances appear to be less infrequent than is generally imagined; and it also seems not improbable that some anomalous forms of intractable skin disease may be vestiges or obscure manifestations of a partially leprous diathesis."

I thank the physicians to St. John's Hospital for Diseases of the Skin, London, for permission to publish their cases; Dr. R. G. Cochrane, honorary consultant leprologist to St. John's Hospital for Diseases of the Skin, for his helpful advice and criticism; and Dr. Maurice Lee, of the Homes of St. Giles, Chelmsford, and Dr. W. J. Jopling, of the Hospital for Tropical Diseases, London, for further information concerning these cases.

REFERENCES

- Benson, H. (1877). *Dublin J. med. Sci.*, **63**, 562.
 Canizares, O., and Combes, F. C. (1952). *N.Y. St. J. Med.*, **52**, 2123.
 Cochrane, R. G. (1952). *Publ. Hlth. (Lond.)*, **65**, 197.
 — (1959). *Leprosy in Theory and Practice*. Wright, Bristol.
 — (1960). *Med. Press*, **243**, 77.
 Doull, J. A., Guinto, R. S., Rodriguez, J. N., and Bancroft, H. (1945). *Amer. J. trop. Med.*, **25**, 435.
 Haber, H. (1944). *Lancet*, **1**, 629.
 Macleod, J. M. H. (1925). *Brit. med. J.*, **1**, 107.
 Muir, E. (1938). *Leprosy*, 6th ed. Indian Council, B.E.L.R.A., Delhi and Simla.
 Newman, I. (1895). *Prize Essays on Leprosy*. New Sydenham Society, London.
 Royal College of Physicians of London (1867). Report on Leprosy.

The All-India Institute of Medical Sciences has now received the donation of books and journals made to it by the late Professor J. F. Fulton from his personal collection. The Rockefeller Foundation financed the transport of the consignment, which included 451 books—mostly physiological and neurological works, a complete set of the *Journal of Neurophysiology*, and a number of volumes of other journals. A special bookplate was sent with the books, and when the library of the institute is completed it is hoped that a separate section will house the collection and be named after Professor Fulton.

STUDIES IN PORTAL HYPERTENSION

FACTORS AFFECTING OPERATION MORTALITY

BY

A. G. RIDDELL, M.B.E., M.S., F.R.C.S.

AND

W. K. JONES, F.F.A.R.C.S.

*From the Departments of Surgery and Anaesthesia,
University of Manchester*

The evaluation of shunt surgery for portal hypertension awaits a more definite appraisal of the natural history of the disease. One frequent criticism is that operation can be offered only to patients with good liver function who would survive for a long period of time without operative treatment. Should the mortality from the operation be high, a group of such patients might demonstrate a poorer survival time than a group treated conservatively. Nachlas (1958) suggests that the operative mortality may be greater than the risk of death from haemorrhage. He records an operative mortality of 18% among 440 cirrhotic patients collected from the literature between 1948 and 1956.

It is our belief that the operation is of value, but its application will be extremely limited unless the operative mortality is low. We report our series of 33 patients in whom a portacaval or splenorenal shunt operation was carried out. The operative mortality was zero. Comment is made on the factors that will keep the mortality at a low level. This series is a consecutive one and includes our total experience of the operation.

Clinical Findings.—As operative mortality may be defined in a variety of ways it is important at the outset to give our own definition. We believe that a patient who dies before leaving the hospital after an operation must be counted as an operative death. No patient in this series was discharged before his wound was soundly healed or before he was able to walk and was well enough to go to his own home.

The data relevant to the patients in this series are given in the Table. In all except one of the patients the indication for operation was haemorrhage from the varices which had required transfusion for its correction. In the one exception the indication was the size of the spleen. As this patient had portal hypertension it was decided to perform a splenorenal anastomosis.

Criteria of Operability.—In this we have been guided by some of the criteria laid down by Linton (1951), particularly the following: (1) the serum albumin should be greater than 3 g./100 ml.; (2) the serum bilirubin should be below 1 mg./100 ml.; and (3) ascites should be absent. As time has gone on and we have gained more experience with the operation, these criteria have been relaxed and no patient has been rejected for operation on account of any one of these factors alone. Patients with gross ascites are not accepted unless this can be substantially reduced by medical treatment. The preoperative levels for serum albumin and bilirubin in the patients in this series are given in the Table.

Factors Contributing to a Lowering of the Operative Mortality.—The operative techniques of portacaval and splenorenal anastomosis have been described elsewhere (Riddell and Cassie, 1959) and do not differ substantially from those in use at other centres. The causes of death

to be anticipated from these operations are liver-cell failure, ammonia intoxication, and the effects of haemorrhage. We believe that the following factors rather than the operative technique contribute to the safety of the operation. These are thorough preoperative preparation, the greatest attention and care in controlling haemorrhage during operation with immediate and

In the prevention of ammonia intoxication in the immediate post-operative period the patient is taken off his high-protein diet three days before operation. Two days before operation he is given 1 g. of neomycin orally every four hours, and this is continued for a week post-operatively. This reduces bacterial fermentation in the gut and consequently lowers the production of ammonia from the alimentary tract.

Details of Cases

Case No.	Sex and Age	Diagnosis	Preoperative Level		Ascites	Type of Shunt
			Serum Alb. (g./100 ml.)	Serum Bilirubin (mg./100 ml.)		
1	M 23	Portal cirrhosis	4.0	0.5	Absent	Splenorenal
2	M 52	" "	3.1	1.2	"	"
3	M 53	" "	3.7	1.6	"	Portacaval
4	M 33	" "	3.3	0.7	"	Splenorenal
5	M 63	" "	3.5	2.0	"	"
6	F 42	" "	3.0	0.6	"	"
7	M 64	" "	3.5	0.5	"	Portacaval
8	F 59	" "	2.4	0.3	"	"
9	M 36	" "	3.3	0.6	"	"
10	M 37	" "	3.6	0.7	"	"
11	F 57	" "	3.2	0.4	"	"
12	F 42	" "	2.9	0.3	"	"
13	F 42	" "	2.8	0.7	"	"
14	M 49	" "	3.0	0.4	"	"
15	F 68	" "	2.4	0.4	"	"
16	F 71	" "	3.8	0.9	Present	"
17	M 57	" "	3.7	1.3	Absent	"
18	F 53	" "	4.2	0.6	"	"
19	F 58	" "	3.7	0.2	"	Splenorenal
20	F 46	" "	3.6	1.2	"	Portacaval
21	F 67	" "	3.1	0.5	"	Splenorenal
22	F 58	" "	3.1	0.5	"	Portacaval
23	F 59	" "	3.5	0.4	"	"
24	F 56	" "	3.4	0.8	Present	Splenorenal
25	F 64	" "	2.4	2.4	"	Portacaval
26	F 57	" "	3.5	0.8	Absent	"
27	M 40	" "	3.7	0.2	"	"
28	F 63	" "	3.1	0.9	Present	"
29	F 53	" "	3.4	0.4	Absent	"
30	F 54	" "	2.7	2.5	Present	Splenorenal
31	M 40	" "	4.3	0.2	Absent	Portacaval
32	M 46	" "	4.5	0.4	"	"
33	F 74	Hydatid disease. Budd-Chiari syndrome	2.8	1.6	Present	Splenorenal

quantitative replacement of blood loss, planned anaesthesia, and careful post-operative care. The more theoretical aspects of the surgical care of patients with liver disease have been reported (Riddell and Jones, 1959), and although it is not our intention to review this again we describe our present programme for the care of these patients.

Pre-operative Care

The first essential of preoperative care is to bring liver function to the best achievable standard, and the main therapeutic weapon is a high-protein, high-calorie diet. This should be started with caution because of the danger of producing ammonia intoxication from high-protein feeding (McDermott, Adams, and Riddell, 1954). If the patient can tolerate it, the diet is increased to 120 g. of protein and 3,000 calories a day. Tube feeding, especially at night, is useful in patients with anorexia. Generous vitamin supplements are given, with the addition of vitamin K intramuscularly. In a few patients the level of the serum albumin has been raised by intravenous infusion of salt-free albumin with apparently beneficial results.

Haemoglobin deficits should be corrected before operation by transfusion with packed red cells. However, care is required as substantial increases in blood volume may raise the portal pressure and precipitate further haemorrhage. Attempts to increase the haemoglobin level above 10 g./100 ml. (70%) are inadvisable.

Care During Operation

Careful blood replacement is of the utmost importance during operations on patients with cirrhosis, for two reasons: firstly, oligæmia will result in hypoxia of the liver cells which may precipitate liver-cell necrosis; and, secondly, patients with liver damage are more liable to citrate intoxication than the normal patient, as the damaged liver is less able to deal with the citrate in the transfused blood. Quantitative replacement of any blood loss is therefore required. To achieve this the blood loss is continuously estimated by the technique of swab weighing and any deficit is immediately replaced.

Anaesthetic Technique

These patients tolerate hypotension and its attendant hypoxia badly, and every precaution should be taken to prevent it occurring, especially in the immediate post-operative period. It is therefore mandatory not to return these patients to the ward until any blood loss is replaced and the blood-pressure is restored to its preoperative level. These factors have become the guiding principles in the development of the anaesthetic technique.

In our early cases hypotensive anaesthesia was not employed to reduce bleeding, but in the second half of the series halothane has been used with increasing frequency to lower the systolic blood-pressure to between 80 and 100 mm. Hg. It is very difficult to define the degree of hypotension that is permissible and for how long it may be continued. At present we feel that the levels mentioned above are satisfactory, but the arterial pressure must be returned to its normal value before the end of the operation. Hypotensive anaesthesia, together with the employment of a positive/negative-phase respiratory pump, has reduced the bleeding from the operative field and the total blood loss has been approximately halved. Halothane produces marked vasodilatation, and with blood-pressures of the order mentioned above there is an efficient peripheral circulation. Although in some instances the administration of halothane has continued for four or five hours no adverse effect on liver function has been detected. By using halothane as an adjunct to nitrous oxide and oxygen it is possible to reduce the induction dose of thiopentone to the minimum, and the morphine group of drugs are unnecessary until the post-operative period.

Complete muscular relaxation is essential throughout the operation, and for this both suxamethonium chloride and D-tubocurarine have been used, but never both drugs in the same patient. Better relaxation is obtained with suxamethonium given intermittently, but the technique is tedious and the patient requires careful observation both to maintain full relaxation and to prevent an excess of the drug being given as the interval between successive doses increases. There has been no difficulty in establishing normal respiratory exchange at the end of operation. Neostigmine is always given after

D-tubocurarine in dosage sufficient to produce a normal tidal air. Whichever drug is used to produce relaxation it is essential to keep the patient on the respiratory pump until it is absolutely certain that the action of the relaxant has been completely reversed.

The use of a mechanical ventilator for long operations of this kind offers three main advantages: (1) it ensures even ventilation, which is necessary to maintain proper oxygenation and carbon dioxide elimination; (2) the regular rhythm of the pump is a great help when performing a difficult dissection in the hilum of the liver; and (3) the use of a negative-phase respirator reduces venous congestion and so bleeding.

The use of halothane with a respiratory pump is permissible only if the concentration of vapour in the inspiratory gases is accurately known and controllable. Probably the most economical technique is to pass 3-4 litres of 1% halothane per minute, using nitrous oxide and oxygen as a vehicle, into a non-return circuit and to make up the balance of the minute-volume with air. If the whole of the inspired gases are passed through a halothane vaporizer then concentrations of less than 0.5% must be used. Anaesthesia must be kept light enough for the patient to regain consciousness before he returns to the ward.

We make it a rule never to return a patient to the ward unless: (a) blood loss has been completely replaced; (b) blood-pressure is at its preoperative level; and (c) reflexes are present with a normal respiratory function—that is, respiratory exchange is sufficient to maintain full oxygenation without the aid of oxygen.

Post-operative Care

The immediate post-operative care of the patient is similar to that of any patient who has had a thoraco-abdominal incision. These patients tolerate all forms of sepsis badly, and it is important to get the lung expanded at the earliest possible moment and remove any retained secretions.

In patients with liver disease there are abnormalities of water and salt metabolism, and it is easy to produce water and sodium retention. It is our practice to give no sodium intravenously on the first post-operative day and to limit the intake of water to 1,000 ml. for this period. The gastric tube is removed on the second day, and shortly after this the intravenous infusion is discontinued. In a few patients 500 ml. of 0.9% NaCl has been given on the second day.

After surgery, protein feeding is resumed with caution during the fifth to seventh post-operative days. The diet is then built up in 10-g. increments every two days until 70 g. is reached. The protein intake is maintained at this level for three months before increasing it to 120 g. Increasing the dietary protein above 70 g. too soon after operation may precipitate portasystemic encephalopathy.

Summary and Conclusions

For the operation of portacaval and splenorenal anastomosis to be of value in the treatment of bleeding oesophageal varices the operative mortality must be low.

We report a series of 33 consecutive shunt operations without operative mortality.

The factors which are important in maintaining a low operative mortality are described.

REFERENCES

- Linton, R. R. (1951). *Ann. Surg.*, **134**, 433.
 McDermott, W. V., Adams, R. D., and Riddell, A. G. (1954). *Ibid.*, **140**, 539.
 Nachlas, M. M. (1958). *Ibid.*, **148**, 169.
 Riddell, A. G., and Cassie, A. B. (1959). *Recent Advances in Surgery*, edited by S. Taylor, p. 437. Churchill, London.
 — and Jones, W. K. (1959). *Ibid.*, p. 420.

ASPECTS OF CONVALESCENCE AFTER HERNIORRHAPHY

A REVIEW OF 762 PATIENTS

BY

R. J. JARRETT,* M.B., B.Chir.

Late House-Surgeon, Bolingbroke Hospital, London

AND

J.-C. GAZET,† F.R.C.S.

Late Surgical Registrar, Bolingbroke Hospital, London

The purpose of this investigation was to study the actual convalescence of patients treated at a general surgical hospital and to relate this to other factors affecting the results of operation. All operations for herniae performed at the Bolingbroke Hospital in the years 1948-57 have therefore been reviewed.

"Nobody seems to be able, or prepared, to answer the question: When shall a patient return to work? Perhaps this is because the convalescent is in a therapeutic no-man's-land." And, again, convalescence "is an aspect of hospital treatment which after 20 years remains almost unchanged" (*Lancet*, 1957).

The Oxford Dictionary defines convalescence as the period of regaining health, and one can therefore regard surgical convalescence as beginning as the operation ends. The concept of health is incapable of mensuration. So in discussing the problems of convalescence it is more profitable to consider two periods of time: firstly, the length of stay in hospital and/or convalescent home; and, secondly, the length of time between discharge and return to work or full activity.

In practice these periods are often a compromise between what the patient considers economically possible and what the physician considers ideally necessary. Although the former is beyond the scope of the present study, its importance is reflected in the results obtained in this series.

The length of convalescence considered desirable after individual operation is a subject which has received little study. Medical students are taught much about the criteria for operation, the methods of operation, and to a less extent about post-operative management. But authoritative guidance on the further stages is lacking. Each surgeon has his own views on how long he likes to keep his patients in bed post-operatively and whether or not they should spend a period of convalescence at a recognized home or hostel. But the time of return to work may be decided upon by the registrar or house-surgeon who sees the patient in the follow-up clinic, by the patient's own general practitioner, or even by the patient himself.

*Present address: Department of Pharmacology, Guy's Hospital Medical School, London.

†Present address: Guy's Hospital, London.