prostatic cavity is completely epithelized. Cystoscopy before this time shows deposits of fibrin, surface sloughs, and pieces of unabsorbed catgut in the area of the operation. It is surprising how these appearances are generally associated with the passage of a good, well-controlled stream of clear urine.

Summary and Conclusions

A series of 120 consecutive prostatectomies performed in two small non-teaching hospitals is described. Ogier Ward's vesico-capsular operation has been used. There was no undue selection of cases, and the results are comparable with those which have prevailed in large centres for many years.

Cardiovascular accidents are the most serious menace after prostatectomy, and this is probably the reason why the mortality rate rises steadily with the age of the patients. Acute retention of urine also raises the mortality rate, the reason probably being that the retention is often precipitated by some cardiovascular or other complaint. Where acute retention has occurred preliminary bladder drainage with an indwelling urethral catheter is maintained until the patient is fit for operation. If necessary, it may be continued for many months.

In the absence of obvious systemic disease the patient's mental state is the most important single factor in the assessment of the operative risk. Old men with senile cerebral degeneration are very poor risks, while the patient whose mind is clear will probably survive prostatectomy whatever his age.

About 90% of patients who need prostatectomy can be operated upon with safety. There is a hard core of a little under 10% (represented in this series by a preoperative rejection rate of 3% and a mortality rate of 4%) which constitutes a very serious problem not yet satisfactorily solved.

Low spinal or epidural anaesthesia is thought to be the best form for the bad risk.

The vesico-capsular operation is described.

Post-operative management is described. It is based on a closed bladder-irrigation system. Every effort should be made to avoid cross-infection, which occurs very easily.

Post-operative urinary infection has been reduced to a minimum, but wound infection was not uncommon and has been the most troublesome complication in the series.

The prostatic cavity is not healed until about eight weeks after the operation, but good function has usually been achieved well before this time.

REFERENCES

- Bourque, J.-P. (1954). *J. Urol.*, 72, 918.
 Dawson-Edwards, P. (1956). *Lancet*, 2, 332.
 Ellis, H. and Leatherdale, R. A. L. (1958). *Ibid.*, 2, 1189.
 Freyer, P. J. (1920). *Clinical Lectures on Enlargement of the Prostate*. Baillière, London.
 Galbraith, W. W. (1948). *Proc. roy. Soc. Med.*, 41, 73.
 Harris, S. H. (1914). *Brit. J. Surg.*, 21, 434.
 Hey, W. H. (1945). *Ibid.*, 33, 41.
 Jacobs, A. (1951). *Lancet*, 1, 1088.
Lancet, 1956, 2, 343.
 Millin, T., Macalister, C. L. O., and Kelly, P. M. (1949). *Lancet*, 1, 381.
 Riches, E. W. (1949). *Proc. urol. Soc. Aust.*, No. 3, 39.
 — (1954). *Ann. roy. Coll. Surg. Engl.*, 15, 128.
 Stewart, H. H. (1945). *Brit. med. J.*, 2, 724.
 Ward, R. Ogier (1948). *Lancet*, 1, 472.
 Watts, G. T. (1956). *Brit. med. J.*, 2, 1283.
 Wells, C. (1953). *Modern Trends in Urology*, edited by E. W. Riches. Butterworth, London.

PANCREATIC LITHIASIS A FOLLOW-UP STUDY OF 17 CASES*

BY

R. MAHADEVAN, M.S., F.R.C.S., F.R.C.S.Ed.

Formerly Professor of Surgery, Madras Medical College;
 Surgeon, General Hospital, Munnar, South India

A follow-up study of 17 cases of pancreatic lithiasis showed that the calculi gradually disappeared after operations mainly directed towards relief of the severe pain characteristic of this condition. In some cases the pancreatic involvement was total, and it was thought that total pancreatotomy would ultimately become inevitable. Yet the hundreds of calcified areas in calcareous pancreatitis and the numerous calculi in the ducts gradually but almost completely disappeared.

Pancreatic lithiasis, while not a common condition, is commoner than is generally thought. Many cases are missed because a plain radiograph is not taken routinely in cases of recurrent abdominal pain.

FIG. 1.—Case 4. Barium-meal radiograph of patient with recurrent attacks of severe colicky abdominal pain.

FIG. 2.—Case 4. Plain radiograph showing almost entire calcification of pancreas with myriads of calcified areas. These were almost completely hidden by the barium.

Barium-meal series alone may hide even large and multiple calculi (Figs. 1 and 2), or the calculus shadows may pass unrecognized in the confusion caused by the barium shadows. Eliason and Welty (1948), by subjecting pancreatic glands removed at necropsy to radiographic studies, showed that the incidence of calcification was 1 in 20. Calcification occurs in approximately half the cases of chronic pancreatitis (Allbritten and Pressly, 1950).

*An Arasthu Memorial Lecture delivered at Hyderabad on August 30, 1959.

Of the 17 cases studied, 14 were calcareous pancreatitis, and in three the stones were in the duct. Sixteen were in males. The main symptom was attacks of severe upper abdominal pain, radiating to the back and sometimes to either shoulder or scapula. Two patients threatened suicide because of unbearable pain. Some became drug addicts. Many were emaciated and anaemic at the time of admission. Some had mild icterus. None was associated with cholecystitis or gall-stones. Sometimes calcareous pancreatitis may be an accidental finding, and the patient may have no symptoms referable to it. Further, as much as 90% of the gland may be involved without giving rise to symptoms if the body and tail only are affected, while a small lesion near the head obstructing the flow of secretion may cause early and severe symptoms (Allbritten and Pressly, 1950).

Pancreatic calculi near the head or tail may resemble renal stones, and those near the head may mimic gall-stones. A point worth noting, to which, so far as I am aware, attention has not hitherto been drawn, is that in the lateral views of straight radiographs shadows of pancreatic calculi lie just in front of those of vertebral bodies, a fact not so well known as the fact that the shadows of renal stones overlap shadows of vertebral bodies, and those of gall-stones lie just behind the anterior abdominal wall.

Palpation may reveal a mass due to a coexisting cyst, chronic pancreatitis, or growth. In this series five patients had associated pancreatic cysts.

Treatment

Medical Treatment of Acute Attack.—The analgesic most often recommended is pethidine. Morphine is said to cause spasm of the sphincter of Oddi, but sometimes, when pethidine and other drugs failed, it gave the required relief (Case 4). Other drugs used are atropine and ephedrine to reduce pancreatic secretion and ganglion-blocking drugs (said to act by reducing vagal secretion). Sympathetic block or continuous epidural block may give great relief, sometimes for even days and weeks (Cases 3 and 4). Attention to nutrition and control of associated diabetes, if present, are important. Alcohol should be avoided, as it may precipitate an attack.

Surgical Treatment When Calculi Are in Ducts.—The duct is incised and the calculi are removed. If there is associated gross infection the duct is drained by direct external drainage or through the bile passages (T-tube drainage). Calculi impacted in the ampulla require removal by the transduodenal route. In Case 6 several calculi were found not only in the main duct but also in its various ramifications. Some could be removed by incising the duct; others could not be removed, but prolonged external drainage was maintained for nine and a half months. The drainage tubes were removed when the attacks of pain ceased, despite several calculi being left behind. Later it was found that all the calculi except a solitary one in the head had completely disappeared (Fig. 7).

Surgical Treatment of Calcareous Pancreatitis.—This condition is more difficult to treat. Resection of the affected area may be the logical procedure, but this, especially when it involves the head or entire gland, is risky. Total resection has the serious disadvantage of permanent loss of pancreatic secretion. The effect of indirect operation should be tried first. Sphincterotomy

is not very successful, though some have claimed good results after it. It is necessary if the calculi are impacted in the ampulla.

Diversion of Biliary Flow

Diversion of the biliary flow sometimes gives excellent results. It consists of some form of biliary-intestinal anastomosis like cholecystojejunostomy or choledochojejunostomy. In Cases 1 and 2, where cholecystojejunostomy was performed for calcareous pancreatitis, the patients were completely free from symptoms 14 and 11 years after operation. But the method failed in some cases, and various additional procedures like splanchnicectomy and sympathectomy were required later. In two such cases (Nos. 3 and 4) the calculi almost completely disappeared after these procedures.

Case 1.—A man of 30 was admitted on June 21, 1945, with a history of recurrent attacks of abdominal pain, agonizing on occasions, for 18 months. A mass was palpable in the right hypochondriac and epigastric regions. He had a mild icterus. Urine showed a trace of sugar. X-ray examination: Entire pancreas calcareous; barium-meal-filling defect of pyloric area and stretched out C curve of the duodenum. Operation on July 9 showed a hard, nodular, and calcified pancreas. A large multilocular cyst in the head was pushing the duodenum forwards. The stomach and duodenum were dilated. The cyst was tapped in four different places and about 15 oz. (425 ml.) of thick, glairy fluid drawn. There were no calculi in the duct. Because of the jaundice and the likelihood of an increasing accumulation in the cyst causing bile-duct obstruction, cholecystojejunostomy was performed, utilizing a loop of jejunum 16 in. (40 cm.) below the duodeno-jejunal flexure. The pain disappeared and the patient has been comfortable for 14 years.

Case 2.—A woman of 20 was admitted on May 7, 1948, for attacks of pain in right hypochondrium, epigastrium, and back of 15 years' duration, occurring almost daily in the last two years. X-rays revealed calcareous pancreatitis. Laparotomy showed the entire pancreas to be hard and nodular; no stones in the duct. The caudal portion of the duct was wider than the cephalic. A No. 6 catheter was passed towards the head and a No. 8 towards the tail, and these were stitched in place. The incision in the duct was sutured around them and this area reinforced with omentum. The catheters were brought out through the left loin. A hard mass was felt near the ampulla. Duodenotomy showed no stones in the ampulla. Cholecystojejunostomy was performed, Morison's pouch was drained for 48 hours, and the tubes draining the pancreas were removed on the eleventh day. The patient was last seen in August, 1959, when she had been free from symptoms for 11 years.

Nerve Resection

Various forms of nerve reaction have been successfully adopted for the relief of the pain. As the nerve supply to the pancreas is still doubtful, diverse procedures have been suggested—unilateral (right or left) or bilateral thoraco-lumbar sympathectomy. The segments recommended to be resected have varied from D3 to L2 to as few as D8 to L1. Others have resected only the nerves passing through the pancreas from the coeliac ganglion. A study of the literature suggests that the best method to adopt is that if the pain is mainly on one side the operation should be performed on that side, and if bilateral on both sides. A sympathetic block beforehand may help in deciding. In one case where cholecystojejunostomy failed, left-sided nerve resection gave relief only on that side; so right-sided resection was also performed and the patient became free from pain (Case 3). The minimal operation found to provide total analgesia of the pancreas, and the one proposed as most suitable, is bilateral resection

of ganglionated chains from D 11 to L 1 and of the lowest 8 cm. of the greater splanchnic nerves. Unilateral sympathectomy alone may sometimes succeed (Ray and Console, 1949).

In two cases (Nos. 3 and 4) where cholecystojejunostomy failed to give relief, thoraco-lumbar sympathectomy succeeded. Serial radiographs in both cases showed a progressive decrease in the number and size of the calculi.

Case 3.—A boy aged 12 was admitted on October 23, 1952, with a history of recurrent attacks of colicky abdominal pain for seven years. He had had mumps at 8, measles at 9, and attacks of dysentery off and on. X-ray examination showed calcareous pancreatitis, confirmed at operation on October 31. There were no stones in the duct, which was drained by a self-retaining catheter. As an induration was felt near the head, sphincterotomy was performed. The operation was completed by cholecystojejunostomy, and the catheter removed on the sixth day. Pain recurred after a year. On October 18, 1954, resection of the left dorsilumbar sympathetic trunk and left greater and lesser splanchnic nerves was performed. *This resulted in relief of pain in the left side only. The attacks of pain on the right side persisted.* On July 25, 1955, right-sided nerve resection was performed. He is now free from pain, has put on weight; radiographs showed a progressive almost complete disappearance of calcification.

FIG. 3.—Case 4. Plain radiograph taken on August 4, 1953, showing extensive calcareous pancreatitis involving the entire pancreas.

FIG. 4.—Case 4. Radiograph taken in 1956, showing disappearance of majority of the calculi.

Case 4.—A man aged 21 was admitted on August 1, 1953, with a history of recurrent attacks of severe right hypochondriac pain for nine years. He became an addict to toddy and cannabis indica. He was very ill-nourished. During attacks he used to feel a lump in the right hypochondrium. While in hospital, morphine was the most effective pain-killing drug; pethidine was not so effective. Laparotomy showed that the entire pancreas was the seat of calcareous deposits. A cyst 5 cm. in diameter was found in the head of the pancreas. On aspiration, 4 oz. (114 ml.) of slightly bile-stained fluid was withdrawn. The cyst was then opened and drained externally, and cholecystojejunostomy performed. Pain recurred after six weeks. *Bilateral paravertebral block gave immediate relief, which lasted for a fairly long time.* However, the pain recurred early in 1956. A bilateral thoraco-lumbar and splanchnic resection, done in two stages, prevented further attacks, and his general condition has greatly improved. Serial radiographs showed disappearance of the greater part of the calcified areas (Figs. 3 and 4).

Whether absorption of the calcified areas was due to control of the infection after cholecystojejunostomy or to a revascularization of the pancreas after sympathectomy is not clear. Possibly both play a part. Such explanations cannot, however, be the entire answer to the problem, for in at least one patient nothing was done at operation apart from mobilization and inspection of the pancreas, and yet the calcified masses entirely disappeared in three months (Mohan Rau, 1956).

Vagotomy has sometimes been recommended on the principle that it lessens vagal secretion, but very few have adopted it.

Case 5.—A weaver aged 25 was admitted on September 13, 1954, with recurrent attacks of pain in the left epigastric and lumbar regions of five years' duration. A vague, diffuse intra-abdominal swelling in the epigastrium felt doughy and was immobile and dull to percussion. A radiograph showed calculi in the head of the pancreas. Barium-meal examination revealed irregular filling in the pyloric region. A cholecystogram disclosed a normal gall-bladder. Laparotomy on October 13 showed calculi in the head. There was a hemispherical cyst about 4 in. (10 cm.) in diameter in the body. Tapping drew about 100 ml. of colourless, glairy fluid. The cyst was anastomosed to the stomach. The operation was completed by cholecystojejunostomy. The coeliac ganglion was exposed and the right great splanchnic nerve resected. The left one could not be located. The patient was comfortable until November 5, when he developed epigastric pain, which he volunteered was entirely different from his previous pain. It was more like peptic-ulcer pain.

Possibly this patient had a pyloric ulcer also, of which the radiographic appearances were suggestive, but it is difficult to be sure, as such appearances are not unknown in chronic pancreatitis itself. However, he has remained comfortable with medical treatment for peptic ulcer.

Calculi Situated in the Pancreatic Duct

There were three cases of calculi in the pancreatic duct. Case 6 is one in which the calculi were occupying every nook and cranny of the duct system. Some of these could be removed by incising the duct. As the rest, dispersed in rather inaccessible parts of the duct, could not be removed, free external drainage of the duct was established. The pain soon disappeared. The drainage tubes were removed after nine and a half months, though several calculi were still present. Some months later all these, except a solitary one in the head, disappeared (see Fig. 7). The patient has been free from symptoms ever since.

Case 6.—A weaver aged 27 was admitted on October 16, 1955, for recurrent attacks of pain in the epigastric and

umbilical region of five years' duration. At operation the entire pancreas was found to be a calcareous mass, the head markedly enlarged. The duct was incised and 11 large calculi were removed. Others were in the branches. It was not easy to decide whether they were in the duct or in the glandular substance. Those in the duct slipped out easily on incision as out of a thin paper bag. External drainage was established (Figs. 5 and 6). The operation was completed by cholecystojejunostomy. As the patient had no relief the abdomen was reopened on November 30 with the idea of sphincterotomy, but the ampulla could not be located. The duct was reopened and several more calculi were removed. Some of these were dislodged by manipulation with the finger through the duodenal opening and introducing Desjardins forceps into the pancreatic duct, but attempts at passing a catheter through the ampulla failed.

FIG. 5.—Case 6. Radiograph showing two catheters draining the pancreatic duct. For further details see text.

FIG. 6.—Case 6. One of the serial radiographs showing a few calculi in the tip of the draining Malecot's catheter. For further details see text.

FIG. 7.—Case 6. Plain radiograph taken in May, 1959—four years after the original operation. Note that only a solitary calculus remains. All the other calculi have disappeared.

The duct was periodically washed. The patient discharged himself on April 24, 1956, with the drainage tubes still in place. He returned on July 7. A small amount of foul-smelling discharge was escaping from the catheter. Periodical irrigation was performed. The catheter was removed on August 8. The wound healed rapidly and he was discharged on August 23. He has been free from symptoms except for a small incisional hernia. When last seen (May 16, 1959) he was in good health. All the calculi had disappeared except for a solitary one in the head (Fig. 7).

This happy result has possibly been brought about by the following process. Through the catheters draining the pancreatic duct, periodical wash-outs were performed. Gravel used to be present in the returns. In later stages it was possible to change the self-retaining catheters very much like changing a suprapubic catheter. When these catheters were changed some small calculi used to come away, lying in their lumen or stuck in the hollow of their tips (Fig. 6). The increased vascularity of the gland after handling at the two operations has probably played a part in the absorption of some of the calculi in the glandular substance.

That infection can play a large part in the production of symptoms and calcification is borne out by the following case.

Case 7.—A boy aged 16, admitted in February, 1957, had severe attacks of colicky abdominal pain. The clinical features and radiographic appearances suggested calcareous pancreatitis. As he had unbearable pain a splanchnic block was given, and within 15 minutes the anaesthetist found him in a state of collapse. Cardiac arrest quickly followed. Cardiac massage resulted in recovery, but he died 24 hours later. A partial post-mortem examination was made. The ampulla was pin-point. On slitting it open the pancreatic duct was found very dilated and filled with pus. Calcareous nodules were felt in the head of the pancreas.

Two other patients had calculi removed from the ducts. One developed cholaemia and died after a week. In the other an associated cyst was externally drained; a pancreatic fistula occurred and attempts at closing it failed. He died six months later. Details of this case have been given elsewhere (Mahadevan, 1947).

Two patients who were not operated upon have been symptom-free for seven and for over four years respectively, though periodical radiographs showed persistence of the calculi. One has developed diabetes but is at full work as an engineer. The other, a boy of 16, came in an emaciated condition, but is symptom-free on medical treatment and his general condition has improved.

Summary

Various operations designed to give relief to the unbearable pain in pancreatic lithiasis not only relieved the pain but brought about quite unexpectedly complete absorption of the numerous calcified areas in two cases of calcareous pancreatitis and of the several calculi situated in the duct in one case. Control of infection by free drainage of the duct and the increased vascularity after sympathetic denervation of the pancreas seemed to have brought about this happy result. Mechanical wash-out of the pancreatic duct also seems to have helped.

REFERENCES

- Allbritten, F. F., jun., and Pressly, L. C. (1950). *Surg. Clin. N. Amer.*, **30**, 1627.
- Eliason, E. L., and Welty, R. F. (1948). *Ann. Surg.*, **127**, 150.
- Mahadevan, R. (1947). *Indian J. Surg.*, **9**, 73.
- Mohan Rau, U. (1956). *Ibid.*, **18**, 471.
- Ray, B. S., and Console, A. D. (1949). *Surg. Gynec. Obstet.*, **89**, 1.