among diabetics, but since both diabetes and infections of the urinary tract occur most commonly in middle-aged and elderly women, it is not surprising that the number of diabetics with infections should seem large. The greater hazard of infections in diabetic patients serves also to make them more memorable.

Previous reports have, with one exception (Huvos and Rocha, 1959), found a higher incidence of infection in diabetics than in non-diabetics. The great majority, however, base their findings either on the results of necropsies on patients dying in hospital or on hospital in-patients. The lack of controlled studies on a diabetic out-patient population and "healthy" non-diabetics is surprising. Thus Barnard *et al.* (1953), in a series of 100 necropsies on diabetics dying in hospital, found that 30% had suffered significant infection of the urinary tract. Merritt and Sanford (1958) examined 100 consecutive ambulatory and non-ambulatory admissions to a medical ward and found a significant excess of patients with diabetes in the group with positive cultures. It is probably invalid to apply results from an investigation such as this, based on hospital admissions, to a general population, either diabetic or non-diabetic, particularly as Rengarts (1960), studying diabetic admissions to the Joslin Clinic Service, found a much higher incidence of urinary infections in patients confined to bed than among ambulatory diabetics. In this latter study the incidence of infection among ambulatory diabetics (14% of 43 patients) was very similar to the present findings.

That infections of the urinary tract are commoner in women than in men has long been recognized, and a marked excess of infected females is seen in the present series, as in others (Kass, 1955, 1956; Rengarts, 1960). The striking significance of age in determining the incidence of infection is not so widely appreciated, but where it has been taken into account the number of patients infected has risen steadily with age in both diabetics (Huvos and Rocha, 1959) and non-diabetics (Loopuyt, 1946). Kass, in an extensive study, found that 18% of 54 female diabetics and only 6% of 337 females attending as out-patients had infections of the urinary tract, but makes no mention of the ages of either group, so that it is difficult to assess the significance of his findings. Failure to take into account the large numbers of middle-aged and elderly women in the diabetic population is probably the main reason for the present confusion of thought on this subject.

Finally, although adequate figures are available regarding the incidence of infections of the urinary tract in hospital in-patients (Marple, 1941; Loopuyt, 1946), there is a striking lack of information about the general population. The need to obtain catheter specimens of urine from the female for reliable bacteriological and microscopical study has in the past been an understandable deterrent. Quantitative examination of clean-catch voided specimens now provide a safe and relatively simple method of evaluating infections of the urinary tract, and its use in "healthy" populations such as the controls in the present study could with advantage be extended further. Only in this way will the significance of asymptomatic infections be recognized, their possible association with other disorders such as hypertension be elucidated, and methods for their control be put on a more effective and adequate basis.

Summary

The use of quantitative bacterial counts on clean voided urine specimens is described.

A comparison of the incidence of infections of the urinary tract in 150 randomly selected diabetics, representative of a large diabetic clinic, with a similar number of non-diabetic "healthy" controls shows no significant difference between the two groups.

The high incidence of infection of the urinary tract in middle-aged and elderly women in both groups is noted and its significance discussed.

We wish to thank Miss M. A. Butcher for technical assistance and the staff of the casualty department of the Birmingham General Hospital for their co-operation. This work was in part supported by a grant from the Diabetic Association.

REFERENCES

- Baldwin, A. D., and Root, H. F. (1940). *New Engl. J. Med.*, **223**, 244.
 Barnard, D. M., Story, R. D., and Root, H. F. (1953). *Ibid.*, **248**, 136.
 Beeson, P. B. (1955). *Yale J. Biol. Med.*, **28**, 81.
 — (1958). *Amer. J. Med.*, **24**, 1.
 Boshell, B. R., and Sanford, J. P. (1958). *Ann. intern. Med.*, **48**, 1040.
 Guze, L. B., and Beeson, P. B. (1956). *New Engl. J. Med.*, **255**, 474.
 Huvos, A., and Rocha, H. (1959). *Ibid.*, **261**, 1213.
 Kass, E. H. (1955). *Amer. J. Med.*, **18**, 764.
 — (1956). *Trans. Ass. Amer. Phycns.*, **69**, 56.
 Loopuyt, L. (1946). *Acta med. scand.*, **125**, 245.
 Marple, C. D. (1941). *Ann. intern. Med.*, **14**, 2220.
 MacDonald, R. A., Levitin, H., Mallory, G. K., and Kass, E. H. (1957). *New Engl. J. Med.*, **256**, 915.
 Merritt, A. D., and Sanford, J. P. (1958). *J. Lab. clin. Med.*, **52**, 463.
 Monzon, O. T., Ory, E. M., Dobson, H. L., Carter, E., and Yow, E. M. (1958). *New Engl. J. Med.*, **259**, 764.
 Ophüls, W. (1926). *Stan. Univ. Publ. med. Sci.*, **1**, 127.
 O'Sullivan, D. J., Fitzgerald, M. G., Meynell, M. J., and Malins, J. M. (1960). *J. clin. Path.*, **13**, 527.
 Perla, D., and Marmorston, J. (1941). *Natural Resistance and Clinical Medicine*. Little, Brown, Boston.
 Rengarts, R. T. (1960). *Amer. J. med. Sci.*, **239**, 159.
 Robbins, S. L., and Tucker, A. W. (1944). *New Engl. J. Med.*, **231**, 865.
 Schofield, R. A., and Baker, R. D. (1956). *Arch. Path. (Chicago)*, **61**, 407.
 Sharkey, T. P., and Root, H. F. (1935). *J. Amer. med. Ass.*, **104**, 2231.

VARICOCELE: A TREATABLE CAUSE OF SUBFERTILITY

BY

L. STUART SCOTT, M.D., Ch.M., F.R.C.S.Ed.
F.R.F.P.S.

Consultant Surgeon to the Male Subfertility Clinic, Western Infirmary, Glasgow

Disappointment at the results of drug therapy in the treatment of male subfertility (Scott, 1959) has given way to increasing optimism as regards the results now being achieved by surgery. Before Tulloch (1952) first reported restoration of fertility in an azoospermic man after varicocele ligation, a varicocele was seldom if ever considered as a possible cause of subfertility. Russell (1954) later noted that the incidence of varicocele was much greater in men attending a subfertility clinic than in the husbands of women attending his antenatal clinic. Finally, I showed that subfertility could be detected in 66% of varicocele patients compared with only 39% of those with no varicocele (Scott, 1958a). Persistence of a varicocele beyond the age of 30 years resulted in progressive deterioration of fertility in the cases studied by Russell (1957), and I subsequently reported three cases of progressive subfertility over a period of five years which was finally checked and returned to acceptable fertility by varicocele ligation (Scott, 1960a).

The incidence of subfertility appears to be in direct proportion to the size of the varicocele (see Chart). In a previous communication (Scott, 1958b) I showed that the effects on the sperm count and sperm motility were also related to the size of the varicocele. A later study (Scott, 1960b) confirmed this clinical impression by demonstrating that the greatest improvement followed

operation on the largest varicoceles, ligation of small varicoceles gave the poorest results, and control ligation of the spermatic veins in subfertile volunteers with no varicocele produced no change in fertility at all.

can be attributed to the normal fluctuation of sperm count and motility, as described by MacLeod and Gold (1956). In order to refute these criticisms and overcome the possibility of error in assessing the value of this operative procedure, the seminal findings have been "ranked" in this study, and only improvement to a higher rank was considered significant.

Material and Method

Of the 108 subfertile men whose varicoceles have been ligated in this clinic, 11 have since left the district and have not been traced, while four were operated on too recently to be included in this study. Pre- and post-operative seminal studies were carried out on 82 cases with pre-operative subfertility and on six cases of pre-operative azoospermia. Another five men have declined further seminal investigation but have notified a successful pregnancy since operation.

Varicocele ligation and division at the deep inguinal ring was performed on 108 men with left-sided varicoceles which, with the patient standing upright, were greater than the diameter of the index finger.

As it had previously been shown (Scott, 1958b) that the deleterious effects of varicocele were most pronounced on the sperm count and sperm motility, "subfertility" was taken to denote a sperm count below 20 million per ml. or a sperm motility showing less than 50% of the sperms actively progressive five hours after ejaculation. Two or more seminal samples were studied

TABLE I.—Method of Ranking Sperm Counts and Motility

Rank No.	Sperm Count (millions/ml.)	Rank No.	Active Motility
1	Over 60	1	Over 50%
2	21-60	2	11-50%
3	6-20	3	Under 11%
4	Under 6		
5	Azoospermia		

before and after operation, and the figures referred to in the text represent an average of these readings. Allowance for normal fluctuation in count and motility was made by using the ranking system shown in Table I. Post-operative readings represent the seminal findings six months after ligation. All samples were produced by coitus interruptus after three days' abstinence from intercourse.

Results

Overall Effect on Fertility.—Before operation all cases were judged to be subfertile, because of a low sperm count or poor motility or both. Within six months of ligation, 67 cases (83.7%) were judged to be fertile (63 having vastly improved seminal rankings, and 4 producing a pregnancy without further analysis). Eight cases (10%) improved to higher seminal rankings but not to acceptably fertile levels. Four cases (5%) remained in their original ranking, and one case dropped to a lower rank.

Effect on Sperm Count.—Serial analyses over many months showed that the sperm count invariably rose slowly and steadily for as long as two years. Two such cases are cited by way of example:

G.F., aged 29, had a pre-operative sperm count of 5 million/ml. Six months after varicocele ligation the count had risen to 21 million, and at the end of two years it was 126 million.

D.C., aged 26, had a pre-operative sperm count of 0.25 million/ml. Four months after ligation it had increased to 32 million, and, at the end of a year, it was 62 million.

As shown in Table II, 50 out of 72 cases with a pre-operative sperm count ranking of 2 or more improved to a higher rank within six months; 21 cases were still in the same rank; and one case dropped to a lower rank. Changes in the sperm counts of men with very severe pre-operative subfertility or complete sterility are shown in Table III, which demonstrates plainly that the improvement is much greater than could possibly be accounted for by "normal fluctuation." These readings were taken at six months; many of them were much higher at a later date.

TABLE II.—Sperm Count Rankings Before and Six Months After Varicocele Ligation

Before Operation		After Operation				
Sperm Count Rank	No. of Cases	No. of Cases in each Rank				
		1	2	3	4	5
1	16	16				
2	30	16	13	1		
3	17	7	8	2		
4	19	3	6	7	3	
5	6		1	1	1	3

TABLE III.—Effect of Varicocele Ligation on Cases of Azoospermia and Gross Oligospermia. (All Sperm Counts Expressed in Millions per ml.)

Case No.	Pre-operative Sperm Count	Post-operative Sperm Count	Case No.	Pre-operative Sperm Count	Post-operative Sperm Count
1	Azoospermia	38	14	2	9
2	"	14	15	2	8
3	"	0.5	16	2	7
4	"	Azoospermia	17	2	2
5	"	"	18	3	55
6	"	"	19	3	13
7	0.25	32	20	4	70
8	0.25	15	21	4	46
9	0.25	3	22	5	87
10	0.5	55	23	5	42
11	0.5	35	24	5	21
12	0.5	5	25	5	5
13	2	15			

Effect on Sperm Motility.—A visible improvement in the quality of sperm motility was the most striking feature of the post-operative analyses. Table IV shows, however, that there was also a significant improvement in the percentage of active sperms; no fewer than 48 of the 57 cases whose pre-operative motility was persistently under 10% had a post-operative activity of 50% or better within six months.

Effect on Pregnancy Rate.—This effect, so far as the patient is concerned, is the supreme test of success or failure of the operation, but it is appreciated that production of a successful conception cannot, by itself, be used as a yardstick to measure the value of any subfertility treatment as there are too many factors in both partners that must be taken into account. The seminal changes in 23 post-operative pregnancies are shown in Table V. All these cases had failed to produce

TABLE IV.—*Sperm Motility Rankings Before and Six Months After Varicocele Ligation*

Before Operation		After Operation		
Sperm Motility Rank	No. of Cases	No. of Cases in Each Rank		
		1	2	3
1	7	7		
2	18	13	5	
3	57	40	8	9

TABLE V.—*Seminal Changes in 23 Men Whose Wives Became Pregnant After Varicocele Ligation*

Case No.	Months Married Without Conception	Months Between Ligation and Conception	Before Operation		After Operation	
			Count Rank	Motility Rank	Count Rank	Motility Rank
1	122	24	3	3	2	1
2	120	6	2	3	1	2
3	84	3	1	2	1	1
4	72	2	3	2	3	2
5	72	3	2	3	1	1
6	60	11	3	3	1	1
7	62	5	2	3	2	1
8	64	3	3	2	1	1
9	49	3	1	3	1	1
10	48	5	2	3	1	1
11	47	5	2	3	1	1
12	44	6	4	3	2	1
13	37	4	4	2	3	2
14	35	3	2	2	1	1
15	29	8	1	3	1	1
16	27	5	3	3	2	3
17	26	3	3	3	2	1
18	24	9	2	2	1	1
19	51	4	2	2	Not received	
20	50	4	2	2		
21	37	3	3	3	"	
22	26	12	3	3		
23	84	3	2	3	"	

a pregnancy for more than two years without recourse to contraception. This period of "barren marriage" was chosen because Tietze (1956) has shown that under similar circumstances in the United States 91% of first conceptions would be expected to have occurred. It is therefore considered significant that within a few months of varicocele ligation 29% of these subfertile men had successfully impregnated their wives; although it is freely admitted that two of them showed no post-operative improvement (Cases 4 and 17) and one was still below the acceptable fertile level (Case 13).

Discussion

Although many American and Continental seminologists still fail to accept varicocele as a contributory factor to male subfertility, there is now, in my opinion, ample evidence to the contrary, although the exact mechanism is not yet clearly understood. There is considerable evidence to suggest that the seminal

changes are produced by increased intrascrotal temperature (Hanley, 1957). However, as similar high intrascrotal temperatures accompany large hydroceles, and as I have never been able to demonstrate any adverse effect on fertility in such cases, the increased temperature may only be an incidental finding. An alternative explanation may be that the damage is trophic in origin, due to some reversible type of reflex alteration in the calibre of the nutritional vessels (Harrison, 1958), because I have previously shown (Scott, 1958b), from testicular biopsy studies, that the tubular changes are bilateral although the varicocele is general unilateral.

The criteria for elective surgery (Scott, 1960b) demand careful selection of cases, the principle contraindications being previous disease of the opposite testis, particularly mumps orchitis (Scott, 1960c), and maldescent (Scott, 1961).

Whatever the explanation, a profound improvement occurs in the vast majority of cases, and a significant increase in the pregnancy rate can also be anticipated provided the patient's wife is fertile; if she is sterile there is no indication for operating on the husband.

Summary

The overall effects of varicocele, together with specific effects on the sperm count and motility, have been discussed. Significant post-operative improvement has been demonstrated in the sperm count in 78% of the cases, and in sperm motility in 81% of the cases; while the overall fertility was significantly improved in 83%. A pregnancy rate of 29% was achieved within a few months of operation. Possible mechanisms of the effect of varicocele on spermatogenesis are briefly discussed. The use of a ranking system for comparison of the pre- and post-operative seminal factors has clearly demonstrated that the changes after varicocele ligation are significantly greater than could be accounted for by normal fluctuation.

I thank my senior colleague, Mr. W. S. Mack, for his unfailing support and constructive criticism throughout the five years of this study.

REFERENCES

- Davidson, H. A. (1954). *Practitioner*, **173**, 703.
 Hanley, H. G. (1957). *Proceedings of 2nd World Congress on Fertility and Sterility*, Naples, 1956, **2**, 953.
 Harrison, R. G. (1958). In *La Fonction Spermatogénétique du Testicule Humain*, p. 63. Masson, Paris.
 MacLeod, J., and Gold, R. Z. (1956). *Fertil. and Steril.*, **7**, 394.
 Russell, J. K. (1954). *Brit. med. J.*, **1**, 1231.
 — (1957). *Lancet*, **2**, 222.
 Scott, L. S. (1958a). *J. Obstet. Gynaec. Brit. Emp.*, **65**, 504.
 — (1958b). *Proc. Soc. Study Fertil.*, **10**, 33.
 — (1959). *Scot. med. J.*, **4**, 354.
 — (1960a). M.D. thesis, University of Glasgow.
 — (1960b). *J. Reprod. Fertil.*, **1**, 45.
 — (1960c). *Brit. J. Urol.*, **32**, 183.
 — (1961). *J. Reprod. Fertil.* In press.
 Tietze, C. (1956). *Fertil. and Steril.*, **7**, 88.
 Tulloch, W. S. (1952). *Edinb. med. J.*, **59**, 29.
 — (1955). *Brit. med. J.*, **2**, 356.
 Young, D. (1956). *Brit. J. Urol.*, **28**, 426.

Leprosy: its Challenge and Hope, by Dr. R. G. Cochrane, has recently been published by the Mission to Lepers, and may be obtained from them at 7 Bloomsbury Square, London W.C.1, price 1s., or 1s. 4d. by post. The booklet covers the history of the disease, the position as it is to-day, and what measures can be taken to prevent and treat leprosy. all aspects being presented against a specifically Christian background. Dr. Cochrane is technical medical adviser to the American Leprosy Mission, Inc., and adviser on leprosy to the Ministry of Health, London. He was formerly director of the Christian Medical College, Vellore.