MEASUREMENT OF THE PROTECTIVE EFFECT OF ATTENUATED POLIOVIRUS VACCINE

BY

J. KNOWELDEN, M.D., D.P.H.*Department of Preventive Medicine and Public Health,
University of Sheffield***J. H. HALE, O.B.E., M.D., M.R.C.P.****P. S. GARDNER, M.D., Dip.Bact***Regional Public Health Laboratory, Newcastle upon Tyne*

AND

L. H. LEE, M.B., B.S.*Department of Bacteriology, University of Malaya, Singapore*

The assessment of the protective effect of attenuated poliovirus vaccines is particularly difficult. The technique which has been outstandingly successful with other vaccines, such as inactivated poliovirus vaccine, whooping-cough vaccine, and B.C.G., is that of the controlled prophylactic trial, but there are reasons why the method is not readily applicable to this particular problem. In the first place, many of the countries which experienced high morbidity from poliomyelitis in the past have used inactivated virus vaccines on a wide scale, and it would not be easy to allocate individuals in a population randomly to receive either inactivated or live virus vaccine; allocation at random to either attenuated virus or no vaccine at all is clearly unethical. Secondly, because of the widespread use of inactivated vaccines or for other reasons, the morbidity in most of these countries is now rather low and very large populations would have to be studied to give any chance of seeing sufficient paralytic poliomyelitis to provide an assessment. Thirdly, because there is the possibility of the vaccine virus spreading from individuals who had been given vaccine to their contacts the group given inactivated vaccine might include an unknown proportion of infected persons; as a consequence the poliomyelitis experience of the groups as allocated might not provide a true measure of the difference in performance of the prophylactics given to them.

Also, because of the risk of spread to contacts and the possibility of increase in virulence of the vaccine virus on repeated human passage, live virus vaccine has often been fed so far as possible to whole populations simultaneously and the deliberate withholding of live vaccine from part of the population, as required in a controlled trial, has been specifically avoided.

As an alternative to the controlled trial as a method of assessment, reliance has been placed either on the comparison between areas in which attenuated virus vaccines have or have not been used, or on the trend of morbidity before and after the use of such vaccines. The past experience of this disease has, however, been so unpredictable in both time and place that such comparisons cannot immediately indicate whether a specific prophylactic is responsible for any change in pattern. Repeated demonstrations of the association of vaccination with reduced morbidity would carry more conviction, but such evidence would not provide a quantitative assessment of protective effect; there would be no possibility, as existed, for example, in the Medical Research Council's trials of whooping-cough vaccines

(Medical Research Council, 1956), of measuring the relative performance in the field of different vaccines, or of relating such potency to the results of laboratory tests. In this respect the ability to produce antibodies may not be a valid method of comparing inactivated and attenuated vaccines because the former do not aim to produce alimentary immunity.

The Singapore Epidemic

Attenuated virus vaccines have been used on several occasions in the face of an epidemic. Here the problem of measuring the effect has additional complications; not only is it impossible to forecast what pattern the epidemic would have followed if vaccine had not been offered, but the vaccinated and unvaccinated groups are changing during the very period in which assessment is to be made. This situation existed in the type I epidemic during the later months of 1958 in Singapore, during which Sabin type II virus was fed to a proportion of children under 10 years of age (Hale *et al.*, 1959). The peak number of paralytic cases was recorded in the twelfth week of the epidemic ending October 25, and it was in this week that the first children were fed Sabin virus. Vaccine was given to nearly 200,000 of the half-million children under 10 years of age over a period of 12 weeks extending to mid-January, 1959, by which time the epidemic was virtually over.

In the early part of the vaccination period, when the incidence of poliomyelitis was high, the unvaccinated group was large and the vaccinated group small. By the end of this period the vaccinated group was relatively large, but the whole population, vaccinated and unvaccinated, was subject to a small risk. Even if the vaccine had been completely valueless the attack rate, counting cases occurring since vaccination started, would appear to be far smaller in the vaccinated group than in the remainder who were never vaccinated. It is difficult, therefore, to interpret the finding in the previous report that there were 6 type I paralytic cases among 200,000 vaccinated, compared with 179 among 300,000 unvaccinated. Furthermore, this comparison was made after the exclusion from the vaccinated group of cases whose onset was within a period after feeding less than the incubation period of the disease. Such exclusion might have introduced some bias, since a similar exclusion could not be made from the unvaccinated group. Finally, there was no way of determining what selection might have occurred in the group receiving vaccine, or how this might have influenced the relative state of immunity or risks of exposure to infection of vaccinated and unvaccinated groups.

Reanalysis of Singapore Data

As a solution to these difficulties the data from the Singapore epidemic have been reanalysed. The number of children vaccinated each week has been obtained (Table I), together with the age-distribution of vaccinations for the whole period of feeding (Table II). In the absence of specific information it has been assumed that the age-distribution in any one week was the same as that for the whole period. The population in four age-groups, 3-23 months, 2 years, 3-5 years, and 6-9 years, has been divided each week into three groups—unvaccinated, vaccinated less than eight days previously, and vaccinated eight or more days previously. For this purpose the number vaccinated less than eight days previously in any particular week *x* was taken to be

TABLE I.—*Number of Children Given Sabin Type II Poliovirus Vaccine in Singapore*

Week Ending	No. Vaccinated	Week Ending	No. Vaccinated
October 25	8,695	December 6	32,378
November 1	1,367	.. 13	20,797
.. 8	19,720	.. 20	6,938
.. 15	19,158	.. 27	677
.. 22	43,411	January 3	512
.. 29	42,606	.. 10	206

The total in this table, 196,465, includes a small proportion of children over 10 years of age.

TABLE II.—*Age Distribution of Vaccination*

Age	Total Population	No. Vaccinated	Percentage Vaccinated
3-23 months ..	96,303	42,899	44.5
2 years ..	53,154	22,834	42.9
3-5 ..	147,008	62,233	42.3
6-9 ..	171,861	61,617	35.9
3 months-9 years	468,326	189,583	40.5

the sum of half the vaccinations in the current week, x , plus half the vaccinations in the previous week $x-1$; the number vaccinated eight or more days previously was taken as the sum of half the vaccinations in week $x-1$, plus all the vaccinations in earlier weeks, $x-2$, $x-3$, etc. The cases of paralytic poliomyelitis were similarly divided into three groups according to whether they occurred in unvaccinated children or in children vaccinated less than eight days or eight or more days before onset.

Having ascertained these populations at risk, the ratios of each of the two vaccinated populations to the unvaccinated population in the same week were calculated. These ratios were then applied to the number of cases of paralytic poliomyelitis occurring in unvaccinated children to get expected numbers in those vaccinated less than eight days and eight or more days previously, on the assumption that vaccine was without effect. The expected and observed numbers in these vaccinated groups were then summed over the whole period. The details of the calculation are shown for the youngest age-group in Table III.

Table IV summarizes the analysis for all four age-groups. For those children vaccinated less than eight days before onset the total of 8 observed cases is close to the 10.02 expected on the null hypothesis, and in each

TABLE IV.—*Singapore, Paralytic Poliomyelitis in Vaccinated Children Under 10 Years of Age in the Period Week Ending October 25, 1958, to Week Ending January 24, 1959*

Age	Vaccinated Less than 8 Days before Onset		Vaccinated 8 or More Days before Onset	
	Observed	Expected at Rates in Unvaccinated	Observed	Expected at Rates in Unvaccinated
3-23 months ..	5	6.74	4	23.12
2 years ..	3	1.82	1	9.09
3-5 ..	0	1.15	0	6.06
6-9 ..	0	0.31	0	1.19
Total under 10 ..	8	10.02	5	39.46

of the separate age-groups there is close agreement. On the other hand, for the children vaccinated eight or more days before onset the total of 5 observed cases is very much less than the 39.46 expected, and again the pattern is consistent within each of the age-groups.

The reduction in the incidence of paralytic poliomyelitis in children vaccinated eight or more days before onset compared with that in unvaccinated children is impressive. It has already been noted, however, that one of the difficulties in interpreting such a pattern is the uncertainty whether vaccinated children were in some way selected and naturally subject to lower rates even without prophylaxis. The experience of children vaccinated less than eight days before onset suggests that such selection was not taking place. Within this period after vaccination their incidence was only a little less than that in unvaccinated children; as it is possible that vaccine virus had already begun to exert some protective effect by interfering with wild virus the eight observed cases may be fewer than those which would have occurred without vaccination. Whether such effect was acting or not, it is clear that the vaccinated children were not subject to such a reduced natural liability to infection as could explain their favourable experience after eight days from feeding.

It is concluded, therefore, that in the latter part of this type I epidemic in Singapore, children vaccinated with Sabin type II virus received considerable protection.

Summary

In an attempt to solve the problem of measuring the effectiveness of live poliovirus vaccine in preventing paralytic disease, the data recorded during the type I

TABLE III.—*Singapore, 1958-9. Children 3 Months to 2 Years of Age*

Week (x) (Ending)	No. of Children			Ratio Vaccinated/ Unvaccinated (%)		Poliomyelitis Cases				
	Unvaccinated	Vaccinated <8 Days	Vaccinated 8+ Days	<8 Days	8+ Days	Unvac- cinated (Observed)	Vaccinated <8 Days		Vaccinated 8+ Days	
							Obs.	Exp.	Obs.	Exp.
11 Oct. 18	96,303	—	—	—	—	—	—	—	—	—
12 " 25	95,354	949	—	1.00	—	20	—	0.20	—	—
13 Nov. 1	94,255	1,099	949	1.17	1.01	14	—	0.16	—	0.14
14 " 8	91,953	2,302	2,048	2.50	2.23	16	—	0.40	—	0.36
15 " 15	87,708	4,245	4,350	4.84	4.96	12	—	0.58	1	0.60
16 " 22	80,877	6,831	8,595	8.45	10.63	12	—	1.01	1	1.28
17 " 29	71,486	9,391	15,426	13.14	21.58	16	2	2.10	—	3.45
18 Dec. 6	63,300	8,186	24,817	12.93	39.21	8	2	1.03	—	3.14
19 " 13	57,495	5,805	33,003	10.10	57.40	10	1	1.01	—	5.74
20 " 20	54,467	3,028	38,808	5.56	71.25	4	—	0.22	—	2.85
21 " 27	53,636	831	41,836	1.55	78.00	2	—	0.03	1	1.56
22 Jan. 3	53,506	130	42,667	0.24	79.74	1	—	0.00	—	0.80
23 " 10	53,428	78	42,797	0.15	80.10	2	—	0.00	—	1.60
24 " 17	53,404	24	42,875	0.04	80.28	1	—	0.00	1	0.80
25 " 24	53,404	—	42,899	—	80.33	1	—	—	—	0.80
						Total	5	6.74	4	23.12

(1) At all ages a total of 196,465 children were vaccinated, of which 42,899 (21.8%) were aged 3 months to 2 years. It has been assumed that 21.8% of the children vaccinated each week were under 2 years of age.

(2) For any week, x , the number of children vaccinated (a) less than 8 days before onset has been taken as $\frac{1}{2}$ (vaccinations in week x) + $\frac{1}{2}$ (vaccinations in week $x-1$); (b) 8 or more days before onset has been taken as $\frac{1}{2}$ (vaccinations in week $x-1$) + (vaccinations in week $x-2$) + (vaccinations in week $x-3$), etc.

epidemic in Singapore in late 1958 and early 1959 have been analysed. Sabin type II vaccine was fed to about 200,000 of the half-million children under 10 years of age, beginning in the twelfth week of the epidemic. For each subsequent week estimates were made of the child populations in three categories: unvaccinated, vaccinated less than eight days, and vaccinated eight or more days previously. It is shown that the number of patients (8) who developed paralytic poliomyelitis less than eight days after vaccination was close to that expected at the rates in unvaccinated children (10.02), whereas the five cases observed in children vaccinated a longer period before onset were many fewer than the expected number of 39.46. It is concluded that a week after feeding the Sabin type II strain a substantial reduction in the risk of paralytic disease occurred as a result of vaccination.

REFERENCES

Hale, J. H., Doraisingham, M., Kanagaratnam, K., Leong, K. W., and Monteiro, E. S (1959). *Brit. med. J.*, 1, 1541.
Medical Research Council (1956). *Ibid.*, 2, 454.

RESULTS OF SURGERY IN CANCER OF LARGE INTESTINE IN A NON-TEACHING HOSPITAL

BY

G. S. ANDERSON, M.B., F.R.C.S.ED.

Lewis Hospital, Stornoway, Isle of Lewis

The large bowel is regarded by many as a site in which carcinoma, surgically treated, has a favourable prognosis, and this belief is supported by many reviews. Boyd (1955, p. 223) gives five-year survival rates of 55–60% for the right colon, 45–50% for the left colon, and 35–40% for the rectum. He adds that the five-year survival rate for early cases exceeds 80%. Many of the series reviewed have been of cases treated in special centres which do not provide a surgical service for the whole community.

Review of Series

The present group of cases includes all the patients with carcinoma of the large bowel, including those with acute obstruction, admitted to the two surgical units of the Royal Northern Infirmary, Inverness, a hospital of 220 beds, during the period 1944–53. All the deaths in the series are attributed to the carcinoma which brought the patient to hospital. Those untraced are presumed to have succumbed to their disease.

A total of 284 cases were admitted in the 10-year period. During this time there were 17,281 new surgical admissions, giving an incidence of colon and rectum carcinoma of 1 in 61. This is an average of four cases per general practitioner working in the area, so that

TABLE I.—Age Incidence

Age	10–19	20–29	30–39	40–49	50–59	60–69	70–79	80+
No. of patients	1	1	10	22	66	98	68	18

TABLE II.—Sex Incidence and Site

	Right Colon	Transverse and Left Colon	Rectum	Total
Males	16	63	81	160
Females	24	67	33	124

the family doctor referred one patient with carcinoma of the large bowel every two and a half years.

Age and Sex Incidence.—The ages of the patients ranged from 19 to 87 years, with the largest number in the seventh decade, and a rapid fall off in numbers in the preceding and following decades (Table I). There was an overall male preponderance of 160 males to 124 females, due to the larger number of rectal tumours in the male group (Table II).

Site Incidence (Fig. 1).—Approximately two-thirds of the tumours occurred in the rectum and pelvic colon, with rather more than one-third in the rectum. The remaining one-third were fairly evenly distributed round the remainder of the colon.

It is difficult to separate the high rectal growth from the low growth in the pelvic colon, but this was regarded as unimportant for the purpose of this review.

The average duration of the presenting symptom was fairly constant in each yearly group at seven to eight months. This is an inaccurate and minimum estimate, as many of the acutely ill obstructed cases gave a short history of a few days. It is not possible to divide this delay into that which was attributable to the patient and that to the medical adviser.

The investigation of these patients followed the well-established pattern and calls for no comment except to draw attention to a considerable delay in diagnosis which occurred on a few occasions when a barium follow-through was relied upon to show the colon. These were dyspeptic patients who turned out to have a right-sided colon tumour.

Treatment (Fig. 2).—In order to avoid a series of insignificantly small numbers, the treatment is divided

FIG. 1.—Site incidence.

FIG. 2.—Total number of cases in each year, with number in which a radical operation was attempted and the operative mortality.

into three simple groups—no surgery, palliative surgery, and radical surgery. The main radical operations used were right hemicolectomy where applicable and a V