6. Should it be necessary to stop giving poldine for some days—for example, during pyrexia—it is important to build up the dose again gradually to avoid the possibility of marked side-effects on restarting with a full dose. In every instance the omission of one dose has abolished all side-effects.

7. The full effect of the drug on gastric secretion and on symptoms may take up to 21 days to appear. In the meantime the dyspepsia can be treated with antacids.

8. It is important to realize that poldine does not completely abolish the dyspepsia in all patients. In our experience it greatly reduces the number of occasions on which there is pain and it reduces the severity of the pain. However, overwork—for example, decorating a flat in the evening, long hours of overtime in a bank, or concreting a garden path—has raised the gastric secretion in response to test meals on the following day, and sometimes apparently caused dyspepsia in spite of continuous treatment with poldine. It is remarkable how slow even intelligent patients are in relating their symptoms to overwork. They have in our experience always responded to antacids.

Summary

Fourteen men with duodenal ulcers and two with gastric ulcers have been studied for a mean period of three years.

During this time they have been under treatment with four daily doses of poldine methosulphate ("nacton") just small enough to produce no side-effects. No tolerance developed.

The secretion of acid, as assessed at monthly intervals with test meals, was reduced by one-half by the drug as compared with the secretion with dummy tablets. There was no change in gastric emptying.

The patients were free of dyspepsia for four-fifths of the months of treatment. As compared with an average absence from work of four weeks per patient per annum before treatment, the mean loss of time from work was half a day per annum during the three years of the study. During this period the deformity of the duodenum was reduced in 6 out of 14 patients with duodenal ulcer. These six had less dyspepsia than the others.

The trial has established that poldine can reduce the secretion of acid by about one-half in out-patients with peptic ulcer. Simultaneously, dyspepsia and loss of time from work were much reduced. The trial does not establish that the reduction of secretion of acid and the clinical improvement are cause and effect, although the results are consistent with this view.

We thank Drs. C. G. Baker, E. R. Boland, R. Kauntze, K. S. Maclean, and W. N. Mann for their co-operation in our studies of patients under their care. We are indebted to Beecham Research Laboratories Ltd. for generous supplies of nacton (poldine methosulphate) and prodexin (aluminium glycinate and magnesium carbonate).

REFERENCES

- Bettarello, A., Tuttle, S. G., and Grossman, M. I. (1960). *Gastroenterology*, **39**, 340.
 Douthwaite, A. H., and Hunt, J. N. (1950). *Lancet*, **1**, 1019.
 — (1958). *Brit. med. J.*, **1**, 1030.
 Clark, T., and Hunt, J. N. (1958). *Practitioner*, **180**, 334.
 Hunt, J. N. (1954). *Guy's Hosp. Rep.*, **103**, 161.
 — (1957a). *Lancet*, **1**, 132.
 — (1957b). *Brit. med. J.*, **1**, 681.
 — and Macdonald, I. (1952). *J. Physiol. (Lond.)*, **117**, 289.
 Seidelin, R. (1961). In press.
 Shay, H., Sun, D. C., Dlin, B., and Weiss, E. (1958). *J. appl. Physiol.*, **12**, 461.

ORAL TREATMENT OF PERNICIOUS ANAEMIA WITH VITAMIN-B₁₂-PEPTIDE

BY

N. K. SHINTON, M.B., M.R.C.P.

Consultant Haematologist, Coventry Hospital Group

The earliest treatment of pernicious anaemia was the inclusion of large quantities of liver or kidney in the patient's diet (Minot and Murphy, 1926; McCann, 1929). Oral administration of extracts and digests from these organs followed (Porter *et al.*, 1929; Davidson, 1932; Davis *et al.*, 1943), and remissions were also obtained by giving desiccated hog's stomach (Sturgis and Isaacs, 1929; Sharp, 1929; Wilkinson, 1930, 1949). All of these preparations, though effective, were unpleasant and could not be sterilized. They were therefore largely replaced by parenteral injections of liver extract and later of cyanocobalamin (vitamin B₁₂). Small amounts of the latter have been combined with extracts of desiccated hog's duodenal and pyloric mucosa to provide palatable oral substitutes, but in spite of good initial therapeutic results most long-term maintenance studies have shown high relapse rates (Adams, 1957; Bastrup-Madsen, 1957; Killander, 1957; Kristensen *et al.*, 1957; Lowenstein *et al.*, 1957; Schwartz *et al.*, 1957; Berlin *et al.*, 1958; Pedersen and Ebbesen, 1958; Stokes and Pitney, 1958; Blackburn *et al.*, 1959; Ellenbogen *et al.*, 1960).

Orally administered crystalline vitamin B₁₂ alone, either in single massive doses of 3,000–9,000 µg. or in smaller daily doses of 5–50 µg., have been shown to produce rises in haemoglobin concentration, red-cell count, and packed cell volume when given to cases of pernicious anaemia in relapse (Chalmers and Hall, 1954; Estren and Wassermann, 1956). Clinical and haematological remissions have been maintained by giving orally 1,000–3,000 µg. every one to four weeks (Conley and Krevans, 1955; Reisner *et al.*, 1955; Unglaub and Goldsmith, 1955) or 50–500 µg. each day (Chalmers and Shinton, 1958; Brody *et al.*, 1959; McIntyre *et al.*, 1960). However, with doses less than 150 µg. daily or 1,000 µg. weekly the serum vitamin B₁₂ concentration remained abnormally low in some patients.

It has more recently been claimed that patients with pernicious anaemia can be adequately maintained on oral daily doses of 10 µg. of vitamin B₁₂ when the latter is combined as a peptide complex derived from fermentation (Heathcote and Mooney, 1958). This has since been disputed by Varadi and Elwis (1960). Having made a study of orally administered crystalline vitamin B₁₂ alone in cases of pernicious anaemia (Chalmers and Shinton, 1958), the opportunity was taken of making a comparison, using identical dosage and test methods, with this vitamin-B₁₂-peptide in similar patients.

Materials and Methods

Seventeen consecutive cases of pernicious anaemia in relapse were studied. The diagnostic criteria included macrocytic anaemia, megaloblastic erythropoiesis, histamine-fast achlorhydria, absence of steatorrhoea or other known cause of megaloblastic anaemia. All had serum-vitamin-B₁₂ concentrations below 100 µg./ml. At the start of the trial none showed any neurological mani-

festations. All the patients were treated with oral vitamin-B₁₂-peptide from the stage of relapse and completed a short-term 15-day trial. Ten of them were maintained on the same regime for a further 3-13 months. The dose of oral vitamin B₁₂ given as B₁₂-peptide complex was 100, 200, or 500 µg. daily. It was taken in tablet form, each tablet containing 100 µg., as a single dose early each morning. Each patient took a standard dose throughout the trial. Haematological techniques and serum-vitamin-B₁₂ estimations were performed as described by Chalmers and Shinton (1958).

Results

Short-term (15-day) Studies

The changes in haemoglobin, red-cell count, packed cell volume, and serum-vitamin-B₁₂ concentration after 15 days of treatment with each dose are summarized in Table I. In all of the patients there was an improvement

TABLE I.—Pernicious Anaemia in Relapse Treated for 15 Days with Daily Oral Doses of Vitamin-B₁₂-Peptide

Case No.	Sex and Age	Daily Oral Dose (µg.)	Hb (g./100 ml.)		R.B.C. (mil./c.mm.)		P.C.V.		Serum Vitamin B ₁₂ (µg./ml.)	
			0*	15†	0	15	0	15	0	15
1	M 74	100	6.9	9.1	1.6	2.5	20	28	10	100
2	M 53	100	6.6	8.8	1.5	2.5	20	28	10	100
3	M 53	100	7.8	11.4	2.0	3.3	24	39	40	70
4	F 35	100	6.5	7.9	1.7	2.6	20	30	20	70
5	F 63	100	4.5	8.3	1.1	2.7	14	29	20	200
6	F 34	100	12.0	13.4	3.6	4.0	38	42	50	130
7	F 67	100	8.9	10.5	2.2	2.8	29	35	90	230
8	M 64	200	4.1	6.7	1.0	2.0	13	25	40	140
9	F 56	200	6.7	10.4	1.8	3.4	21	35	30	130
10	M 58	200	12.3	12.5	3.0	3.5	38	42	40	180
11	F 59	200	6.1	10.7	1.4	2.8	17	31	40	190
12	F 54	200	7.3	9.5	1.8	2.9	22	30	50	150
13	F 64	200	5.7	7.6	1.4	2.4	17	25	90	260
14	M 54	200	7.3	9.8	1.8	2.8	22	31	10	80
15	F 63	500	10.1	11.7	2.5	3.0	33	38	10	250
16	F 60	500	5.6	7.5	1.3	2.1	17	27	40	290
17	F 38	500	5.0	8.8	1.4	3.1	14	31	30	370

* Before treatment. † 15 days after treatment.

in their haematological values irrespective of the daily dosage received. The serum-vitamin-B₁₂ level from day 0 to days 5, 10, and 15 was found to rise progressively with the corresponding increase in the daily amount of vitamin-B₁₂-peptide given. These levels were compared with those previously obtained when giving similar doses of crystalline vitamin B₁₂ (Table II). Some of these results have been already reported (Cases 5-21, Chalmers and Shinton, 1958).

The mean serum-vitamin-B₁₂ concentration of all patients on days 0-15 for each respective preparation and dosage showed a progressive rise (see Chart), but when doses of only 100 µg. daily were given there was

a greater increase of these levels in those patients receiving the peptide preparation compared with those receiving an equivalent dose of crystalline vitamin B₁₂. An analysis of variance was therefore carried out on all these figures in order to extract the variations due to individual patients and to the rate of rise from day 0 to day 15. From the residual variance the significance of the overall rise in each group was assessed (Table III). From this it can be seen that the difference in the vitamin-B₁₂ levels between the two preparations when administering 100 µg. daily is significant, whereas with both 200 and 500 µg. daily no significant difference was found.

Maintenance Studies

Ten patients continued taking daily oral doses of the vitamin-B₁₂-peptide preparation from the stage of

Mean serum-vitamin-B₁₂ values in relation to the daily oral dose of crystalline vitamin-B₁₂ and vitamin-B₁₂-peptide during the first 15 days of treatment.

TABLE II.—Rise in Serum-Vitamin-B₁₂ Concentration (µg./ml.) from 0 to 15 Days in Cases of Pernicious Anaemia in Relapse Treated by Two Oral Preparations of Vitamin B₁₂

Case No.:		Vitamin-B ₁₂ -Peptide Complex								Crystalline Vitamin B ₁₂									
		1	2	3	4	5	6	7	Mean	5	6	8	22	23	24	25			Mean
100 µg. daily	{ Day 0	20	10	40	20	20	50	90	35	50	15	20	10	20	60	30			30
	{ " 5	40	20	40	20	120	80	150	70	50	30	30	20	40	130	40			50
	{ " 10	80	50	50	30	150	120	210	100	50	60	50	40	50	160	40			60
	{ " 15	100	100	70	70	200	130	230	130	75	85	60	40	80	160	70			80
Case No.:		8	9	10	11	12	13	14		11	12	13	26	27	28	29	30	31	
200 µg. daily	{ Day 0	40	30	40	40	50	90	10	40	20	30	30	40	40	30	10	30	60	30
	{ " 5	70	40	110	80	70	140	40	80	40	60	40	70	60	70	90	100	140	70
	{ " 10	100	80	150	170	130	180	70	130	60	150	50	80	120	110	110	110	260	120
	{ " 15	140	130	180	190	150	260	80	160	90	180	90	160	160	170	180	120	260	160
Case No.:		15	16	17						14	15	16	17	18	19	20	21		
500 µg. daily	{ Day 0	10	40	30					30	25	20	35	30	30	20	50	30		30
	{ " 5	160	200	180					180	190	200	180	310	150	170	150	110		180
	{ " 10	230	280	360					290	230	240	260	430	230	220	230	150		250
	{ " 15	250	290	370					300	260	280	280	490	260	260	250	210		290

relapse for periods varying from 3 to 18 months (Table IV). All but one of those receiving 100 μ g. daily (Cases 3-7) showed normal haematological values within two months, and these were maintained throughout the time of study. The exception (Case 4) failed to reach normal values after three months' therapy, and was therefore given intramuscular injections of 250 μ g. of crystalline vitamin B₁₂ every two weeks. Two months later a complete remission resulted.

All the patients receiving 200 and 500 μ g. daily reached and maintained haemoglobin, red-cell counts, and packed cell volumes within the normal range. The serum-vitamin-B₁₂ levels in each dosage group were broadly similar to those found on the 15th day of the study. Of those receiving 100 μ g. daily, only Case 4 showed any appreciable difference in level after three months' therapy, the levels ranging at the end of the trial from 80 to 260 μ g./ml. In the three cases receiving 200 μ g. daily (Cases 8, 9, and 10) rises after 15 days were only small, the figures ranging from 160-170 μ g./ml. after periods of three to nine months. A progressive rise for three months was found in the two patients (Cases 15 and 16) receiving 500 μ g. daily, in both cases the levels being maintained around 400 μ g./ml. for a further 18 months. One patient (Case 9) who received 200 μ g. daily developed peripheral neuritis in the upper limbs after three months' therapy and was therefore changed to intramuscular injections. The remainder of the patients developed no abnormal neurological signs or symptoms.

Discussion

For an oral preparation of vitamin B₁₂ to be accepted as a satisfactory form of treatment of pernicious anaemia it must be established that it will reliably cause a complete clinical and haematological remission which will be maintained. It would also be desirable for it to be capable of replenishing the body stores, which are mainly in the liver. When there is a deficiency of the vitamin, the levels in the liver are similar to those in the plasma (Grasbeck, 1959), and it is therefore reasonable to assume that low serum levels in all circumstances reflect a deficiency in the body stores. If this is accepted,

TABLE III.—Residual Variances Between Serum-Vitamin-B₁₂ Concentrations After Treatment with Vitamin-B₁₂-Peptide as Compared with Crystalline Vitamin-B₁₂

Daily Dosage	Variance Ratio	Degrees of Freedom	P
100 μ g.	36	1/36	≤ 0.001
200 "	1.4	1/42	> 0.2
500 "	1.0	1/27	> 0.2

TABLE IV.—Long-term Maintenance Therapy with Daily Oral Doses of Vitamin-B₁₂-Peptide

Case No.	Values at Beginning of Trial					Months of Treatment	Values at End of Trial				
	Hb (g./100 ml.)	R.B.C. (mil./c.mm.)	P.C.V. (%)	Serum Vitamin B ₁₂ (μg./ml.)	Daily Oral Dose (μg.)		Hb (g./100 ml.)	R.B.C. (mil./c.mm.)	P.C.V. (%)	Serum Vitamin B ₁₂ (μg./ml.)	
3	7.8	2.0	24	30	100	6	13.8	4.7	43	80	
4	6.5	1.7	20	20	100	3	10.2	4.2	35	140	
5	4.5	1.1	14	30	100	3	12.4	4.7	41	200	
6	12.0	3.6	38	60	100	18	11.5	3.6	37	170	
7	8.9	2.2	29	90	100	6	13.1	4.6	43	260	
8	4.1	1.0	13	40	200	3	13.6	4.6	41	160	
9	6.7	1.8	21	40	200	9	15.2	5.0	46	170	
10	12.3	3.0	38	40	200	5	14.6	4.8	44	160	
15	10.1	2.5	33	10	500	18	13.3	4.5	39	420	
16	5.6	1.3	17	40	500	18	13.7	4.6	40	390	

satisfactory treatment of pernicious anaemia also requires that the serum-vitamin-B₁₂ concentration should be at least within the normal range and preferably up to the mean level of about 400 μ g./ml. (Shinton, 1959). The importance of maintaining a normal serum-vitamin-B₁₂ level in patients receiving oral treatment has been emphasized by reports that where relapse is about to occur decline in this level precedes that of other haematological values (Bastrup-Madsen, 1957; Killander, 1957; Kristensen *et al.*, 1957; Lowenstein *et al.*, 1957; Blackburn *et al.*, 1959).

Furthermore, the development of neurological manifestations has been described in patients maintained on oral preparations with serum levels of 50-150 μ g./ml. (Bastrup-Madsen, 1957; Kristensen *et al.*, 1957; Varadi and Elwis, 1960). The peripheral neuritis that developed in one patient of the present series where the serum vitamin B₁₂ was 170 μ g./ml. was probably another example of this danger. Regular estimations of serum-vitamin-B₁₂ levels must therefore be made when an oral preparation is being used.

The claims by Mooney and Heathcote (1960, 1961) that vitamin-B₁₂-peptide is superior to crystalline B₁₂ in the oral treatment of pernicious anaemia were based on therapeutic trials in four cases where one preparation followed the other without interval. The responses observed while the patients were receiving the vitamin B₁₂-peptide could therefore have been a carry-over effect of the crystalline vitamin B₁₂ given primarily, such a delayed response having been previously reported (Chalmers and Shinton, 1958). However, in the present investigation it has been shown that when 100 μ g. daily of crystalline vitamin B₁₂ or its equivalent dose of vitamin-B₁₂-peptide is given to individual but similar groups of cases there is a greater rise in the mean serum-vitamin-B₁₂ level in those receiving the peptide complex. This concurs with the demonstration by Milhaud (1961) that in patients who have had a partial gastrectomy the absorption of radioactive vitamin B₁₂ in small doses was greatly increased when given in the form of a peptide complex.

These considerations, while of academic interest, have little bearing on the possible therapeutic advantages of vitamin-B₁₂-peptide, as, like the crystalline form, it is necessary to give a daily dose of at least 200 μ g. in order to be sure that the serum-vitamin-B₁₂ level is being maintained near the normal mean. At this dosage no difference was found between the two preparations. There may be special circumstances where oral treatment of pernicious anaemia is desirable (*Brit. med. J.*, 1959), but in the routine management of these patients the view previously expressed by Chalmers and Shinton (1958) remains that intramuscular injections of vitamin B₁₂ provide the most reliable method of ensuring a rapid response, and maintenance free of the risk of neurological complications.

Summary

The use of vitamin-B₁₂-peptide as a possible form of oral treatment for pernicious anaemia has been assessed clinically, haematologically, and by serum-vitamin-B₁₂ estimations.

While it has been shown to be similar in effect to crystalline vitamin B₁₂ when given orally in doses of 200 μ g. daily or more, with 100 μ g. daily there is a comparatively greater rise of the serum-vitamin-B₁₂ level over 15 days.

Parenteral therapy remains the most reliable method of treating pernicious anaemia, but should this be contraindicated then generous daily doses of either crystalline vitamin B₁₂ or vitamin-B₁₂-peptide are as effective as any other form of oral therapy yet developed.

I thank the physicians who referred cases to me for these trials; the patients for their co-operation; Dr. J. A. H. Waterhouse for help with the statistical analysis; Dr. J. N. Marshall Chalmers for his co-operation and advice; Mr. C. I. D. Wilson for technical assistance; and Distillers Co. (Biochemicals) Ltd. for supplying the vitamin-B₁₂-peptide.

REFERENCES

- Adams, J. F. (1957). *Scot. med. J.*, **2**, 151.
 Bastrup-Madsen, P. (1957). *Acta med. scand.*, **159**, 323.
 Berlin, R., Berlia, H., Brante, G., and Sjöberg, S.-G. (1958). *Ibid.*, **161**, 143.
 Blackburn, E. K., Spray, G. H., Swan, H. T., Tudhope, G. R., and Wilson, G. M. (1959). *Brit. med. J.*, **2**, 535.
Brit. med. J., **1959**, **2**, 554.
 Brody, E. A., Estren, S., and Wasserman, L. R. (1959). *New Engl. J. Med.*, **260**, 361.
 Chalmers, J. N. M., and Hall, Z. M. (1954). *Brit. med. J.*, **1**, 1179.
 — and Shinton, N. K. (1958). *Lancet*, **2**, 1298.
 Conley, C. L., and Krevans, J. R. (1955). *Ann. intern. Med.*, **43**, 758.
 Davidson, L. S. P. (1932). *Brit. med. J.*, **2**, 347.
 Davis, L. J., Davidson, L. S. P., Riding, D., and Shaw, G. E. (1943). *Ibid.*, **1**, 655.
 Ellenbogen, L., Williams, W. L., and Lichtman, H. C. (1960). *Ibid.*, **2**, 1066.
 Estren, S., and Wasserman, L. R. (1956). *Proc. Soc. exp. Biol. (N.Y.)*, **91**, 499.
 Grasbeck, R. (1959). *Advanc. clin. Chem.*, **3**, 336.
 Heathcote, J. G., and Mooney, F. S. (1958). *Lancet*, **1**, 982.
 Killander, A. (1957). *Ibid.*, **1**, 1041.
 Kristensen, H. P. Ø., Lund, J., Sjøberg Ohlsen, A., and Pedersen, J. (1957). *Ibid.*, **1**, 1266.
 Lowenstein, L., Brunton, L., Shapiro, L., de Leeuw, N., and Dufresne, M. (1957). *Canad. med. Ass. J.*, **77**, 923.
 McCann, W. S. (1929). *Proc. Soc. exp. Biol. (N.Y.)*, **25**, 255.
 McIntyre, P. A., Hahn, R., Masters, J. M., and Krevans, J. R. (1960). *Arch. intern. Med.*, **106**, 280.
 Milhaud, G. (1961). *Nature (Lond.)*, **189**, 33.
 Minot, G. R., and Murphy, W. P. (1926). *J. Amer. med. Ass.*, **87**, 470.
 Mooney, F. S., and Heathcote, J. G. (1960). *Lancet*, **2**, 291.
 — (1961). *Brit. med. J.*, **1**, 232.
 Pedersen, J., and Ebbesen, I. (1958). *Acta med. scand.*, **161**, 413.
 Porter, W. B., Williams, J. P., Forbes, J. C., and Irving, H. (1929). *J. Amer. med. Ass.*, **93**, 176.
 Reisner, E. H., Weiner, L., Schittone, M. T., and Henck, E. A. (1955). *New Engl. J. Med.*, **253**, 502.
 Schwartz, M., Lous, P., and Meulengracht, E. (1957). *Lancet*, **1**, 751.
 Sharp, E. A. (1929). *J. Amer. med. Ass.*, **93**, 749.
 Shinton, N. K. (1959). *Clin. Sci.*, **18**, 389.
 Stokes, J. B., and Pitney, W. R. (1958). *Brit. med. J.*, **1**, 322.
 Sturgis, C. C., and Isaacs, R. (1929). *J. Amer. med. Ass.*, **93**, 747.
 Unglaub, W. G., and Goldsmith, G. A. (1955). *Sth. med. J. (Bgham, Ala.)*, **48**, 261.
 Varadi, S., and Elwis, A. (1960). *Lancet*, **2**, 653.
 Wilkinson, J. F. (1930). *Brit. med. J.*, **1**, 236.
 — (1949). *Lancet*, **1**, 291.

Dr. C. S. P. HAMILTON, of Glenageary, Eire, has presented a pewter medal to the Maritime Institute Museum, Dun Laoghaire. The medal was one of a number presented to the crew of the British sloop *Calliope* by the citizens of Apia on the occasion of the hurricane which struck Apia and wrecked several German and American warships in 1889. The *Calliope* was the only ship to sail for the open sea when the hurricane struck. The others threw out extra anchors and attempted to ride out the storm. Some foundered and others were driven ashore. When the *Calliope* returned after the hurricane her Irish commander, Captain Kane, and his crew set to work to bring order out of the destruction in the harbour. In gratitude for their efforts and in admiration of Captain Kane's superb seamanship, the leading residents of Apia gave the captain a silver medal, and each member of the crew a pewter medal. One of these pewter medals, which was given to an Irish sailor, eventually came into the possession of Dr. Hamilton. (*New Zealand News*.)

HAEMOGLOBIN Q-ALPHA-THALASSAEMIA

BY

KATHARINE M. DORMANDY, M.B., B.S.
M.R.C.P.

S. P. LOCK, M.A., M.B., B.Chir.

Hospital for Sick Children, Great Ormond Street, London

AND

H. LEHMANN, M.D., Sc.D., M.R.C.P., F.R.I.C.
St. Bartholomew's Hospital, London

Three years ago a new haemoglobin, Q, was described by Vella, Wells, Ager, and Lehmann (1958); it was found in combination with haemoglobin H. Since then there has been much progress in our knowledge of the fine structure of haemoglobin, which we owe largely to Ingram, Schroeder, Itano, and their colleagues. The present paper describes the case of a child in whom haemoglobin Q was found together with haemoglobin Bart's. Taken together with the previous case (Vella *et al.*, 1958), and utilizing the new information on the structure and inheritance of human haemoglobin, our findings make it possible to define a new syndrome—Hb Q- α -thalassaemia.

Case Report

A 10-months-old boy from Thailand, who had developed severe atopic eczema at the age of 7 months, was admitted to the Hospital for Sick Children for treatment. Apart from the eczema, physical examination revealed no abnormal findings; examination of the blood film, however, showed numerous target cells, and further investigations were therefore carried out.

Laboratory Investigations

The results of routine laboratory investigations are summarized in the Table. A stained blood film (Fig. 1, A) showed considerable hypochromia, moderate anisocytosis and poikilocytosis, and many target cells. Except for an eosinophilia, the white cells and platelets were normal. On supervital staining with brilliant cresyl blue a moderate number of red cells were found to contain inclusion bodies (Fig. 2), similar to those found in cells containing haemoglobin H. Red-cell osmotic fragility was greatly diminished, as shown in Fig. 3. The direct Coombs test was negative. The serum bilirubin was 0.2 mg./100 ml. and serum-protein electrophoresis was normal.

The combination of a high serum-iron level with a hypochromic blood picture suggested a diagnosis of thalassaemia. On examination of the haemoglobin by paper electrophoresis at alkaline pH three fractions were observed; the fastest, comprising about 5%, moved in the position of Hb Bart's; the second and major component was found just behind Hb A, in the position of Hb Q; and the third, which amounted to 2% of the whole, was just behind the position of Hb A₂ (Fig. 4). Hb A and Hb A₂ themselves were not found. The finding of Hb Bart's was confirmed by paper electrophoresis at acid pH. On chromatography in "amberlite IRC 50" at pH 6, no Hb A was discovered, and the major component again moved in the position of Hb Q. Hb F was not found, although 2.4% alkali-resistant haemoglobin was shown to be present by the one-minute alkali-denaturation test (Singer, Chernoff, and Singer, 1951). This was explained by the presence of 5% Hb Bart's, which is about half as resistant to alkali as Hb F in this test (Ager and Lehmann, 1958). In several experiments Hb H was not discovered on electrophoresis, although the inclusion bodies (Fig. 2) suggested that some had been present. However, in one experiment with large amounts of very