(4) The mortality from tumours of bones other than limb bones shows only a faint peak at ages 20–24 years.

(5) At ages under 45 years the mortality is approximately equal in both sexes.

(6) At ages 45 years and over there is a sharp increase in the mortality among men but very little increase among women. In the absence of Paget's disease the discrepancy is more marked and among women the mortality remains practically constant. The male excess in this group is due principally to tumours of the ribs and shoulder-girdle.

No significant difference in the geographical distribution of bone-tumour mortality has been demonstrated.

We are grateful for the helpful advice of Dr. R. Acheson, Dr. J. H. Marshall, and Dr. J. M. Tanner.

APPENDIX

In 25 cases it was impossible to decide whether a primary bone tumour was present. There was no reference to Paget's disease on any of the death certificates. The descriptions of these cases given on the death certificates were:

Case No.	Sex	Age (Yrs.)	Affected Bone	Histological Description	Place of Death
1	M	3	Maxilla	Osteosarcoma	Home
2	M	6	Jaw	Sarcoma	Hospital
3	M	8	Spine	"	Home
4	M	18	Calcis	Ewing's tumour	"
5	M	37	Ilium	Sarcoma	Hospital
6	M	45	Femur	"	Home
7	M	46	Ribs	"	Hospital
8	M	50	Pelvis	Osteosarcoma	Home
9	M	56	Skull	Sarcoma	"
10	M	59	Femur	"	"
11	M	59	Humerus	Malignant neoplasm	"
12	F	9	Lumbar spine	Tumour	"
13	F	23	Fibula	Ewing's sarcoma	"
14	F	26	Ribs	Sarcoma	"
15	F	33	Rib	"	Hospital
16	F	48	Ilium	"	Home
17	F	50	Pelvis	Osteosarcoma	"
18	F	50	"	Chordoma	Hospital
19	F	57	Vertebrae	Sarcoma	"
20	F	60	Ethmoid	"	Home
21	F	60	Rib	"	"
22	F	61	Femur	"	Hospital
23	F	63	Orbit	"	Home
24	F	63	Lumbar spine	Carcinoma	"
25	F	64	Pelvis	Sarcoma	Hospital

REFERENCES

- Aub, J. C., Evans, R. D., Hempelmann, L. H., and Martland, H. S. (1952). *Medicine (Baltimore)*, **31**, 221.
- Brues, A. M., Lisco, H., and Finkel, M. (1946). U.S. Atomic Energy Commission Report.
- Court Brown, W. M., Doll, R., Heasman, M., and Sissons, H. A. (1961). In press.
- Finkel, M. P. (1956). *Proceedings of International Conference on Peaceful Uses of Atomic Energy*, **2**, 160. United Nations, New York.
- Looney, W. B. (1956). *Amer. J. Roentgenol.*, **75**, 559.
- Hasterlik, R. J., Brues, A. M., and Skirmont, E. (1955). *Ibid.*, **73**, 1006.
- Marinelli, L. D. (1958). *Ibid.*, **80**, 729.
- Martland, H. S. (1931). *Amer. J. Cancer*, **15**, 2435.
- Price, C. H. G. (1955). *Brit. J. Cancer*, **9**, 558.
- (1958). *J. Bone Jt. Surg.*, **40B**, 574.
- Registrar-General of England and Wales (1953). *Statistical Review of England and Wales for the Year 1952*. Tables, Part I, Medical, H.M.S.O., London.
- (1954). *Statistical Review of England and Wales for the Year 1953*. Tables, Part I, Medical, H.M.S.O., London.
- (1955). *Statistical Review of England and Wales for the Year 1954*. Tables, Part I, Medical, H.M.S.O., London.
- Tanner, J. M. (1961). *Growth at Adolescence*, 2nd ed. Blackwell, Oxford.
- Turner, R. C., Radley, J. M., and Mayneord, W. V. (1961). *Nature (Lond.)*, **189**, 348.
- United Nations Scientific Committee on the Effects of Atomic Radiation (1958) Report. New York.
- Vaughan, J. (1956). In *Lectures on Scientific Basis of Medicine*, **4**, 196.
- World Health Organization (1949). *Manual of the International Statistical Classification of Diseases, Injuries and Causes of Death*, 6th revision, 1948. H.M.S.O., London.

PLEA FOR GREATER USE OF CYTOLOGICAL STUDIES IN EARLY DETECTION OF CANCER

BY

J. G. S. CRABBE, M.R.C.S., L.R.C.P.

AND

MARY E. EGERTON, M.B., B.S., M.R.C.O.G.

From the Department of Pathology, Royal Free Hospital, London

Among the most important factors influencing the prognosis of malignant disease are the extent of the growth and the degree of its spread. By the time a tumour gives rise to symptoms it may have reached proportions which render treatment difficult and the prognosis poor. The examination of body fluids by the cytological method makes possible the discovery of malignant disease when the tumour is still of microscopic proportions, and thus detection at such an early stage can help to improve prognosis. Furthermore, the method is also of considerable assistance in helping to differentiate malignant disease from other pathological conditions, particularly where other diagnostic tests may be inconclusive.

The cytological method originated in England over 100 years ago, when Walshe (1851) observed fragments of malignant tissue in sputum. From that time onwards other workers, both in this country and elsewhere, carried out much work in this field, and in 1927 Dudgeon and Patrick, experimenting with the wet-fixation technique at St. Thomas's Hospital, examined sputum, urine, and other body secretions for the presence of malignant cells. At about the same time in the United States of America, Papanicolaou (1928) used a similar technique for diagnosing cancer of the female genital tract, and by 1940, having realized the potentialities of the method, he devoted his time wholly to its development, so that to-day, after many years of extensive trial all over the world, it has become inevitably associated with his name and now is used as an everyday routine for cancer detection in most hospitals in the Americas and on the Continent. In Great Britain much excellent cytological work has also been done in a number of centres for some years, but, as Saxton (1960) points out, this country lags far behind both the U.S.S.R. and the U.S.A., where cytological examinations for cancer of the uterus have become routine procedures, and, although in a few localities schemes for this purpose are in operation, he pleads for the establishment of more such schemes in all centres of population where these examinations could be carried out regularly.

Principal Uses of the Cytological Method

Female Genital Tract.—Undoubtedly the most successful application of the cytological method has been its use for the detection of early invasive or pre-invasive carcinoma of the cervix uteri. In Great Britain, Anderson (1950), Wachtel and Plester (1952), Way (1953), McLaren (1953), and Egerton (1957), after trying out the method as a research project, found that their results were so encouraging that they now continue to use it as a routine test in the gynaecological clinics of the hospitals where they work. Wrigley (1960) maintains that it is essential as an additional means of investigation where there is the slightest sign or suspicion

of malignant disease. He also states that it should be used as a routine procedure when changing a ring pessary and for following up treatment for cervical carcinoma. The method is much less reliable for diagnosing carcinoma of the body of the uterus; nevertheless, Wachtel and Plester (1952) were able to detect not only five cases of unsuspected endometrial carcinomata among 2,000 patients examined but also four unsuspected ovarian adenocarcinomata. Moore (1958) at the Royal Society of Medicine demonstrated a case from the Royal Free Hospital in which an unsuspected carcinoma in a Fallopian tube was discovered by this means.

Urinary Tract.—Cytology is now used as an essential method in the control of bladder tumours of occupational origin, such as those occurring among workers in the dyestuffs and rubber industries (Crabbe, Cresdee, Scott, and Williams, 1956). Since its introduction as a diagnostic measure for this purpose it has already helped to prolong the lives of most of the men who have been affected. Between the years 1951 and 1956 some 1,800 men had their urine examined routinely by the method and 62 new cases of bladder tumours were discovered. Although similar work has been carried out in America, Germany, and Italy for some years no results had been published from these countries until last year, when Melamed, Koss, Ricci, and Whitmore (1960) in America reported on some observations on bladder tumours developing in workers exposed to para-aminodiphenyl, a powerful bladder carcinogen. Apart from diagnosing occupational bladder tumours the examination of urine may be of help in hospital routine in detecting bladder growths where cystoscopy is contraindicated or refused and in locating or confirming the diagnosis of tumours in the renal pelvis or ureters. It is of little or no use for detecting tumours of the renal parenchyma.

A further application in the genito-urinary field is for the detection of carcinoma of the prostate by the examination of fluid obtained by massage, and Fergusson and Gibson (1956) found that the method was reliable for indicating prostatic malignancy as it gave results comparable with those of biopsy without the attendant risk from trauma, haemorrhage, and infection. They found that it could be safely repeated many times and that it affords a useful method for indicating response or otherwise to hormonal therapy.

Respiratory System.—With the increase in the incidence of lung cancer during the last few decades the examination of sputum and bronchial washings for the presence of malignant cells is a most valuable addition to current methods for helping to detect these growths. Sometimes it is the only means of detecting the disease at a stage early enough for successful treatment to be instituted or of confirming a clinical diagnosis when x-ray examination is equivocal and bronchoscopy negative. Some of the earliest systematic cytological investigations on sputum were performed by British pathologists (Dudgeon and Patrick, 1927; Gloyne, 1937; Schuster, 1943; Bamforth, 1946).

Pleural and Ascitic Fluids.—The cytological examination of serous effusions can be of great diagnostic and prognostic significance, as the occurrence of ascites or a pleural effusion in malignant disease is a grave and often terminal phenomenon. The results of the examination can be of great help not only in making a differential diagnosis but sometimes as a guide to the effectiveness of treatment.

Alimentary Tract.—Washings from the stomach are also examined for malignant cells, and Schade (1958), using a simple lavage technique, states that small malignant lesions, often confined to the mucosa, were easier to detect than the larger advanced tumours. Burnett, MacFarlane, Scott Park, and Kay (1960), also using a similar technique, found that cytology was a useful adjunct to other routine methods of diagnosis for gastric carcinoma, but owing to the high incidence of false-negative results (38%) cytology was unreliable as a screening test. Oakland (1957) uses the method for the examination of washings from the colon and rectum, and he found that it was most successful in the diagnosis of carcinoma of the descending and sigmoid colons.

Technique

The making and staining of the smears is so simple that it can be done in any small histological laboratory. The actual details of technique vary in different cytological schools. Pathologists tend to prefer fixation of the smear while it is still wet and to stain with haematoxylin and eosin (Bamforth, 1946; Philips, 1954), while the haematologists prefer drying and fixation followed by one of the Romanowsky stains (Spriggs, 1957). On the other hand, the followers of Papanicolaou and those engaged in endocrine work favour wet fixation and the use of the polychrome stains of Papanicolaou (1954) or Shorr (1940).

All of these techniques are simple, but it really does not matter which method is used, it is purely a question of preference. The best method is that with which the cytologist is most familiar. The most popular is probably that of Papanicolaou, and in his *Atlas* he describes fully the different ways of obtaining the materials, making the smears, and how to stain them by his method. Finally, for the proper examination of the smears the equipment must include a good binocular microscope with a mechanical stage and a reasonably powerful and accurate source of illumination for dry high-power work. The reading of the slides is greatly facilitated if $\times 10$ or $\times 15$ wide-field eyepieces are used. The interpretation of the smears and the criteria of malignancy have been described many times over in the copious literature on the subject, and these points again are fully discussed by Papanicolaou in his *Atlas*.

Use of the Cytological Technique at the Royal Free Hospital

The encouraging results of the cytological research work of Egerton (1957) in the gynaecological department of the Royal Free Hospital since 1952 have led to the adoption of the technique as a routine cancer detection test which is now applied to all women aged 30 and over attending the clinics. The method is also used in the department as an indication of ovarian function in the endocrine disorders of women (Egerton, 1957).

Although the cytological study of other body secretions, notably sputum (Findlayson, 1956) and serous effusions, has been done in the hospital for some years it was not until 1958 that a department was set up devoted entirely to the cytological examination of all types of materials.

The specimens examined include gastric washings as a screening test for carcinoma, but so far the experience here has been similar to that of Burnett *et al.* (1960) in that the incidence of false-negative results is too high

for the method to be reliable for this purpose. The accompanying Table is a summary of the results from May, 1958, to December, 1960.

Summary of Results

Specimens	No. of Patients	Tumours Found	Positive Smears	Con-firmed	False Positives	False Negatives	Not Con-firmed
Gynaecological	3,000	60	60 (100%)	48 (80%)	11 (18.3%)	12 (20%)	1
Respiratory ..	275	46	32 (69.6%)	30 (65.2%)	1 (2.03%)	16 (34.8%)	1
Serous effusions	147	57	51 (89.5%)	46 (80.7%)	0	11 (19.3%)	5
Urine	40	9	8 (88.9%)	6 (66.7%)	1 (11.1%)	3 (33.3%)	1
Gastric washings	228	15	6 (40%)	5 (33.3%)	0	10 (66.7%)	1

Of the gynaecological cases 16 (26.7%) were unsuspected.

Female Genital Tract

Since May, 1958, 3,000 cases have been examined; 48 tumours have been found by means of the cytological technique (39 cases of carcinoma of the cervix, 8 of carcinoma of the fundus, and 1 case of carcinoma of the ovary). Of these 48 cases, 16 had no signs or symptoms suggestive of carcinoma and were entirely unsuspected. Five proved on biopsy to be carcinoma *in situ*, eight proved to be early invasive carcinoma, and three proved to be adenocarcinoma of the body of the uterus. The remaining 32 cases were all clinically suspected. In four of these 32 cases it was possible to confirm the clinical suspicion by immediate staining and reading the smears before the patients left the department. In 11 cases the smears were not confirmed by biopsy, while one remains to be confirmed. Twelve cases showed no malignant cells in the smears and all of them proved to be carcinomas of the body of the uterus.

Example, Case 1.—The patient had been attending the gynaecological clinic since September, 1959, for treatment of a persistent vaginal discharge due to trichomonal infection. In November, 1959, smears were taken from the cervix and posterior fornix purely as a routine, as she had become 30 years of age during the course of her treatment. Malignant cells were seen in the smears and biopsy confirmed the presence of an early invasive carcinoma of the cervix. Irradiation therapy was followed by total hysterectomy. At the time of writing she was alive and well and was attending the gynaecological clinic for follow-up smears.

Respiratory Tract

Since May, 1958, 275 cases have been examined; 32 of these had malignant cells in the sputum, and 30 of them were proved to have carcinoma of the lung or bronchus. In one case no carcinoma was found at necropsy while the remaining case has yet to be confirmed. In 16 cases with proved carcinoma of the respiratory tract no malignant cells were found in the sputum.

Example, Case 2.—In January, 1960, a man aged 46 had a fit of coughing which was followed by a severe pain in the front of the chest that lasted for two or three days. This cleared up but recurred more severely in July, but this time it lasted longer and much brown sputum was produced. In September he was admitted to hospital, where it was noted that he had night sweats and was losing weight. An x-ray examination on September 8 showed a large opacity with cavitation in the left upper lobe which might have been due to a breaking-down growth, an abscess, or aspergillus infection. Smears of the sputum taken at this time contained malignant cells suggestive of a squamous-cell carcinoma. A week later bronchoscopy showed a stricture of the left upper lobe bronchus probably due to neoplasm. The biopsy

section taken at the same time confirmed the presence of a squamous-cell carcinoma. He was discharged from hospital at the end of September after a lobectomy and is alive and well.

Pleural and Ascitic Fluids

Smears of 147 cases have been examined and 51 of these had malignant cells. In three of these cases the cytological reports were the first indication that the condition was due to malignant disease, and in two other cases the report was of great help in confirming the presence of a suspected neoplasm. However, in 11 cases where a tumour was found to be present no definitely malignant cells were seen. Five cases remain to be confirmed.

Example, Case 3.—A man aged 60 was admitted to hospital in December, 1959, complaining of abdominal discomfort and distension for four days. A tentative diagnosis of cirrhosis of the liver was made, but the ascitic fluid was found to contain many clusters of malignant cells. X-ray and barium-meal examinations were suggestive of oesophageal varices, but gastric carcinoma could not be completely excluded. His condition rapidly deteriorated and he died in January, 1960. Necropsy disclosed cirrhosis of the liver together with a carcinoma of the head of the pancreas, but there was no carcinoma of the stomach. The smears were the only tests to give a definite indication that malignant disease was present.

Urinary Tract

Only 40 specimens of urine have been examined since May, 1958. Eight of these cases had positive smears; six were proved to have tumours on further investigation, while in one case no tumour was found on cystoscopy and one case has yet to be confirmed. Three specimens sent from patients, later proved to have hypernephromata, contained no malignant cells. The usefulness of the technique is illustrated in the following case report.

Example, Case 4.—This patient, aged 68, had a carcinoma of the cervix uteri which was treated successfully in 1938. She came to the hospital in January, 1959, after an attack of haematuria, and it was thought that she might have a recurrence of this tumour which was involving the bladder. On gynaecological examination there was no clinical evidence of a local recurrence, so a catheter specimen of urine was obtained for cytological examination. The smears of the urine sediment contained many clusters of malignant cells which were suggestive of a transitional-cell carcinoma. On cystoscopic examination a bladder tumour was found which, on sectioning, proved to be a transitional-cell carcinoma. A partial cystectomy was performed in February, 1959, and the patient is alive and well.

Gastric Washings

Of 288 cases examined, 15 were suspected of having carcinoma of the stomach. Of these 15, only 6 showed malignant cells in the smears. Five were confirmed at operation, while one remains to be confirmed. Ten cases proved by other means of investigation to have gastric carcinoma showed no malignant cells in smears of the gastric washings.

Example, Case 5.—The patient, aged 44, was admitted to hospital in May, 1960, complaining of severe epigastric pain, which was relieved by vomiting and which had persisted for the last five years. X-ray examinations were inconclusive, but the smears of the gastric washings contained clusters of cells which were highly suspicious. Sections of gastric biopsy material showed the presence of gastric carcinoma, and this was confirmed by gastroscopy. A partial gastrectomy was performed in July, and sections of the stomach showed it to be a multifocal early gastric carcinoma. The patient was discharged from hospital in August and is alive and well.

Conclusions

In the comparatively few centres in this country where the cytological diagnosis of cancer has been practised for some years the successes have been such that the demand for its use has tended to increase. This is exemplified at the Royal Free Hospital, where during a brief period of some six years the successes of one or two research workers have resulted in the establishment of a department devoted exclusively to cytological cancer diagnosis with facilities for both teaching and research work. During this time the great advantages of the technique have become apparent. Firstly, and most important, as a routine procedure it can be used to detect tumours at a very early stage of development, often before signs and symptoms appear. It is simple to perform and need cause no alarm, inconvenience, or discomfort to the patient, and thus it can be repeated as often as may be necessary. The only inconvenience that a patient may be caused is in obtaining those materials where needles or other instruments have been used.

The chief disadvantages of the method are the time that is required to read the slides and the lack of trained personnel to do this. The taking and processing of the smears is simple, but the interpretation of the wide range of the cellular variations in the different types of body fluids requires considerable training. Until recently it was usually found necessary to go to America to gain the requisite experience, but now one or two cytological centres in London and the provinces provide facilities for teaching the technique and the interpretation of the smears. This should help to eliminate the necessity for sending personnel abroad for training, with the concomitant expense which that involves.

A plea is made for the more extended use of the method throughout the British Isles, where much of the original pioneer cytological work was done. The experience of cytological workers all over the world shows that the technique is well worth while, for if the detection of cancer when it is symptomless and at a very early stage of development is going to save or prolong lives—and there is reason to believe that it does—then the method is a most valuable weapon in the fight against cancer.

Summary

An account is given of the cytological technique for the early detection of malignancy which, though evolved in the early days to a great extent by British pathologists, was developed by Papanicolaou in America in such a way that it has practical application.

The principal and most successful uses of the method are described and reference is made to the processing, staining, and interpretation of the smears.

Although most of the published work on cytology comes from American and Continental laboratories, the references in this paper apply chiefly to publications from British cytologists.

A short account is given of the establishment of a cytology department and its use at the Royal Free Hospital during the last two years, with a record of the results obtained. Typical example cases are quoted. The advantages and disadvantages are discussed, and a plea is made for the more extended use in this country of this undoubtedly valuable weapon in the fight against cancer.

We thank Professor K. R. Hill for his encouragement and Miss Anne Robertson for carrying out the technical work and helping to read the slides. We are indebted to the British Empire Cancer Campaign for the generous help with a grant from which technical and laboratory assistance has been provided, thus enabling this work to be carried out.

REFERENCES

- Anderson, A. F. (1950). *J. Obstet. Gynaec. Brit. Emp.*, **57**, 3.
 Bamforth, J. (1946). *Thorax*, **1**, 118.
 Burnett, W., MacFarlane, P. S., Park, S. D. S., and Kay, A. W. (1960). *Brit. med. J.*, **1**, 753.
 Crabbe, J. G. S., Cresdee, W. C., Scott, T. S., and Williams, M. H. C. (1956). *Brit. J. Industr. Med.*, **13**, 270.
 Dudgeon, L. S. and Patrick, C. V. (1927). *Brit. J. Surg.*, **15**, 250.
 Egerton, M. E. (1957). *34th A.R. Brit. Emp. Cancer Campgn.*, p. 237.
 — (1957). *J. Obstet. Gynaec. Brit. Emp.*, **64**, 827.
 Fergusson, J. D., and Gibson, E. C. (1956). *Brit. med. J.*, **1**, 822.
 Findlayson, R. (1956). M. D. Thesis, Univ. Cambridge.
 Gloyne, S. R. (1937). *Tubercle (Lond.)*, **18**, 292.
 McLaren, H. C. (1953). *31st A.R. Brit. Emp. Cancer Campgn.*, p. 181.
 Melamed, M. R., Koss, L. G., Ricci, A., and Whitmore, W. F. (1960). *Cancer*, **13**, 67.
 Moore, J. A. M. (1958). *Proc. roy. Soc. Med.*, **51**, 172.
 Oakland, D. (1957). *Brit. med. J.*, **1**, 1391.
 Papanicolaou, G. N. (1928). *Proceedings of Third Race Betterment Conference*, p. 528.
 — (1954). *Atlas of Exfoliative Cytology*, pp. 3–12, 13–21, Harvard Univ. Cambridge, Mass.
 Philips, F. R. (1954). *Brit. J. Cancer*, **8**, 67.
 Saxton, R. S. (1960). *Brit. med. J.*, **1**, 1657.
 Schade, R. O. K. (1958). *Ibid.*, **1**, 743.
 Schuster, N. (1943). *Practitioner*, **151**, 20.
 Shorr, E. (1940). *Science*, **91**, 321.
 Spriggs, A. L. (1957). *The Cytology of Effusions*, p. 1. Heinemann, London.
 Wachtel, E. and Plester, J. A. (1952). *J. Obstet. Gynaec. Brit. Emp.*, **59**, 323.
 Walshe, W. H. (1851). *A Practical Treatise on the Diseases of the Lungs, Heart, and Aorta*, p. 412. Walton and Maberly, London.
 Way, S. (1953). *31st A.R. Brit. Emp. Cancer Campgn.*, p. 186.
 Wrigley, A. J. (1960). *Practitioner*, **184**, 459.

THYROGLOBULIN ANTIBODIES IN 1,297 PATIENTS WITHOUT THYROID DISEASE

BY

OSCAR W. HILL, B.A., M.B., M.R.C.P.*

Registrar, Professorial Medical Unit, Middlesex
Hospital, London

Since the work of Roitt and Doniach (1960) and others it is known that the sera of patients suffering from thyroid disorders, especially Hashimoto's disease, often show antibodies to thyroglobulin. The present investigation reports the frequency with which antibodies may be found in subjects not suffering from any apparent thyroid abnormality.

Materials and Methods

The sources of the sera tested were 597 samples sent to the laboratory for performance of the Wassermann test and 700 sent to the blood transfusion service for grouping, the former preserved at 4° C. for a week and the latter at -20° C. The notes were examined of all patients whose sera contained thyroid antibodies, and cases with clinically observed thyroid abnormalities were removed from the main series and made into a subgroup for the purpose of comparing the incidence of complement-fixing antibodies in the two groups (see Table II).

*Now at the Maudsley Hospital, London.