Figure 1 Periorbital xanthogranuloma. (A) Clinical appearance with infiltration of all four eyelids and mild blepharoptosis with an S-shape configuration. (B) Visibly enlarged lacrimal gland. (C) Coronal T1 fat-suppressed gadolinium-enhanced orbital MRI demonstrating diffuse hyperintensity of the periorbital tissues including the lacrimal gland. Informed consent was obtained for publication of this figure.

lacrimal glands and follicular conjunctivitis (fig 1). Ophthalmological examination was otherwise unremarkable, with normal symmetric exophthalmometry measurements and full extraocular motility. Orbital MRI demonstrated bilateral diffuse periorbital infiltration with lacrimal gland involvement (fig 1C). A left anterior inferior orbitotomy was performed. All tissue planes (epidermis to orbit fat) were

involved and obscured with diffusely infiltrative, firm, bright yellow material.

H&E-stained sections displayed several large follicles surrounded by lipid-laden histiocytes and Touton giant cells infiltrating the fat, muscle and fibrous tissue (fig 2). Immunohistochemical staining confirmed the benign nature of the follicles, staining positive for CD20 and negative for κ , λ and bcl-2. A few CD3 cells were seen in the mantle zone and scattered throughout the surrounding tissue, particularly around the xanthogranulomatous cells. They stained positive for κ and λ with a ratio of 5:1. Flow cytometry suggested a κ monoclonal B cell population (ratio 10:1). PCR using three sets of primers (FR3A/VLJN, FR2A/VLJH and FR2B/VLJH) detected no heavy-chain gene rearrangements. No abnormality consistent with a lymphoproliferative disorder or ECD was detected on systemic evaluation (chest and abdominal CT, bone scan, extremity plain films and protein electrophoresis of serum and urine).

Oral prednisone (60 mg daily) achieved a partial response, with recurrence on tapering the drug. External beam radiation (2080 rad in 13 divided doses) resulted in complete clinical resolution, which persisted 1 year later.

Comment

This report is unique in that xanthogranulomatous inflammation seems to have been caused or at least precipitated by periorbital

surgery. In most reported cases, no mention is made of periocular surgery.¹⁻³ However, this is not the first case presumably related to trauma. A recent report suggested a relationship between xanthogranuloma of the ear lobes and mechanical injury due to earrings.⁴ In our patient, an entirely causal relationship seems unlikely; however, surgical insult might have stimulated/amplified subclinical disease. Admittedly, even this is unconfirmed and the exact nature of the relationship between surgical insult and xanthogranulomatous inflammation remains to be determined.

Christopher I Zoumalan

Department of Ophthalmology, University of California
San Francisco, San Francisco, California, USA

**Melanie H Erb, Narsing A Rao, Robert See,
Michael A Bernstein**

University of Southern California, Doheny Eye Institute,
Los Angeles, California, USA

Samir B Shah, Timothy J McCulley

Department of Ophthalmology, University of California
San Francisco, San Francisco, California, USA

Correspondence to: Dr T J McCulley, Department of
Ophthalmology, University of California San Francisco,
10 Koret Way, San Francisco, CA 94143-0336, USA;
mcculleyt@vision.ucsf.edu

This paper was presented in part at the Walsh Society
meeting at Snowbird, Utah, February 2003.

doi: 10.1136/bjo.2006.107821

Accepted 9 December 2006

Competing interests: None declared.

References

- 1 Jakobeic FA, Mills MD, Hidayat AA, *et al.* Periorbital xanthogranulomas associated with severe adult-onset asthma. *Trans Am Ophthalmol Soc* 1993;**91**:99-125.
- 2 Hammond MD, Niemi EW, Ward TP, *et al.* Adult orbital xanthogranuloma with associated adult-onset asthma. *Ophthalmic Plast Reconstr Surg* 2004;**20**:329-32.
- 3 Karioglu ZA, Sharara N, Boles TL, *et al.* Orbital xanthogranuloma: clinical and morphologic features in eight patients. *Ophthalm Plast Reconstr Surg* 2003;**19**:372-81.
- 4 Sueki H, Saito T, Iijima M, *et al.* Adult-onset xanthogranuloma appearing symmetrically on the ear lobes. *J Am Acad Dermatol* 1995;**32**(Pt 2):372-4.

The Finger iridectomy technique for glaucoma

Surgical iridectomy is a standard method of treatment for narrow-angle glaucoma.¹ However, the development of laser iridectomy has largely replaced the need for incisional surgery.¹ There are cases where patients are unable or unwilling to submit to laser iridectomy, when surgical manipulation of the iris is required and when the cornea is not sufficiently clear. This case demonstrates the first use of a 25-gauge aspiration cutter through a 1 mm self-sealing corneal incision to perform a surgical iridectomy for glaucoma.

Case report

An 80-year-old woman was noted to have a variably pigmented inferonasal iris tumour, lenticular pseudoexfoliation and narrow angles in her left eye. The tumour was documented to grow and cause a sector cataract (prompting her referral to The New York Eye Cancer

Figure 2 Periorbital xanthogranuloma (H&E $\times 200$). Chronic inflammatory cells and lipid-laden histiocytes infiltrating the muscle fibres and fibrous tissue as well as multiple Touton giant cells (arrows) displaying the characteristic central eosinophilic cytoplasm surrounded by multiple nuclei and peripheral foamy/vacuolated cytoplasm are seen.

Figure 1 (A) A slit-lamp photograph of the Finger iridectomy technique (FIT) result. Also note the complex melanotic and amelanotic iris melanoma in the inferonasal quadrant. (B) Transillumination through the superonasal iridotomy demonstrates its patency. (C) A 35 MHz high-frequency B-scan ultrasonography (ultrasound biomicroscopy) prior to FIT iridectomy demonstrates a severely narrowed anterior chamber angle (before small-incision surgical iridotomy for glaucoma). (D) At 3 weeks after successful FIT iridectomy, the anterior chamber angle has widened.

Centre, New York City, New York, USA). Ophthalmic examination included a 35 MHz high-frequency ultrasound (ultrasound biomicroscopy) in movie mode. Both her narrow angles and tumour were evaluated (fig 1).

Methods of procedure

This study conformed to the tenets of the Declaration of Helsinki² and the Health Insurance Portability and Accountability Act of 1996.³ The Finger iridectomy technique (FIT) was performed for glaucoma immediately after an FIT the iris tumour biopsy was found positive for melanoma.⁴ A 0.3 forceps was used to stabilise the eye at the limbus. A 25-gauge trochar was used to create an incision through clear juxta-limbal cornea into the anterior chamber. The first incision was made on the same side as the tumour to avoid the pupil and lens. Acetylcholine chloride 10 mg/ml was introduced to induce miosis, then sodium hyaluronate 1% was filled to maintain the anterior chamber.

For iridectomy, a second 25-gauge incision was made in the superotemporal cornea.⁵

Using sodium hyaluronate 1%, the aspiration cutter was inserted and the port rotated for occlusion by the superonasal iris stroma. Aspiration cutting (suction of 300 mm Hg and cutting rate of 600 cuts/min) was used to perform a single full-thickness superonasal iridectomy (fig 1). Once engaged with iris tissue, aspiration, cutting and full-thickness iridectomy are almost instantaneous. At the end of surgery, the sodium hyaluronate 1% was removed, and peribulbar antibiotic-steroid and topical ocular hypotensive agents were used. No adverse side effects were noted.

At 3 weeks after iridectomy, slit-lamp photography and high-frequency ultrasound were used to document the patient's patent iridectomy and deepening of her anterior chamber

(fig 1). Her intraocular pressure decreased by 5 mm Hg (off medications).

Comment

A review of the literature shows that relatively large aspiration cutters have been used to perform iridectomy.⁶ For example, Ghanem *et al*⁶ used an aspiration cutter during phacemulsification in patients with iridoschisis. The FIT is different in that it is a minimally invasive approach using a smaller 25-gauge aspiration cutter probe to perform localised iridectomy through a self-sealing 25-gauge incision. No large incision, irrigation or sutures are required in FIT.⁴

The FIT introduces the concept of using a 25-gauge aspiration cutter to perform a minimally invasive iridectomy for glaucoma. Unlike standard surgical iridectomy, the FIT 25-gauge corneal incision allows for a relatively safe self-sealing corneal wound.

Paul T Finger

The New York Eye and Ear Infirmary, New York City, New York, USA

Correspondence to: Dr P T Finger, The New York Eye Cancer Center, 115 East 61st Street, New York City, NY 10021, USA; pfinger@paultfingermd.com

doi: 10.1136/bjo.2006.108126

Accepted 18 October 2006

Funding: This work was supported by The EyeCare Foundation, New York City, NY, USA.

Competing interests: None.

References

- 1 Quigley HA. Surgery of the glaucomas. In: Rice TA, Michels RG, Stark WJ, eds. *Ophthalmic surgery*, 4th edn. St Louis: Mosby, 1984, 177–81 (Chapter 7).
- 2 World Medical Association. Declaration of Helsinki (2000). Ethical Principles for Medical Research Involving Human Subjects. *Bull Med Eth* 2000;162:8–11.
- 3 Health Insurance Portability and Accountability Act of 1996. Public law 104–191. 104th Congress, United States of America, Aug 21, 1996.
- 4 Finger PT, Latkany P, Kurli M, *et al*. The Finger iridectomy technique: small incision biopsy of anterior segment tumours. *Br J Ophthalmol* 2005;89:946–9.
- 5 Finger PT. Small incision surgical iridectomy and iridectomy. *Graefes Arch Clin Exp Ophthalmol* 2006;244:399–400.
- 6 Ghanem VC, Ghanem EA, Ghanem RC. Iridectomy of the anterior iris stroma using the vitreocutter during phacemulsification in patients with iridoschisis. *J Cataract Refract Surg* 2003;29:2057–9.

Preimplantation genetic diagnosis for retinoblastoma predisposition

Heritable mutations in the *RB1* gene cause an autosomal dominant condition resulting in retinoblastoma^{1,2} and an increased risk of malignancies including pineoblastoma, neuroblastoma, chondrosarcoma, rhabdomyosarcoma, glioma, leukaemia, sebaceous carcinoma, squamous cell carcinoma and cutaneous melanoma.^{3–6} Individuals with heritable retinoblastoma can undergo prenatal diagnosis followed by termination to avoid passing on the mutation to the next generation.⁷ Preimplantation genetic diagnosis (PGD) offers a means of achieving an unaffected pregnancy from the outset. IVF is required for PGD to allow cell biopsy from embryos for genetic testing. Embryos without the germline *RB1* mutation are transferred to the mother for implantation and pregnancy.

Case study

A 24-year-old woman with bilateral retinoblastoma (RB1, OMIM#180200), had a *de novo* M708R mutation in *RB1* and was referred for PGD. She had a one-year-old child with the mutation and also had had a miscarriage and two terminations of affected pregnancies.

Following ovarian stimulation, 15 eggs were collected, of which 12 oocytes matured and eight fertilised normally 18–20 hours post-intracytoplasmic sperm injection (ICSI). Seven embryos were suitable for biopsy on day 3. DNA from biopsied cells was amplified by PCR using fluorescently labelled intragenic (RB120) and linked polymorphic micro-satellite markers (D13S168 and D13S262). The PCR products were analysed using an ABI 3730 Prism. Three embryos (E4, E6 and E7) were diagnosed as carrying the maternal affected chromosome, while two embryos (E5 and E1) did not give clear results. Embryos E3 and E8 were diagnosed as unaffected. Figure 1 shows the haplotype for the family and figure 2 shows electrophoretograms of the parental DNA, a normal and an affected blastomere. All cell negatives and PCR negatives showed no DNA amplification.

On day 4 post-insemination unaffected embryos E3 (5 cell) and E8 (morula), were transferred to the uterus. Six+ weeks gestation ultrasound showed two sacs, one with a viable fetal pole while the other was anembryonic. A healthy boy was delivered at 35/40 weeks. Here we report the first successful use of PGD for retinoblastoma in the UK.