Botulinum toxin for the treatment of acute-onset concomitant esotropia in Chiari I malformation

Acute-onset esotropia (ET) is a rare presentation of Chiari I malformation. The ET may resolve following neurosurgical decompression, although this is not usually immediate.¹ Where neurosurgery is not undertaken, Kowal *et al*² suggest prismatic correction or strabismus surgery. The latter may result in temporary correction of the strabismus, as the strabismus can recur and resolves only following decompression.³ Botulinum toxin (BT) has been reported as successful in one case where the ET did not resolve following neurosurgery.⁴ Despite BT being a common treatment for acute acquired concomitant ET,⁵ no previous case has been reported in which BT was used prior to neurosurgical decompression for Chiari I malformation.

Case report

A 16-year-old girl experienced diplopia during swimming, and presented to the A&E Department on the day of onset. She gave a history of occasional, brief diplopia. On examination, there was 16Δ ET at near and 20Δ ET at distance. The angle of deviation reduced on versions. Saccade to the right was possibly hypometric; no nystagmus was present. Over 2 weeks, the angle increased to 40Δ ET, saccades to the right became definitely hypometric and abducting nystagmus developed. MRI scan revealed Chiari I malformation with tonsillar descent to about 12 mm below the foramen magnum, with loss of cerebrospinal fluid (CSF) spaces indicating impaction of neural tissue (fig 1). Neurosurgery was discussed, but delayed for 8 months.

Three months after onset, the angle had increased to 45Δ ET at near and distance. Four units of BTA were injected into the right medial rectus muscle with electromyographic guidance under topical anaesthesia. Initially there was an overcorrection (12ΔXT, 5ΔRHT, with $-3\frac{1}{2}$ limitation right adduction). Following recovery of medial rectus function, 6Δ exophoria at near and 4Δ esophoria at distance were present. Eight months following BT injection there had been no recurrence of strabismus and the patient underwent neurosurgical decompression. Currently the patient is orthophoric and shows hypometric saccades to the right and left, and fine pendular-like nystagmus on right and left gaze.

Figure 1 MRI scan showing significant tonsillar descent approximately 12 mm below the foramen magnum with loss of cerebrospinal fluid spaces around the foramen magnum indicative of impaction of neural tissue.

Comment

BT restored binocular single vision (BSV) in a patient awaiting neurosurgical decompression for Chiari I malformation. Neurosurgery was delayed to fit in with the patient's education. BT was used in the expectation that repeat injections would be necessary as the aetiological cause had not been treated at this time. Following the BT, BSV was maintained for 8 months prior to decompression surgery. Recurrence of ET following strabismus surgery has occurred over a variety of time frames upwards from 3 months following strabismus surgery.⁶

Surgical decompression improves ocular manifestations such as nystagmus, but this may not be for several months.⁷ In our case, nystagmus and hypometria had not resolved 3 months following surgery.

BT should be considered as a treatment option to restore alignment in acute-onset ET due to Chiari I malformation. Further case series are required to determine the effectiveness of BT compared with strabismus surgery.

Alison Y Firth

Academic Unit of Ophthalmology and Orthoptics,
University of Sheffield, Royal Hallamshire Hospital,
Sheffield, UK

John P Burke

Department of Ophthalmology, Sheffield Teaching
Hospitals NHS Foundation Trust, Royal Hallamshire
Hospital, Sheffield, UK

Correspondence to: Alison Y Firth, Academic Unit of
Ophthalmology and Orthoptics, University of Sheffield,
Royal Hallamshire Hospital, Sheffield, UK; a.firth@
sheffield.ac.uk

doi: 10.1136/bjo.2006.111104

Competing interests: None declared.

References

- 1 Defoort-Dhellemmes S, Denion E, Arndt CF, *et al*. Resolution of acute acquired comitant esotropia after suboccipital decompression for Chiari I malformation. *Am J Ophthalmol* 2002;133:723-725.
- 2 Kowal L, Yahalom C, Shuey NH. Chiari I malformation presenting as strabismus. *Binocul Vis Strabismus Q* 2006;21:18-26.
- 3 Passo M, Shults WT, Talbot T, *et al*. Acquired esotropia. A manifestation of Chiari I malformation. *J Clin Neuro-ophthalmol* 1984;24:83-86.
- 4 Imes RK, Quinn TA. Acute comitant esotropia in Chiari I malformation. *Ophthalmology* 2001;108:834.
- 5 Dawson ELM, Marshman WE, Adams GGW. The role of botulinum toxin A in acute-onset esotropia. *Ophthalmology* 1999;106:1727-1730.
- 6 Weeks CL, Hamed LM. Treatment of acute comitant esotropia in Chiari I malformation. *Ophthalmology* 1999;106:2368-2371.
- 7 Liebenberg WA, Georges H, Demetriades AK, *et al*. Does posterior fossa decompression improve oculomotor and vestibulo-ocular manifestations in Chiari I malformation? *Acta Neurochir* 2005;147:1239-1240.

Microfoam surgical tape as practice object for scleral sutures

Performing scleral sutures in strabismus and retina surgery poses a challenge for residents as it requires high precision and allows only little room for error, with possible complications including endophthalmitis and retinal detachment from scleral perforation or muscle dehiscence.

The human sclera consists of collagen and elastin, with a thickness ranging between 0.33 mm beneath the recti muscles, 0.66 mm at the muscle insertion and 1 mm posteriorly.

Figure 1 3M Microfoam surgical tape, 2-inch width.

The three layers of the sclera are ill defined and comprise episclera, sclera proper and lamina fusca.¹

Pig eyes, which are commonly used as practice objects for surgical procedures, are not always readily available, change their consistency after conservation in formalin and potentially present a risk for contamination or infection. Other commonly used practice objects include fruits such as oranges, bananas and grapes, or hard-boiled eggs.

Microfoam surgical tape manufactured by 3M consists of rolls of closed-cell foam and is designed for securing dressings and compression applications to joints, with a thickness of 0.8 mm and width of 1, 2 or 3 inches (fig 1). It is elastic and has an adhesive lower surface.

When used as a practice object for scleral bites, achievement of appropriate suture depth can be easily checked by lifting the superficial layer from underlying layers (fig 2). In addition, the elastic tape can be easily stretched to 0.4 mm for simulation of a thinner sclera.

Our department's retina and strabismus specialists evaluated Microfoam surgical tape as a practice object and uniformly agreed on its authentic feel, resistance, thickness and curvature when compared with the actual human sclera.

In conclusion, Microfoam surgical tape represents a readily available, cheap, storable and easily transportable alternative to pig eyes or fruits for practicing the placement of scleral sutures with authentic characteristics and immediate feedback.

Jan Niklas Ulrich, Thomas W Wilson

Geisinger Medical Center, Danville, Pennsylvania, USA

Correspondence to: Dr J N Ulrich, Geisinger Medical
Center, 500 Pine Street, Apt 5, Danville, PA 17821,
USA; npilic@web.de

doi: 10.1136/bjo.2006.111336

Competing interests: None declared.

Reference

- 1 Friedman N, Kaiser P, Trattler W. Cornea/external disease. *Review of ophthalmology*. Pennsylvania: Elsevier Saunders, 2005.

The use of voriconazole in the treatment of *Aspergillus fumigatus* keratitis

There are a few reports of the use of voriconazole for the treatment of fungal keratitis.¹⁻³ We report another case of its apparent success in *Aspergillus fumigatus* keratitis and discuss the dilemma of

Figure 2 Scleral bites of different depth (top left, bottom left and bottom centre): too superficial (left suture), too deep (centre suture) and adequate depth (right suture). View from underneath (top right, bottom right): suture too superficial (left suture, not visible), suture perforates sclera (centre suture) and adequate depth (right suture).

prescribing an expensive drug in the absence of defined ophthalmic therapeutic levels.

A 51 year old diabetic man presented a week after being poked in his eye by a child's finger, with a 3.0×3.5 mm central corneal ulcer overlying full thickness stromal infiltrate with associated hypopyon (fig 1). Pinhole visual acuity was 2/60 in this eye. Topical ciprofloxacin 0.3% was started. *Aspergillus fumigatus* was isolated after 48 hours' incubation and topical amphotericin B 0.15%, voriconazole 1% hourly, and oral voriconazole 800 mg twice daily were introduced, the latter reducing to 400 mg twice daily after 48 hours. Topical voriconazole was prepared aseptically with 0.9% sodium chloride, with 48 hour expiry (24 hours from opening). The ulcer size, infiltrate, and hypopyon slowly reduced, although recurrent hypopyon at day 13 led to rescraping, aqueous paracentesis, and administration of intracameral amphotericin. A bacillus and coagulase-negative staphylococcus were isolated following enrichment culture. With no further treatment alteration the ulcer healed. At six weeks the eye appeared quiet with a corneal scar, 30% stromal thinning and visual acuity of 6/24; oral voriconazole was discontinued and topical treatments four weeks later.

Comment

Fungal keratitis is difficult to treat and carries a significant risk of intraocular involvement. Traditional antifungals usually have good topical

transcorneal penetration, but even systemic therapy has limited intraocular penetration. Primary treatment failure is reported in 31%, with predictive factors being *Aspergillus* infection, large ulcer size, and hypopyon.⁶ We therefore used a novel agent in this high risk case.

Voriconazole is a highly potent triazole, with 100% in vitro susceptibility reported in 34 common ocular fungal pathogens, compared to only 60–82.4% for fluconazole, itraconazole, amphotericin B, and ketoconazole.⁷

Although unlicensed in this capacity, topical and oral voriconazole (4–6 mg/kg twice daily for 4–12 weeks) are reported in the successful treatment of *Fusarium*,^{1,2} *Scedosporium*,³ *Alternaria*,⁴ *Candida albicans*, and *Aspergillus fumigatus* keratitis.⁵ Despite initial good response, two of these cases also showed disease exacerbation at 10 to 14 days which settled with intracameral infusion or further/increased systemic treatment or both.^{1,5} Intracameral amphotericin may have been instrumental in our case, but this is difficult to disentangle from the voriconazole effect.

Orally administered, voriconazole reaches therapeutic levels in the aqueous and vitreous of the non-inflamed eye,⁸ and aqueous of the infected eye.³ Its penetration topically is less clear: although levels exceeding the minimum inhibitory concentration are reported in the cornea, vitreous, and chorioretina of rabbit control and infected eyes,⁹ human data are conflicting.^{2,5}

The good ocular penetration of oral voriconazole has encouraged its use. Although the properties of this agent are attractive¹⁰ its cost far exceeds that of traditional antifungals. A 28 day course costs £3645.44 orally (400 mg twice daily) and £2884.00 topically (hourly), compared with only £86.24 for oral fluconazole (200 mg four times a day) and £190.40 for topical (hourly) amphotericin B.

Restricting voriconazole use to cases involving resistant organisms is difficult as there are no ophthalmic data on resistance and susceptibility breakpoints. Guidelines for the treatment of fungal keratitis need to be established.

Figure 1 Central corneal ulcer with dense elevated infiltrate and hypopyon.

Rosalind M K Stewart, Say Aun Quah,
Timothy J Neal, Stephen B Kaye
Royal Liverpool University Hospital, Liverpool, UK

Correspondence to: Mr Stephen B Kaye, St Paul's Eye
Unit, 8Z Link, Royal Liverpool University Hospital,
Prescot Street, Liverpool L7 8XP, UK;
s.b.kaye@liverpool.ac.uk

doi: 10.1136/bjo.2006.111989

Accepted 15 January 2007

Competing interests: None declared.

References

- 1 Reis A, Sundmacher R, Tintelnot K, *et al.* Successful treatment of ocular invasive mould infection (fusariosis) with the new antifungal agent voriconazole. *Br J Ophthalmol* 2000;**84**:932-3.
- 2 Klont RR, Eggink CA, Rijs AJMM, *et al.* Successful treatment of *Fusarium* keratitis with cornea transplantation and topical and systemic voriconazole. *Clin Infect Dis* 2005;**40**:e110-12.
- 3 Nulens E, Eggink C, Rijs AJMM, *et al.* Keratitis caused by *Scedosporium apiospermum* successfully treated with a cornea transplant and voriconazole. *J Clin Microbiol* 2003;**42**:2261-4.
- 4 Ozbek Z, Kang S, Sivalingam J, *et al.* Voriconazole in the management of *Alternaria* keratitis. *Cornea* 2006;**25**:242-4.
- 5 Zinkernagel A, Scherrer M, Burhenne J, *et al.* Outcome and aqueous drug levels of voriconazole in four patients with fungal keratitis treated with voriconazole. *Clin Microbiol Infect* 2004;**10**(suppl3):508-9.
- 6 Lalitha P, Prajna NV, Kabra A, *et al.* Risk factors for treatment outcome in fungal keratitis. *Ophthalmology* 2006;**113**:526-30.
- 7 Marangon FB, Miller D, Giacino JA, *et al.* In vitro investigation of voriconazole susceptibility for keratitis and endophthalmitis fungal pathogens. *Am J Ophthalmol* 2004;**137**:820-5.
- 8 Hariprasad SM, Mieler WF, Holz ER, *et al.* Determination of vitreous, aqueous, and plasma concentration of orally administered voriconazole in humans. *Arch Ophthalmol* 2004;**122**:42-7.
- 9 Sponkel W, Chen N, Dang D, *et al.* Topical voriconazole as a novel treatment for fungal keratitis. *Antimicrob Agents Chemother* 2006;**50**:262-8.
- 10 Muijsers RB, Goa KL, Scott U. Voriconazole: in the treatment of invasive aspergillosis. *Drugs* 2002;**62**:2655-64.

12 weeks, respectively) than others (treatment range 6 weeks to 6 months). Whether these patients had a higher dose of isotretinoin or had any predisposing retinal problems are not adequately explained in the report. It would have been informative if the authors had compared the dose-effect relationships among patients in whom the dose of treatment was known (8 patients with abnormal ERG and 23 patients with normal ERG in the isotretinoin group).

In this particular study, the authors have mentioned a patient who continued to show signs of retinal toxicity 8 years after cessation of treatment, presumably with changes in ERG, but there is no description of this patient either in table 1 or elsewhere in the article. Further, the authors have compared the persistence of retinal toxicity in this particular patient with the study conducted by Oner *et al.*² In their study, Oner *et al.* have looked into the visual acuity, anterior segment changes, intraocular pressure, Schirmer's test, tear film break-up time, colour vision and changes in microbial flora. They specifically mention in their article that they did not perform any electrodiagnostic studies in their patients. Perhaps it is inappropriate to compare the two studies, which have looked at entirely different aspects of side effects of isotretinoin.

Although Mollan *et al.* have mentioned in their methods that colour vision was tested in their patients, they have not elaborated on the relevant results in the article.

It is interesting to note that the authors have not justified in the main report their recommendation for routine electrophysiological screening for professions that are critical for night vision, except in the abstract. It would be appropriate to conduct a prospective study to look into the electrodiagnostic and other described ocular changes following the use of isotretinoin, to precisely address their question.

S Pushpoth, S Sandramouli

Wolverhampton and Midland Counties Eye Infirmary,
Wolverhampton, UK

Correspondence to: Dr S Pushpoth, Wolverhampton
and Midland Counties Eye Infirmary, Compton Road,
Wolverhampton WV3 9QR, UK; drsree@tiscali.co.uk

Accepted 24 October 2006

Competing interests: None declared.

References

- 1 Mollan SP, Woodcock M, Siddiqi R, *et al.* Does use of isotretinoin rule out a career in flying? *Br J Ophthalmol* 2006;**90**:957-9.
- 2 Oner A, Ferahbas A, Karakucuk S, *et al.* Ocular side-effects associated with systemic isotretinoin. *J Toxicol* 2004;**23**:189-95.

Isotretinoin and night vision: authors' response

We thank Pushpoth and Sandramouli¹ for their wide-ranging critique of our paper discussing previous isotretinoin use in potential military aviator recruits.² We cannot agree with their comment that the finding of 2 of 47 candidates, with clinically abnormal dark adaptation (DA) and 11 with electroretinogram (ERG) abnormalities does not justify further screening of this population. Aircrew recruits are a selected population, to a large degree, without any significant history of systemic or ocular abnormalities. In balance of probability, this selection makes it more likely that the retinal function of these individuals falls within the

age-matched normal population, not less. The candidates had, for the most part, applied for a career in aviation long after taking isotretinoin, although we agree that it would have been desirable to prospectively follow the ERG and DA changes before and after treatment.

Pushpoth's and Sandramouli's assumption that isotretinoin-related retinal toxicity is dose related cannot be made from our retrospective report, largely for the reasons outlined in their comments on incomplete dosage information. We do know that none of the individuals had a history, or a family history, of retinal abnormality and that clinical retinal and visual field measurements were normal. Colour vision tested normal in all cases with an Ishihara Pseudoisochromatic Plate Test and Lanthony 15-hue colour-vision test.

Our reference to Oner *et al.*² was not to compare the paper's results, but to disagree with the finding that all side effects of isotretinoin were short lived. Had they performed electrophysiology they might have found a different result.

In this paper, we lend support to the findings of other studies that rod function can be adversely affected by previous isotretinoin use.^{4,5} As aviation is a night-vision critical profession, we consider it justified at present to check the night vision of aircrew candidates with a history of isotretinoin by dark adaptometry, from a flight-safety point of view. We agree with Pushpoth and Sandramouli that a prospective study of the retinal effects of isotretinoin would shed more light on the problem, as much remains to be learnt about the safety of isotretinoin.

Susan P Mollan, Malcolm Woodcock,
Peter Good, Robert A H Scott

Birmingham Midland Eye Centre, Birmingham, UK

Correspondence to: Ms S P Mollan, Birmingham
Midland Eye Centre, 64 B Central, Birmingham B1 2AF,
UK; soozmollan@doctors.org.uk

Accepted 9 November 2006

Competing interests: None declared.

References

- 1 Pushpoth S, Sandramouli S. Isotretinoin and night vision. *Br J Ophthalmol* 2007;**91**:1096.
- 2 Mollan SP, Woodcock M, Siddiqi R, *et al.* Does use of isotretinoin rule out a career in flying? *Br J Ophthalmol* 2006;**90**:957-9.
- 3 Oner A, Ferahbas A, Karakucuk S, *et al.* Ocular side-effects associated with systemic isotretinoin. *J Toxicol* 2004;**23**:189-95.
- 4 Brown RD, Grattan CEH. Visual toxicity of synthetic retinoids. *Br J Ophthalmol* 1989;**73**:286-8.
- 5 Weleber RG, Denman ST, Hanifin JM, *et al.* Abnormal retinal function associated with isotretinoin therapy for acne. *Arch Ophthalmol* 1986;**104**:831-7.

Surgical embolus removal in retinal artery occlusion

I was interested to read the article by Garcia-Arumi and colleagues on "Surgical embolus removal in retinal artery occlusion".¹ The authors claim that "Surgical removal of retinal arterial emboli seems to be an effective and safe treatment for RAO (retinal artery occlusion)".

Briefly, the study was based on six eyes with temporal branch retinal artery occlusion (BRAO) and one with central retinal artery occlusion (CRAO). The surgery was performed in the eyes with BRAO 4, 12, 19, 22, 28, and 33 h after onset and in the eye with CRAO 29 h after visual loss. The first postoperative evaluation, 48 h after surgery, showed reperfusion of the occluded branch retinal artery in four and none in one; in

MAILBOX

Isotretinoin and night vision

We read with interest the article by Mollan *et al.*¹ who have concluded that previous isotretinoin use does not cause a clinically significant reduction in night vision in most people, and that the retinal toxic effects of isotretinoin may be measurable by electroretinography (ERG) and dark adaptation (DA). Although the authors have successfully highlighted the importance of counselling patients for potential irreversible loss of DA following isotretinoin use, their report, in our opinion, has failed to substantiate the need for routine screening of potential military and civilian commercial aviators.

In their study, 2 of 47 patients had both abnormal ERG and DA, whereas 11 others had certain abnormal ERG parameters that may or may not be of practical significance. The interpretation of this finding is debatable in the context of this study being a retrospective analysis, where we cannot assess the electrodiagnostic status of the patients in the study group before treatment. Only two patients in the study had abnormalities in both ERG and DA. Those two patients, X and Y, received treatment for a comparatively shorter period of time (8 and