We recommend that the authors accept the possibility for type II error in their results and conclusion.

C H Lim, V Ramappa, M M Ahmed

Department of Gastroenterology, Good Hope Hospital NHS Trust, Sutton Coldfield, West Midlands, B75 7RR, UK

Correspondence to: Dr C H Lim, Department of Gastroenterology, Good Hope Hospital NHS Trust, Rectory Road, Sutton Coldfield, W Midlands, B75 7RR, UK; chee.lim@goodhope.nhs.uk

Competing interests: None.

Reference

- 1 Moher D, Schulz KF, Altman D. The CONSORT statement: revised recommendations for improving the quality of reports of parallel-group randomized trials. *JAMA* 2001;**285**:1987–91.

Authors' response

We thank Lim *et al* for their keen interest in our article (*Gut* 2006; **55**: 1233–9) on the long-term results of ablation in patients with Barrett's oesophagus (BO). In this prospective, randomised controlled trial, we compared the long-term efficacy of achieving complete reversal (endoscopic and histological) between multipolar electrocoagulation (MPEC) and argon plasma coagulation (APC) in patients with BO. Patients underwent stratified randomisation by BO length, using sealed opaque envelopes. Results of this study showed that complete reversal can be maintained in ~70% of patients, irrespective of the technique, with no predictors identified associated with achieving complete reversal.

In this study, we used stratified randomisation instead of simple randomisation, which allowed us to use the paired Fisher's test for determination and analyses of power and sample size. The paired Fisher's test finds whether there is a difference between two groups when the outcome is a response proportion and the difference is assessed by testing if the odds ratio (OR) is 1 or not. This works only when the subjects are stratified into groups before randomisation, as it is believed that the response proportions will vary across the strata, but that the OR is constant across the strata. In the case of our study, the strata are the BO lengths.¹ We believe that Lim *et al* have assumed a simple randomisation and hence used a non-paired Fisher's exact test for their sample size calculation.

Crucial demographic and endoscopic features of patients and the number of ablation sessions required in the two groups were reported in this study. We agree with Lim *et al* that information such as the duration of each procedure for the two techniques, information on patients with BO who refused participation, and so on was not provided in the manuscript. However, as we have highlighted in our article, we reiterate that techniques such as MPEC and APC are not ready for clinical application in ablation of non-dysplastic BO and should not be offered outside the research arena. The degree of long-term control of neoplasia is not as yet known. Persistence of underlying intestinal metaplasia after ablation (buried glands) carries the potential to progress to dysplasia and adenocarcinoma, a problem that cannot be ignored.² It has also been shown that recurrent and/or persistent BO after ablative therapy still contains genetic

alterations associated with malignant progression to cancer.³ At the present time, the available data do not support the application of ablative therapies for patients with non-dysplastic BO.

Matthew Hall, Sachin Wani, Prateek Sharma
Veterans Affairs Medical Center & University of Kansas School of Medicine, Kansas City, Missouri, USA

Correspondence to: Dr P Sharma, Division of Gastroenterology, Veterans Affairs Medical Center, 4801 E Linwood Blvd, Kansas City, MO 64128-2295, USA; psharma@kumc.edu

Competing interests: None.

References

- 1 Nam J. Sample size determination for case-control studies and the comparison of stratified and unstratified analyses. *Biometrics* 1992;**48**:389–95.
- 2 Van Laethem JL, Peny MO, Salmon I, *et al*. Intramucosal adenocarcinoma arising under squamous re-epithelialisation of Barrett's oesophagus. *Gut* 2000;**46**:574–7.
- 3 Hage M, Siersema PD, Vissers KJ, *et al*. Genomic analysis of Barrett's esophagus after ablative therapy: persistence of genetic alterations at tumor suppressor loci. *Int J Cancer* 2006;**118**:155–60.

Development of a bioartificial new intestinal segment using an acellular matrix scaffold

Intestinal rehabilitation for short-bowel syndrome is an integral part of modern intestinal transplant programmes. The mortality of patients with short-bowel syndrome is most significant in individuals with a residual small bowel of <50 cm, as shown by a 5-year survival rate of 57%.¹ Total parenteral nutrition and intestinal transplantation are options to

extend life but are still plagued by serious complications and, in the case of transplantation, immunosuppression. As an alternative, several bowel elongation procedures have been described,^{2–4} but have had limited clinical success and new techniques are warranted. The minimum length of bowel required to allow sufficient absorption of nutrients has not been confirmed.^{1,5} Elongation of even a few centimetres may allow these patients to receive nutritional rehabilitation and become independent from total parenteral nutrition, and possibly avoid transplantation. We hypothesised that an acellular dermal matrix (ADM, AlloDerm, LifeCell Corporation, Branchburg, New Jersey, USA) scaffold placed in continuity with defunctionalised jejunal limb allows mucosal growth and intestinal elongation. We evaluated the morphology of neofomed intestine in ACI (August × Copenhagen-Irish) rats at different time points using two different types of anastomosis. Tubular scaffolds with an intraluminal diameter of approximately 0.3 cm were constructed using rehydrated ADM segments of 1 cm² and 0.78–1.77 mm thickness, and oriented with a luminal basement membrane and a serosal dermal surface. In group A (n = 5), the ADM graft was interposed in continuity with the jejunum using an interrupted end-to-end single-layer anastomosis. In group B (n = 11), the grafts were placed as blind-ended pouches to the defunctionalised jejunal limb. Postoperatively, animals were maintained on a liquid diet for 48 h followed by solid-rat chow and killed at different time-points postoperatively. Tissue samples for histological examination were obtained across the anastomosis. Survival and body weight were evaluated in both groups. All animals in group A were killed in the first week as a result of peritonitis. All animals in group B survived

Figure 1 Photomicrographs of the anastomosis between the acellular dermal matrix (ADM) and small bowel at 4 months of transplantation. (A) Leading edge of regenerating epithelium with villus formation (V). Note a bit of remaining bare graft surface to the left of the villus, H&E ×20. (B) Regenerating crypts with goblet cells. Some residual collagen from the ADM can be seen nearby (arrow), H&E ×20. (C) Epithelium along the edge of the villus in (A) showing goblet cells (arrows) and absorptive cells with brush border, H&E ×60. (D) Vascularisation of the ADM just beneath the regenerating crypts, H&E ×40.

Table 1 Histology results of group B at different time points

Time after anastomosis	Status of anastomosis	Status of alloderm	Epithelial regeneration
2 days	Intact	Minimal acute inflammation	No
2 weeks	Intact but inflamed	Minimal acute inflammation. Proliferation of fibroblasts and endothelial cells	Early budding of crypts at bowel edge of anastomosis
4–5 weeks	Intact but inflamed	Less acute inflammation. Full-thickness ingrowths of capillaries and myofibroblasts	Same as 2 weeks
4–6 months	Intact with granulation tissue around sutures	Full-thickness capillaries and fibroblasts and haemosiderin-laden macrophages	Regenerating crypts with goblet cells, and rudimentary villi with absorptive cells with brush border

and increased body weight appropriately. Tissue samples showed a progressive increase in the amount of cell infiltrate in the matrix (table 1, fig 1). After 2 weeks, acute inflammation was replaced by full-thickness ingrowths of capillaries and myofibroblasts. Epithelial regeneration into the anastomosis was first seen at 2 weeks, and well-formed branching crypts were seen at 4 weeks of transplantation. Goblet cells and absorptive cells with brush border were present at 4 months of transplantation. Morphologically intact regenerated mucosa extending across the anastomosis to the grafts was observed at 6 months of transplantation. To date, there is limited literature on the bioengineered intestine. Vacanti *et al.*^{6–8} developed a cystic structure in which neomucosa forms in a biodegradable polymer in rodents. Once formed, the neointestinal cyst is anastomosed in continuity with the native bowel without causing feeding problems, but some animals had small bare areas of the cysts that lacked neomucosa.⁷ We did not report such bare areas in our model; furthermore, there was progressive growth of the neomucosa in the ADM over time. It is possible that immediate contact of the ADM scaffold with the intestinal structures and with luminal content provided trophic stimuli for the new intestinal segment.⁹ Another possible factor for the observed growth in our model may be the effects of small-bowel resection on the development of neointestine. It is well known that post resection gut mucosa growth factors have a stimulatory effect on intestinal regeneration.¹⁰ In conclusion, we have demonstrated that ADM can be successfully used as a scaffold to generate a bioartificial new intestinal segment in vivo, and we propose this method as a basis for developing new intestinal elongation techniques.

Mohan P Pahari, Melissa L Brown, Georg Elias, Hannan Nseir

Department of Surgery, Division of Organ Transplantation, University of Massachusetts, Medical School, UMass/Memorial Healthcare Worcester, Massachusetts, USA

Barbara Banner

Department of Pathology, University of Massachusetts, Medical School, UMass/Memorial Healthcare Worcester, Massachusetts, USA

Cristiana Rastellini, Luca Cicalese

Department of Surgery, Division of Organ Transplantation, University of Massachusetts, Medical School, UMass/Memorial Healthcare Worcester, Massachusetts, USA

Correspondence to: Dr Luca Cicalese, Department of Surgery, Division of Organ Transplantation, University of Massachusetts Medical School, 55 Lake Avenue North, S3-844, Worcester, MA 01655, USA; luca_cicalese@yahoo.com

doi: 10.1136/gut.2006.116848

Competing interests: None.

References

- Messing B, Crenn P, Beau P, *et al.* Long-term survival and parenteral nutrition dependence in adult patients with the short bowel syndrome. *Gastroenterology* 1999;117:1043–50.
- Bianchi A. Intestinal loop lengthening: a technique for increasing small intestine length. *J Pediatr Surg* 1980;15:145–51.
- Kim HB, Fauza D, Garza J, *et al.* Serial transverse enteroplasty (STEP): a novel bowel lengthening procedure. *J Pediatr Surg* 2003;38:425–9.
- Kimura K, Soper RT. A new bowel elongation technique for short-bowel syndrome using the isolated bowel segment Iowa models. *J Pediatr Surg* 1993;28:792–4.
- Nightingale JM, Lennard-Jones J E, Gertner D J, *et al.* Colonic preservation reduces need for parenteral therapy, increases incidence of renal stones, but does not change high prevalence of gall stones in patients with a short bowel. *Gut* 1992;33:1493–7.
- Choi RS, Vacanti JP. Preliminary studies of tissue-engineered intestine using isolated epithelial organoid units on tubular synthetic biodegradable scaffolds. *Transplant Proc* 1997;29:848–51.
- Kim SS, Kaihara S, Benvenuto M, *et al.* Regenerative signals for tissue-engineered small intestine. *Transplant Proc* 1999;31:657–60.
- Kaihara S, Kim SS, Kim BS, *et al.* Long-term follow-up of tissue-engineered intestine after anastomosis to native small bowel. *Transplantation* 2000;69:1927–32.
- Thompson JS, Tempero MA, Haun JL, *et al.* The importance of luminal factors in neomucosal growth. *J Surg Res* 1986;40, 126–32.
- Cisler JJ, Buchman AL. Intestinal adaptation in short bowel syndrome. *J Invest Med* 2005;53:402–13.

No evidence of persisting measles virus in the intestinal tissues of patients with inflammatory bowel disease

The hypothesis that measles virus (MV) is implicated in the pathogenesis of Crohn's disease (CD) was initiated by Wakefield *et al*¹ in the early 1990s. Several other reports soon followed, which suggested the presence of MV in CD samples using electron microscopy,

immunohistochemical analysis, in situ hybridisation and nested PCR.^{2–4} Controversy in response to this claim resulted in falling measles vaccination rates, mainly in the UK. Immediate concerns were raised regarding several aspects of these studies, including reagent specificity.⁵ In addition, several groups were unable to replicate their findings using a range of PCR techniques.^{6–9}

In 1998 and 2002, the initial investigators again raised concerns over MV persistence in bowel tissue, this time in children with autism.¹⁰ These allegations further damaged confidence in MV vaccination programmes. As the primer pairs that had been used to implicate MV in inflammatory bowel disease (IBD; Kawashima)⁴ and autism (Uhlmann)¹⁰ were not included in the negative studies,^{6–9} it could still be argued that the negative results arose due to the use of incorrect or suboptimal primers. We therefore used the Kawashima and Uhlmann primers in an attempt to detect MV nucleic acids in patients with IBD and control intestinal biopsy samples. Three real-time and two nested PCR assays targeting the MV nucleocapsid, fusion and haemagglutinin genes using the Kawashima and Uhlmann primers were performed, with detection by SYBR Green I. We also designed our own probe-based F gene assay. Positive samples were analysed by melting curve analysis, electrophoresis and sequencing. A total of 41 biopsy samples were available for this study (table 1). Full details of the primers and methods were published previously.¹¹

We report that all samples yielded positive signals in the Uhlmann real-time PCR assays (fig 1A). Our study generated mean (SD) copy numbers that were very similar to those reported by Uhlmann *et al*—that is, between 5.08E3 ($\pm$ 6.32E3) and 2.94E5 ($\pm$ 5.75E5). However, only a proportion of the PCR products had melting peaks that overlapped with those of the positive control sample (fig 1A,B). Of these positive results, no sample in the Uhlmann fusion or haemagglutinin gene assays yielded an appropriate 150 bp band when analysed by electrophoresis. Only 2 of 17 CD samples produced appropriately-sized bands (150 bp) using the nucleocapsid gene assay. Repeated attempts to clone and sequence were successful for one of the two nucleocapsid gene amplicons. This amplicon had sequence homology to various mammalian genes, including “*Homo sapiens* cDNA FLJ34053 fis, highly similar to histone H1” (National Center for Biotechnology Information (NCBI) accession #AK091372).

In all, 10 samples (4 CD, 1 ulcerative colitis and 5 non-IBD) had the correct band size (181 bp) after amplification using the Kawashima fusion gene assay. Nine of ten amplicons successfully cloned and sequenced had homology to “*Homo sapiens* mitochondrial DNA gene” (NCBI accession #AY289053). The remaining amplicon did not match any genes from the NCBI database. No sample was positive by nested PCR using the Kawashima haemagglutinin gene primers or our own probe-based fusion gene assay.

Both the Kawashima and Uhlmann primers demonstrated significant cross-reactivity. Searches for sequence similarities revealed homology from 43% to 75% between the mammalian sequences that we amplified and the Kawashima and Uhlmann primers.

In summary, by direct application of the Uhlmann and Kawashima primers, we also failed to demonstrate the presence of MV nucleic