Milena Bakrac

Institute of Hematology, Clinical Center of Serbia,
Belgrade, Serbia

Vladimir Jurisic

Institute for Pathophysiology, School of Medicine,
University of Kragujevac, Kragujevac, Serbia

Tanja Kostic

Institute for Molecular Genetics and Genetic
Engineering, Belgrade, Serbia

Vera Popovic, Sandra Pekic

Institute of Endocrinology, Clinical Center of Serbia,
Belgrade, Serbia

Nada Kraguljac, Milica Colovic

Institute of Hematology, Clinical Center of Serbia,
Belgrade, Serbia

Correspondence to: Professor V Jurisic, University of
Kragujevac, School of Medicine, PO Box 124, 34 000
Kragujevac, Serbia

doi: 10.1136/jcp.2006.042671

Accepted 17 November 2006

Published Online First 12 January 2007

Competing interests: None declared.

References

- Handgretinger R, Geiselhart A, Moris A, *et al.* Pure red cell aplasia associated with clonal expansion of granular lymphocytes expressing killer-cell inhibitory receptors. *N Engl J Med* 1999;340:278–84.
- Charles RJ, Sabo KM, Kidd PG, *et al.* The pathophysiology of pure red cell aplasia: implications for therapy. *Blood* 1996;87:4831–8.
- Casadevall N, Dupry E, Mohlo-Sabatier P, *et al.* Autoantibodies against erythropoietin in a patient with pure red cell aplasia. *N Engl J Med* 1996;334:630–3.
- Hara T, Mizuno Y, Nagata M, *et al.* Human $\gamma\delta$ T-cell receptor positive cell-mediated inhibition of erythropoiesis in vitro in a patient with type I autoimmune polyglandular syndrome and pure red cell aplasia. *Blood* 1990;75:941–50.
- Ahonen P, Myllarniemi S, Sipila I, *et al.* Clinical variation of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) in a series of 68 patients. *N Engl J Med* 1990;322:1829–36.
- Betterle C, Greggio NA, Volpato M. Autoimmune polyglandular syndrome type 1. *J Clin Endocrinol Metab* 1998;83:1049–55.
- Myhre AG, Halonen M, Eskelin P, *et al.* Autoimmune polyendocrine syndrome type 1 (APS I) in Norway. *Clin Endocrinol* 2001;54:211–17.
- Nagamine K, Peterson P, Scott HS, *et al.* Positional cloning of the APECED gene. *Nat Genet* 1997;17:393–7.
- Ward L, Paquette J, Seidman E, *et al.* Severe autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy in an adolescent girl with a novel AIRE mutation: response to immunosuppressive therapy. *J Clin Endocrinol Metab* 1999;84:844–52.
- Phillips JH, Gumperz JE, Partham P, *et al.* Superantigen-dependent, cell mediated cytotoxicity inhibited by MHC class I receptors on T lymphocytes. *Science* 1995;268:403–5.
- Engelen W, Verpoeten GA, Van der Planken M, *et al.* Four cases of red blood cell aplasia in association with the use of mycophenolate mofetil in renal transplant patients. *Clin Nephrol* 2003;60:119–24.
- Allison AC, Eugui EM. Mechanisms of action of mycophenolate mofetil in preventing acute and chronic allograft rejection. *Transplantation* 2005;80:181–90.
- Nakamura M, Ogawa N, Shalabi A, *et al.* Positive effect on T-cell regulatory apoptosis by mycophenolate mofetil. *Clin Transplant* 2001;15:36–40.
- Rao VK, Dugan F, Dale JK, *et al.* Use of mycophenolate mofetil for chronic, refractory immune cytopenias in children with autoimmune lymphoproliferative syndrome. *Br J Haematol* 2005;129:534–8.
- Andrikos E, Yavuz A, Bordon V, *et al.* Effect of cyclosporine, mycophenolate mofetil, and their combination with steroids on apoptosis in a human cultured monocytic U937 cell line. *Transplan Proc* 2005;37:3226–9.
- Marazuela M, Garcia-Lopez MA, Fuguerola-Vega N, *et al.* Regulatory T cells in human autoimmune disease. *J Clin Endocrinol Metab* 2006;91:3639–46.
- Arbeiter K, Greenbaum L, Balzar E, *et al.* Reproducible erythroid aplasia caused by mycophenolate-mofetil. *Pediatr Nephrol* 2000;14:195–7.

Metastatic renal oncocytoma

Renal oncocytomas are classified as benign renal neoplasms in the 2004 *World Health Organization classification of renal tumours*.¹ Cases of metastatic tumours have been documented,^{2–4} but subsequent reports have identified subtypes of renal tumours having similar histological morphology but having malignant potential. These tumours have been classified as either eosinophilic variants of renal cell carcinoma (RCC) or chromophobe carcinomas. Differentiating these tumours from oncocytomas has proved to be difficult and relies on a combination of histological morphology, immunohistochemistry and electron microscopy. There is still a presumption that if metastasis occurs then the tumour should not be classified as a renal oncocytoma and should be called an eosinophilic RCC.

We present a patient who presented with liver metastasis 9 years after the removal of a renal oncocytoma.

Case report

A woman in her 70s presented with left-sided abdominal pain. She was noted to have a mobile mass in the left upper quadrant. Baseline blood examinations were normal and she had no other significant medical history. An abdominal ultrasound was suggestive of a left RCC. A CT scan showed a 12 cm renal mass extending just across the midline, arising from the upper anterior part of the left kidney (fig 1). The left renal vein, liver and lymph nodes appeared unaffected. The conclusion was that this was most probably an

RCC, but the pattern of enhancement was suggestive of a renal oncocytoma. A chest x ray was clear.

She underwent a left radical nephrectomy and adrenalectomy en bloc. There was no involvement of adjacent organs. She made an uneventful recovery. She remained well on follow-up, with normal renal function and full blood counts. She was discharged from follow-up 18 months after surgery.

After 9 years she re-presented with left-sided abdominal pain, bloating and nausea, but no vomiting. A chest x ray was normal and an abdominal x ray showed faecal loading; this was thought to be diverticular disease. An abdominal CT scan showed multiple liver masses.

An ultrasound-guided liver biopsy was performed.

A CT scan 4 months later showed multiple lesions throughout the liver, which enhanced in the arterial phase, consistent with multiple hypervascular metastases (fig 1). The tumours appeared much the same as on the previous scan. Her liver function tests were normal. She had few symptoms, consisting of some tiredness, minor weight loss but a good appetite until 18 months after she re-presented when she developed confusion. A CT scan of her head showed acute left parietotemporal haemorrhage with intraventricular extension, but there was no evidence of metastatic tumour. She remained an inpatient for 4 months during which her renal and liver function tests were normal. She was discharged to a nursing home where she died a week later. A post-mortem examination was not carried out.

Pathology

The renal tumour was in the upper and mid-zones of the kidney and measured up to 12 cm in diameter. The tumour was yellow-brown in colour with a central white stellate scar, 1.5 cm in diameter. The perinephric fat, renal vein and adrenal gland were uninvolved. Histology showed packets of regular cells with eosinophilic cytoplasm and occasional highly cellular trabecular areas (fig 2). There was no cellular atypia, and mitotic figures were not seen (0 per 1 mm²). MIB-1 labelling index was <1%. Hales colloidal iron was negative. The conclusion was that this was a renal oncocytoma.

The liver biopsy specimen showed cores of liver parenchyma with well defined islands of oncocytic cells arranged in nests (fig 3). The original renal tumour was reviewed in light of

Figure 1 Enhanced CT scans showing (A) a 12 cm well-defined left renal mass with a central scar and radiating area of low density suggestive of oncocytoma and (B) the liver metastases that appeared 9 years later.

Figure 2 Renal oncocytoma showing high and low cellularity: (A) H&E $\times 16$; (B) H&E $\times 630$; (C) H&E $\times 630$; (D) epithelial membrane antigen immunohistochemistry $\times 100$, highlighting different immunopatterns in different areas; (E) electron microscopy showing numerous mitochondria $\times 7900$.

Figure 3 Metastatic renal oncocytoma in liver: (A) H&E $\times 400$, (B) epithelial membrane antigen immunohistochemistry $\times 200$.

the patient presenting with metastatic disease, and further immunohistochemistry was performed both by our laboratory and by Professor Susani in Vienna.

The renal tumour expressed vimentin and pan-cytokeratin marker (AE1/AE3), and there was focal positivity for epithelial membrane antigen (EMA; fig 2) and Ber EP4. CD10 showed very weak patchy positivity and the tumour was negative for c-kit (CD117) and cytokeratin 7. Limited tissue was present in the liver cores, but the tumour was also AE1/AE3 and EMA positive (fig 4). Professor Susani performed immunohistochemical analysis for the novel marker kidney-specific cadherin (Ksp cadherin), and both the renal tumour and the metastasis showed $<10\%$ marking, which is consistent with a renal oncocytoma.⁵

Electron microscopy on the renal tumour showed cells containing large numbers of mitochondria (fig 2), a finding characteristic of oncocytomas. By contrast, ultrastructural examination of chromophobe carcinomas would typically reveal numerous vesicles. Liver biopsy tissue was insufficient to perform electron microscopy.

Discussion

There have been extensive reports on the biological behaviour and histological characteristics of renal oncocytomas since they were first described over 50 years ago.^{3,4} Since then, only rare cases of metastatic oncocytomas have been described and none has been studied with the modern panel of immunohistochemistry utilised in this case report.

Carcinomas and benign epithelial tumours are characterised by the expression of cytokeratins. Currently, 20 distinct subtypes of cytokeratins are known, and the expression profile of these can help classify tumours. However, there is often a degree of overlap between tumours—for example, Skinnider *et al*⁶ found that 10% of renal oncocytomas expressed cytokeratin 7 as opposed to 73% of chromophobe carcinomas. Another marker studied is c-kit, which encodes the membrane-bound tyrosine kinase KIT, and in a recent study was found to be expressed by both chromophobe and oncocytomas, but was not detected in most other subtypes.⁷

Pan *et al*⁸ studied a variety of markers and found EMA to be expressed similarly by oncocytomas and chromophobe carcinomas, and BerEP4 to be negative in all but one renal oncocytoma while positive in all but two chromophobe carcinomas. CD10 was negative in all the renal oncocytomas they tested, but was negative in 68% of the chromophobe carcinomas. Vimentin was negative in all the oncocytomas and in 79% of the chromophobe carcinomas. This study was limited as only 7 oncocytomas and 28 chromophobe carcinomas were examined.

A larger study was carried out by Mazal *et al*⁵ examined the novel marker kidney-specific cadherin. This is a cell adhesion molecule found on the basolateral membrane of renal tubular epithelial cells and collecting duct cells. This study involved a variety of renal tumours but included 31 oncocytomas and 30 chromophobe tumours. The conclusion was that kidney-specific cadherin was expressed almost exclusively in chromophobe tumours.

The limitation of these immunohistochemical studies is that the tumours have to be classified on the basis of morphology before the immunopanel is performed, but there is

often disagreement among pathologists when chromophobe carcinomas are examined.⁹

Adequate sampling of renal tumours is essential to exclude any heterogeneity, and in this case eight blocks from the renal tumour were examined. This may mean that an atypical area was missed but the metastatic tumour had the same morphology and immunoprofile as the tumour areas sampled, suggesting that this was not the case. DNA microsatellite analysis could be used to confirm this, but liver biopsy tissue was insufficient for this to be performed.

With an ever-expanding repertoire of immunohistochemical markers available, these have been applied to renal tumours in order to differentiate renal oncocytomas from malignant renal neoplasms. Examining the tumours at the ultrastructural level and looking for numerous mitochondria still remains the key to the diagnosis of renal oncocytomas. This case highlights that using these techniques can help classify but cannot predict malignant potential. With this in mind, renal oncocytomas should be considered as having a very low rather than no malignant potential.

Acknowledgements

We thank Professor Martin Susani (Professor of Pathology, Department of Clinical Pathology, AKH-Wien, Vienna, Austria), and his staff for reviewing and performing immunohistochemistry on this case.

J D Oxley

Department of Cellular Pathology, North Bristol NHS Trust, Southmead Hospital, Westbury on Trym, Bristol, UK

J Sullivan

Department of Urology, North Bristol NHS Trust, Southmead Hospital, Westbury on Trym, Bristol, UK

A Mitchelmore

Department of Radiology, North Bristol NHS Trust, Southmead Hospital, Westbury on Trym, Bristol, UK

D A Gillatt

Department of Urology, North Bristol NHS Trust, Southmead Hospital, Westbury on Trym, Bristol, UK

Correspondence to: J D Oxley, Department of Cellular Pathology, North Bristol NHS Trust, Southmead Hospital, Westbury on Trym, Bristol BS10 5NB, UK; jon.oxley@nbt.nhs.uk

doi: 10.1136/jcp.2006.044198

Accepted 17 October 2006

Competing interests: None declared.

References

- 1 Eble JN, Sauter G, Epstein JI, Sesterhenn IA. *World Health Organization classification of tumours. Pathology and genetics of tumours of the urinary system and male genital organs*. Lyon: IARC Press, 2004.
- 2 Amin R, Anthony P. Metastatic renal oncocytoma: a case report and review of the literature. *Clin Oncol (R Coll Radiol)* 1999;11:277-9.
- 3 Amin MB, Crotty TB, Tickoo SK, et al. Renal oncocytoma: a reappraisal of morphologic features with clinicopathologic findings in 80 cases. *Am J Surg Pathol* 1997;21:1-12.
- 4 Perez-Ordóñez B, Hamed G, Campbell S, et al. Renal oncocytoma: a clinicopathologic study of 70 cases. *Am J Surg Pathol* 1997;21:871-83.
- 5 Mazal PR, Exner M, Haitel A, et al. Expression of kidney-specific cadherin distinguishes chromophobe renal cell carcinoma from renal oncocytoma. *Hum Pathol* 2005;36:22-8.
- 6 Skinnider BF, Folpe AL, Hennigar RA, et al. Distribution of cytokeratins and vimentin in adult

renal neoplasms and normal renal tissue: potential utility of a cytokeratin antibody panel in the differential diagnosis of renal tumors. *Am J Surg Pathol* 2005;29:747-54.

- 7 Huo L, Sugimura J, Tretiakova MS, et al. C-kit expression in renal oncocytomas and chromophobe renal cell carcinomas. *Hum Pathol* 2005;36:262-8.
- 8 Pan CC, Chen PC, Ho DM. The diagnostic utility of MOC31, BerEP4, RCC marker and CD10 in the classification of renal cell carcinoma and renal oncocytoma: an immunohistochemical analysis of 328 cases. *Histopathology* 2004;45:452-9.
- 9 Abrahams NA, MacLennan GT, Khoury JD, et al. Chromophobe renal cell carcinoma: a comparative study of histological, immunohistochemical and ultrastructural features using high throughput tissue microarray. *Histopathology* 2004;45:593-602.

Multiple adenomatoid tumours in the liver and peritoneum

Adenomatoid tumours are uncommon tumours, which were first described by Golden and Ash.¹ They occur most often in the male and female genital tracts, but have rarely been reported at other sites such as the omentum,² pleura,³ heart,⁴ small bowel mesentery⁵ and adrenal gland.⁶

We report multiple adenomatoid tumours involving the peritoneum and liver. Our diagnosis was based on morphological examination, and was supported by histochemical, immunohistochemical and ultrastructural examinations.

Materials and methods

Tissue was processed for frozen sections using a cryostat, and sections were stained with H&E.

Formalin-fixed paraffin wax sections of each tumour were stained with periodic acid-Schiff after diastase predigestion, in addition to H&E staining. Immunohistochemical staining for cytokeratins (CK) 5, 6, 8, 17 and 19 (clone MNF116, DAKO cytometry), epithelial membrane antigen (EMA; clone E29, DAKO cytometry), epithelial antigen (clone BERE4, DAKO cytometry), calretinin (clone DAK-calret DAKO cytometry), mesothelial cell (clone human bone marrow endothelial (HBME)-1 DAKO cytometry), CD31 (DAKO) and CD34 (DAKO) was performed using an automated immunostainer (DAKO Autostainer, Ely, UK) and a labelled streptavidin-biotin method.

Electron microscopy was performed on tissue retrieved from paraffin-wax-embedded blocks, post-fixed in osmium tetroxide, and stained with uranyl acetate and lead citrate. Microscopy was performed on 80-90 nm sections using a Hitachi H600 transmission electron microscope (Hitachi, Wokingham, UK).

Results

A woman in her 70s presented with coffee ground vomiting associated with malaena. She also had a history of *Helicobacter pylori* infection and eradication therapy, glaucoma and hypertension. There was no history of previous surgery. Her medication consisted of a proton pump inhibitor, a β -blocker and eye drops for glaucoma. There was no relevant family history.

An oesophago-gastro-duodenoscopy showed a chronic ulcer involving the lesser curve of the stomach and a positive campylobacter-like organism test. A biopsy specimen was taken, which showed the features of a gastric adenocarcinoma. A staging CT scan of the thorax and

abdomen showed a T2/T3 lesion, with no radiological evidence of local invasion or metastasis.

Staging laparoscopy was performed, which was followed 4 weeks later by a radical D2 gastrectomy with end-to-side Roux-en Y oesophago-jejunostomy reconstruction. The patient made a satisfactory and uneventful recovery, and she was discharged 3 weeks postoperatively.

The peritoneal lesions sampled during the laparoscopic staging procedure were located on the left side, below the dome of the diaphragm. They appeared pale and well circumscribed, and each measured 0.2 cm in diameter. They were sent for frozen section examination, with subsequent examination of paraffin wax sections.

Frozen section examination showed both the lesions to have an angiomatoid architecture, consisting of tubular channels lined by epithelioid and flattened cells, set in a fibrous stroma. Many cells had vacuolated cytoplasm and a signet-ring morphology, and these features were suspected to indicate adenocarcinoma, but paraffin wax section examination was required for a definitive diagnosis.

Further examination of H&E-stained paraffin wax sections showed similar features as those noted on frozen section examination (fig 1A), but there was a distinct lack of cytological atypia, and mitotic figures were not identified. These lesions appeared benign, and the possibility of them being of mesothelial derivation was considered. Immunohistochemical examination was requested, including a panel of mesothelial markers. Unfortunately, the material available was insufficient to perform the full set of markers on one of the peritoneal lesions. The morphological features, supported by immunohistochemistry, were diagnostic of adenomatoid tumours, and the possibility of them representing metastatic adenocarcinoma was therefore excluded.

The surgical team were now confident to proceed to gastrectomy, at which time a further lesion was detected, involving the surface of the posterior aspect of the left lobe of the liver. It appeared pale yellow and well circumscribed, similar to the peritoneal lesions, and measured 0.5 cm in diameter. On frozen section examination, it appeared unencapsulated, with a non-infiltrating and well-circumscribed margin, and with a morphology similar to that seen in the peritoneal lesions. Its circumscription, together with a lack of cytological atypia, led to an opinion that it was not malignant.

The architecture was angiomatoid, raising the possibility of it representing a vascular neoplasm, which was probably benign owing to a lack of cytological atypia (fig 1B). On frozen section examination, it was stated that the lesion appeared unusual, and hence paraffin wax sections were required to make a confident diagnosis.

The liver lesion had a histochemical and immunohistochemical profile identical to that of the peritoneal lesions, and the features were thought to be classic of an adenomatoid tumour (fig 1C,D).

Examination of the gastric tumour (fig 1E) showed a well-moderately differentiated adenocarcinoma, with focal penetration into the submucosa (stage pT1pN0pMx). Interestingly, the morphology of this tumour showed many similarities to the adenomatoid tumours from the peritoneum and liver. There was no evidence of lymphovascular space invasion, and none of the lymph nodes sampled showed any evidence of metastatic disease. A similar