Table 1 Clinicopathological features of pulmonary myelolipoma cases

Case	Author (reference)	Age (years)/sex	Background	Symptoms	Tumour location	Tumour size (cm)	Adipose tissue (%)	Bone spicula	Diagnosis
1	Saleeby ¹	81/F	Pneumonia	–	B; P	2.5	ND	–	Autopsy
2	Hunter ²	70/F	RA, steroid	–	B, Mu; P	ND	50	ND	Biopsy
3	Ziólkowski <i>et al</i> ³	49/M	Pneumonia	–	LLL	7x5x5	ND	ND	Autopsy
4	Ziólkowski <i>et al</i> ³	59/M	Pneumonia	–	RLL	2	ND	ND	Resection
5	Zunarelli <i>et al</i> ⁴	52/M	MGUS, BC	–	RUL; C	ND	ND	ND	Lobectomy
6	Sabate and Shahian ⁵	54/M	Hypercholesterolaemia	–	LUL; C	2.5	60–100	+	Enucleation
7	Lu and Xiao ⁶	45/M	Pneumonia	–	LUL; C	1.5	60	+	Lobectomy
8	Sato <i>et al</i> (present case)	71/M	Lung cancer	–	LLL; C	2	60	+	Autopsy

B, bilateral; BC, bronchial carcinoid; C, centrally; F, female; LLL, left lower lobe; LUL, left upper lobe; M, male; MGUS, monoclonal gammopathy of undetermined significance; Mu, multiple; ND, not determined; P, peripherally; RA, rheumatoid arthritis; RLL, right lower lobe; RUL, right upper lobe; +, present; –, absent.

think that the multiple bone metastases of the lung carcinoma were scarcely associated with its development.

Gowitt and Zaatari¹¹ reported a patient with bronchial EMH in whom chronic myelogenous leukaemia later developed. The bronchial wall was thickened by calcification and ossification of the cartilage, and the EMH in the ossified area. The EMH extended into the surrounding tissues, and the margin of the lesion was ill defined. In the present case, the bronchial wall structure including bronchial cartilage was well preserved, which is a finding different from the EMH case discussed above.

Shuangshoti¹² described a patient with osseous metaplasia of the bronchus who had tuberculosis for 6 years. Multiple bronchi contained small foci of submucosal osseous metaplasia continuous with the bronchial cartilage. A bone marrow component in the foci was not mentioned in that report, but haematopoietic cells were probably scattered with fat tissue in the lesion, judging from the associated picture. Chronic inflammation might be associated with the development of such lesions. In the present case, the lesion itself was histologically similar to this metaplastic lesion, but there were differences in the solitary lesion, the non-ossified bronchial cartilage and the fact there was no history of chronic inflammatory pulmonary diseases.

A bone element in myelolipoma is controversial; Fowler *et al*¹⁰ described that no bone spicules were observed in myelolipomas, and cases in which bone spicule was observed were excluded from myelolipomas. In the lung, two previously reported cases and the present case show trabecular bone.^{6,7} Centrally located tumours in two cases and in the present case were located close to the bronchus.^{5,7} A bone component in pulmonary myelolipoma might be associated with the site of tumour development that is adjacent to the bronchial cartilage.

Myelolipomas can be detected incidentally at autopsy and do not often exceed 5 cm in diameter, and rarely present as a giant mass.¹³ Pulmonary myelolipomas are mostly small nodules, but a case of a myelolipoma, 7 cm in maximal diameter was reported.³ The lesion in this case had not changed size for 10 years, and had no apparent neoplastic feature. Chang *et al*¹⁴ reported an adrenal myelolipoma with balanced chromosomal translocation that suggested a true neoplastic nature. Myelolipoma is a heterogeneous concept in adrenal and extra-adrenal sites. Some cases are genuine neoplasms, but others might be tumour-like lesions. The histogenesis of myelolipoma is not evident, although there are several theories: embolic origin, heterotopic bone marrow

and metaplasia from multipotential mesenchymal stem cells. The present case suggests that some pulmonary myelolipomas are not true neoplasms but can be metaplastic lesions arising in the peribronchial region.

Acknowledgements

We thank Hideaki Ninomiya, Department of Pathology, Kanazawa Medical University Hospital for assistance with immunohistochemistry.

**Katsuaki Sato, Yoshimichi Ueda,
Shogo Katsuda**

Department of Pathophysiological and Experimental Pathology, Kanazawa Medical University, Ishikawa, Japan

Katsuma Tsuchihara

Department of Respiratory Medicine, Kanazawa Medical University, Ishikawa, Japan

Correspondence to: Dr K Sato, Department of Pathophysiological and Experimental Pathology, Kanazawa Medical University, 1-1 Daigaku, Uchinada, Kahoku, Ishikawa 920-0293, Japan; sato-k@kanazawa-med.ac.jp

doi: 10.1136/jcp.2005.033340

Accepted 14 October 2005

Competing interests: None declared.

References

- 1 Saleeby ER. Heterotopia of the bone marrow without apparent cause. *Am J Pathol* 1925;1:69–78.
- 2 Hunter SB, Schemankewitz EH, Patterson C, *et al*. Extra-adrenal myelolipoma. A report of two cases. *Am J Clin Pathol* 1992;97:402–4.
- 3 Ziolkowski P, Muszczyńska-Bernhard B, Dzięgiel P. Myelolipoma: the report of two cases in pulmonary location. *Pol J Pathol* 1996;47:141–2.
- 4 Zunarelli E, Criscuolo M, Malavolta L, *et al*. A case of lung myelolipomatosis in a patient with bronchial carcinoid. *Panminerva Med* 1999;41:175–8.
- 5 Sabate CJ, Shahian DM. Pulmonary myelolipoma. *Ann Thorac Surg* 2002;74:573–5.
- 6 Lu X, Xiao L. Myelolipoma of the lung: a case report. *Chin Med J* 2003;116:951–3.
- 7 Franiel T, Fleischer B, Raab BW, *et al*. Bilateral thoracic extra-adrenal myelolipoma. *Eur J Cardiothorac Surg* 2004;26:1220–2.
- 8 Kumar PV, Arasteh M, Musallaye A, *et al*. Fine needle aspiration diagnosis of extramedullary hematopoiesis presenting as a right lung mass. *Acta Cytol* 2000;44:698–9.
- 9 Asakura S, Colby TV. Agnogenic myeloid metaplasia with extramedullary hematopoiesis and fibrosis in the lung: report of two cases. *Chest* 1994;105:1866–8.
- 10 Fowler MR, Williams RB, Alba JM, *et al*. Extra-adrenal myelolipomas compared with extramedullary hematopoietic tumors: a case of

presacral myelolipoma. *Am J Surg Pathol* 1982;6:363–74.

- 11 Gowitt GT, Zaatari GS. Bronchial extramedullary hematopoiesis preceding chronic myelogenous leukemia. *Hum Pathol* 1985;16:1069–71.
- 12 Shuangshoti S. Metaplasia of bone in lungs and bronchi: report of 2 cases. *J Med Assoc Thai* 1995;78:103–7.
- 13 Boudreaux D, Waisman J, Skinner DG, *et al*. Giant adrenal myelolipoma and testicular interstitial cell tumor in a man with congenital 21-hydroxylase deficiency. *Am J Surg Pathol* 1979;3:109–23.
- 14 Chang KC, Chen PI, Huang ZH, *et al*. Adrenal myelolipoma with translocation (3;21)(q25;p11). *Cancer Genet Cytogenet* 2002;134:77–80.

Adverse effects of dyes used in sentinel node biopsy on immunocytochemical determination of hormone receptors in breast cancer cells

Sentinel lymph node biopsy (SLNB) is currently used in the UK for axillary staging in breast cancer. Node localisation is achieved with the combination of injected dye and radioactive tracer.¹ The commonly used dyes are the isomers isosulphan blue and Patent blue V (PBV).^{1–3} Methylene blue (MB) is also used in some centres as it is cheaper and easily available.⁴ In the Far East, other dyes such as Indigocarmine (IDC) are also being used.⁵

In the 1980s, when dyes were used to localise occult breast cancers before surgery, studies were carried out on the effect of the dyes on oestrogen receptor (ER α) assessment by ligand-binding assay. These showed a decrease in the binding capacity after treatment with MB but not with isosulphan blue.^{6,7} Given the importance of the determination of ER α expression in the treatment of breast cancer and the wider use of SLNB to stage the axilla, we investigated whether the dyes used change the immunocytochemical profile of these receptors.

Materials and methods

The ER α -positive and progesterone receptor (PR)-positive malignant breast cell line MCF-7 was cultured under standard conditions and exposed to the dyes diluted 1:10, 1:100 and 1:1000 in triplicate for 4 and 24 h. The dyes used were 2.5% PBV (Guerbert, France), 1% MB (Martindale Pharmaceuticals, UK) and 4% IDC (Amino), which are the concentrations used clinically for the SLNB procedure. Appropriate controls were established in parallel. After incubation, cells were trypsinised and embedded in paraffin wax blocks.⁸

Figure 1 MCF-7 cells treated with the dyes for 24 h and stained for oestrogen receptor (ER α). (A) Treated with methylene blue 1%, 1:10 dilution; ER α is negative. (B) Treated with Patent blue V 2.5%, 1:10 dilution; ER α shows positive staining. (C) Treated with Indigocarmine 0.4%, 1:10 dilution; ER α shows strong positivity. (D) Negative controls. (E) Positive controls.

Sections were cut and stained as per the procedures routinely used for ER α , PR and cytokeratin MNF116. The primary antibodies used ER-6F11 (Novocastra, Newcastle upon Tyne, UK) diluted 1:30, PgR 636 (DakoCytomation, Glostrup, Denmark) diluted 1:50 and MNF116 (DakoCytomation) diluted 1:100. Secondary detection was achieved using the ABC technique (DakoCytomation) for ER α and Envision (DakoCytomation) for PR and MNF116. Slides were incubated with DAB and counterstained with haematoxylin. Negative and positive controls were included with each batch. MNF116 was used to determine whether there was evidence to the fact that general immunoreactivity of the samples was affected by the overall effect of the dyes.

Slides were scored blindly using the Allred (quick) score for ER α and PR by two specialist breast histopathologists (AMH and AMS), and for any evidence of diminution in staining

intensity for MNF116 by AMH.⁸ Statistical analysis was performed using InStat software V.3.06 with Dunn's multiple comparison test comparing the different dyes and dilutions with the controls. *p* Values of <0.05 were considered significant.

Results

With higher concentrations of MB dye (1:10 and 1:100 dilutions of the 1% MB preparation) there was a significant decrease in both ER α and PR scores after 4 and 24 h exposure except for the PR 1:100 dilution, 4 h exposure (fig 1). No significant change was seen with the MB 1:1000 dilution (table 1). On 24 h exposure, cells incubated with PBV showed a non-significant decrease in ER α score on the 1:10 dilution, whereas IDC showed a non-significant decrease in PR score with the 1:10 dilution.

All the slides stained strongly positive for the MNF116 on 24 h exposures, and, as no effect was seen, they were not stained for the shorter exposure.

Discussion

The presence of ER α in breast cancer is an established prognostic marker determining the likelihood of response to anti-hormonal therapy. PR is a marker of ER α functionality. ER α -positive and PR-positive tumours are more likely to respond to anti-hormonal treatment than ER α -positive and PR-negative tumours; there is some evidence that ER α -positive and PR-negative tumours are more likely to respond to aromatase inhibitors than tamoxifen.⁹⁻¹¹ In this study, we have explored the hypothesis that the dyes used in SLNB could interfere with hormonal receptor determination. A significant diminution in scoring was seen with cells treated with MB at 1:10 and 1:100 dilutions, with some samples rendered negative. This was not seen with PBV and IDC (table 1). These results suggest that an inherently weak ER α expresser could be rendered negative after exposure to MB with consequent withholding of potentially beneficial treatment. However, it is difficult to accurately relate *in vitro* findings to an *in vivo* situation. No similar effect was seen on the results of MNF116 immunolabelling, and the effects of these dyes on other markers would need case-by-case evaluation.

Hirsch *et al*⁶ measured MB tissue concentrations in breast tissue samples from a patient who had 0.1–0.2 ml of 1% MB injected, using spectrophotometry. This showed tissue concentration of 0.08, which is roughly equal to 1:10 dilution of the dye. The volume used for SLNB is larger, with 2 ml routinely injected for PBV and up to 5 ml for MB.^{1 4 12 13} Although the recommended site of injection for PBV in the UK is subdermal, around the areola, this is not the case with MB because of potential injection site complications, especially skin necrosis.¹⁴ The site of injection for MB is deeper in the breast tissue.^{13 15} Deep subareolar injections of MB have been previously described.¹² Probably, with deeper peritumoral injections, the concentration of MB in the tumour could be in the region of 1:10 or even higher because of the larger volume used, whereas it should be lower with injections at sites more distant from the tumour. These concentrations are comparable to those that reduced ER α and PR status in our experimental system. The time durations of exposure to the chosen dyes were pragmatic:

Table 1 Median (range) oestrogen and progesterone receptor score for each dilution of the three dyes

Dye (dilution)	Oestrogen receptor				Progesterone receptor			
	After 4 h exposure		After 24 h exposure		After 4 h exposure		After 24 h exposure	
	Median (range)	<i>p</i> Value	Median (range)	<i>p</i> Value	Median (range)	<i>p</i> Value	Median (range)	<i>p</i> Value
MB (1:10)	4 (0–6)	<0.01	3 (2–4)	<0.05	1.5 (0–6)	<0.01	3 (2–4)	<0.05
MB (1:100)	6.5 (6–7)	<0.05	2.5 (2–6)	<0.05	7 (6–8)	NS	4 (0–7)	<0.05
MB (1:1000)	8 (8)	NS	7 (7–8)	NS	8 (7–8)	NS	8 (8)	NS
PBV (1:10)	8 (8)	NS	6 (5–6)	NS	8 (7–8)	NS	8 (7–8)	NS
PBV (1:100)	8 (8)	NS	8 (7–8)	NS	8 (8)	NS	8 (8)	NS
PBV (1:1000)	8 (8)	NS	8 (7–8)	NS	8 (8)	NS	8 (8)	NS
IDC (1:10)	8 (8)	NS	8 (7–8)	NS	8 (7–8)	NS	7 (6–8)	NS
IDC (1:100)	8 (7–8)	NS	8 (8)	NS	8 (7–8)	NS	7.5 (6–8)	NS
IDC (1:1000)	8 (8)	NS	8 (8)	NS	8 (8)	NS	8 (8)	NS

IDC, Indigocarmine; MB, Methylene blue; NS, not significant; PBV, Patent blue V.

The *p* value is calculated using Dunn's multiple comparison test comparing the scores of the dye with the positive controls.

the shorter 4 h exposure to represent the duration after a straightforward excision transferred promptly to pathology, whereas the longer time was chosen to represent delayed fixation.

The concentration dynamics and the cell time exposures are likely to be highly complex. In cell line studies, direct and rapid exposure of the cells to the dye occurs. In the context of a complex tissue, even when the dye can be directly visualised pervading the tumour on gross analysis, the uniformity and certainty of cell exposure can only be hypothesised.

In summary, MB; which is one of the dyes used for SLNB worldwide, can reduce ER α and PR staining levels but not MNF116, *in vitro*. We suggest that if MB is used for SLNB, ER α should be checked on the core biopsy, if it was negative in the excised tumour.

Y A Masannat

Department of General Surgery, The Leeds Teaching Hospitals NHS Trust, Leeds General Infirmary, Leeds, UK

A M Shaaban

Department of Pathology, The Leeds Teaching Hospitals NHS Trust, Leeds General Infirmary, Algernon Firth Institute, Leeds, UK

V Speirs

Pathology & Tumour Biology, Wellcome Trust Brenner Building, Leeds Institute for Molecular Medicine, St James's University Hospital, Leeds, UK

G Coast, P Jackson

Department of pathology, The Leeds Teaching Hospitals, NHS Trust, Leeds General Infirmary, Leeds, UK

K Horgan

Department of General Surgery, The Leeds Teaching Hospital NHS Trust, Leeds General Infirmary, Leeds, UK

A M Hanby

Pathology & Tumour Biology, Wellcome Trust Brenner Building, Leeds Institute for Molecular Medicine, St James's University Hospital, Leeds, UK

Correspondence to: Mr Y A Masannat, Breast Unit/ Surgical Department, Leeds General Infirmary, Great George Street, Leeds LS1 3EX, UK; yazanmas@hotmail.com

doi: 10.1136/jcp.2006.044974

Accepted 28 December 2006

Funding: We thank the Breast Cancer Research Action Group for their continuous support.

Competing interests: None.

References

- Mansel RE, Fallowfield L, Kissin M, *et al*. Randomized multicenter trial of sentinel node biopsy versus standard axillary treatment in operable breast cancer: the ALMANAC Trial. *J Natl Cancer Inst* 2006;**98**:599–609.
- Giuliano AE, Haigh PI, Brennan MB, *et al*. Prospective observational study of sentinel lymphadenectomy without further axillary dissection in patients with sentinel node-negative breast cancer. *J Clin Oncol* 2000;**18**:2553–9.
- Masannat Y, Shenoy H, Speirs V, *et al*. Properties and characteristics of the dyes injected to assist axillary sentinel node localization in breast surgery. *EJSO* 2006;**32**:381.
- Simmons R, Thevarajah S, Brennan MB, *et al*. Methylene blue dye as an alternative to isosulfan blue dye for sentinel lymph node localization. *Ann Surg Oncol* 2003;**10**:242–7.
- Imoto S, Wada N, Hasebe T, *et al*. Sentinel node biopsy for breast cancer patients in Japan. *Biomed Pharmacother* 2002;**56**(Suppl 1):192S–5S.
- Hirsch JI, Banks WL Jr, Sullivan JS, *et al*. Effect of methylene blue on estrogen-receptor activity. *Radiology* 1989;**171**:105–7.

- Hirsch JI, Banks WL Jr, Sullivan JS, *et al*. Noninterference of isosulfan blue on estrogen-receptor activity. *Radiology* 1989;**171**:109–10.
- Allred DC, Harvey JM, Berardo M, *et al*. Prognostic and predictive factors in breast cancer by immunohistochemical analysis. *Mod Pathol* 1998;**11**:155–68.
- Dowsett M, Cuzick J, Wale C, *et al*. Retrospective analysis of time to recurrence in the ATAC Trial according to hormone receptor status: an hypothesis-generating study. *J Clin Oncol* 2005;**23**:7512–17.
- Early Breast Cancer Trialists' Collaborative Group. Tamoxifen for early breast cancer: an overview of the randomised trials. *Lancet* 1998;**351**:1451–67.
- Barnes DM, Hanby A. Oestrogen and progesterone receptors in breast cancer: past, present and future. *Histopathology* 2001;**38**:271.
- Nour A. Efficacy of methylene blue dye in localization of sentinel lymph node in breast cancer patients. *Breast J* 2004;**10**:388–91.
- Blessing WD, Stoller AJ, Teng SC, *et al*. A comparison of methylene blue and lymphazurin in breast cancer sentinel node mapping. *Am J Surg* 2002;**184**:341–5.
- Sulhab M, Al Sarakbi W, Mokbel K. Skin and fat necrosis of the breast following methylene blue dye injection for sentinel node biopsy in a patient with breast cancer. *Int Semin Surg Oncol* 2005;**28**:26.
- Simmons RM, Smith SM, Osborne MP. Methylene blue dye as an alternative to isosulfan blue dye for sentinel lymph node localization. *Breast J* 2001;**7**:181–3.

Comparison of estimated glomerular filtration rate with routine creatinine clearance using a kinetic alkaline picrate assay from Olympus Diagnostica

Established chronic kidney disease (CKD) requires treatment with dialysis or transplantation, both of which are expensive. In addition, the late referral of patients requiring renal replacement therapy can have negative effects.^{1–4} As a result, it is recommended that estimated glomerular filtration rate (eGFR) using the Modification of Diet in Renal Disease (MDRD) equation be used to identify people with CKD and those at risk of end-stage CKD.^{5–7} However, MDRD-derived eGFRs are reportedly less precise above 60 ml/min/1.73 m² when a kinetic alkaline picrate assay is used.⁸

We derived eGFR using the four-variable MDRD equation in 100 patients undergoing the creatinine clearance test (CCT) as assessed by 24 h urine collections using a kinetic alkaline picrate assay (Olympus Life and Material Science Europe (Irish Branch) Lismeehan, O'Callaghan's Mills, Ireland).

$GFR (ml/min/1.73 m^2) = 186 \times ((\text{serum creatinine } (\mu\text{mol/l})/88.4)^{-1.154}) \times \text{age (years)} \times 0.742$ if female and $\times 1.21$ if African American.

Urine protein was requested for 85 patients and was measured using pyrogallol red-molybdate (Olympus Diagnostica). Absolute and relative difference plots were drawn (fig 1) and the median bias was calculated for all stages of CKD. For each level of proteinuria, the median percentage difference was calculated (table 1).

As neither the CCT nor the eGFR is a reference method, the mean of both these values was used for the classification of the 100 patients. The individual means ranged from 7 to 178 ml/min/1.73 m², with 22, 22, 39, 12 and 4 patients having values consistent with CKD stages 1, 2, 3, 4 and 5, respectively. The absolute difference plot showed that the difference between the CCT and eGFR increased as the average of the eGFR and CCT increased. This is not unexpected as the range

of the mean of the eGFR and CCT evaluated in numerical terms is relatively large; furthermore, these data are similar to those obtained when deriving the MDRD equation.⁹ However, the percentage difference plot showed no increase in the relative difference with increasing mean of the eGFR and CCT values (fig 1). Unfortunately, the authors of the MDRD equation did not provide any data on relative difference. Compared with the mean of the eGFR and CCT, the CCT had a median positive (95% CI) bias of 49.4% (31.0% to 60.8%), 43.3% (24.4% to 54.1%), 36.1% (31.3% to 41.8%), 34.5% (11.1% to 53.1%) and 60.4% (36.4% to 75.9%) in CKD stages 1, 2, 3, 4 and 5, respectively, with an overall median bias of 39.3% (35.3% to 47.3%). The Kruskal–Wallis one-way analysis of variance by ranks¹⁰ was used to see whether there was a difference between at least two of the medians of the five CKD stages. H was computed to be 5.75 for 4 degrees of freedom. As H was less than $\chi_{0.05} = 9.49$, there was no significant difference between any of the medians. The Passing Bablok linear regression equation¹¹ with creatinine clearance as x and eGFR as y was $y = 0.613x + 2.9$ with the 95% CIs for the slope and intercept being 0.55 to 0.682 and –0.7 to +6.2, respectively. Thus, the slope is significantly different from 1.0 and is consistent with a bias of approximately 38.7% between the two methods. The intercept is not significantly different from 0.0. The correlation coefficient r was calculated as 0.857 (0.869 if the data point with an absolute difference of 157 ml/min is excluded).

The relationship between proteinuria and the median percentage difference between creatinine clearance and eGFR was investigated. The protein concentration ranged from below the limit of detection (0.05 g/l) to 10.5 g/day (table 1). The Kruskal–Wallis one-way analysis of variance by ranks¹⁰ was used to see whether there was a difference between at least two of the medians of the five CKD stages. H was computed to be 3.38 for 6 degrees of freedom. As H was less than $\chi_{0.05} = 12.59$, there was no significant difference between any of the medians.

The CCT overestimates the true GFR because of renal secretion of creatinine. Ensuring accurately timed and complete urine collection is also notoriously difficult. Despite the labour-intensive CCT being far from ideal, it is routinely used in clinical practice. The four-variable MDRD equation has been criticised for its potential for patient misclassification due to analytical interference, imprecision or calibration issues.¹² However, these issues also affect the CCT—for example, the intraindividual coefficient of variation for the CCT and serum creatinine are 13.6% and 4.3%, respectively (<http://www.westgard.com/biodatabase1.htm>). As serum creatinine is the only one of the four variables that has imprecision, the eGFR should have a much smaller CI for individual patients than the CCT. Accordingly, the routine use of eGFR would result in fewer individual patients having a CI that crosses one of the CKD classification cut-offs compared with the CCT. The large intraindividual coefficient of variation for the CCT is probably the main reason for the poor correlation coefficient by method comparison standards ($r = 0.857$) found in this study. Added to this is the relatively large positive bias in the CCT owing to active secretion of creatinine. Compared with the CCT, the eGFR in our study is negatively biased (as expected) by about 39%, with no statistically significant difference