Figure 2 Amniotic membrane (AM) sutured around an Illog shell and left in place on the eye for 2 weeks does not show signs of mechanical irritation or enzyme degradation or inflammation. The photograph of the bio-onlay after 2 weeks on the ocular surface (A,B) does not show signs of destruction or severe inflammation. Histological examination for acid mucopolysaccharides (using Alcian blue staining; C) and periodic acid-Schiff (D) demonstrates intact anatomy of amniotic membrane. Immunohistochemical analysis does not reveal significant accumulation of CD45 leucocytes (E) or CD68 macrophages (F) on the surface or within the bio-onlay.

F E Kruse

Department of Ophthalmology, Friedrich-Alexander University, Erlangen-Nuremberg, Erlangen, Germany

Correspondence to: Dr C Cursiefen, Department of Ophthalmology, Friedrich-Alexander University, Erlangen-Nuremberg, Schwabachanlage 6, 91054 Erlangen, Germany; claus.cursiefen@augen.imed.uni-erlangen.de

doi: 10.1136/bjo.2006.108134

Accepted 4 November 2006

Funding: Interdisciplinary Center for Clinical Research (IZKF) Erlangen (A9), ELAN Fonds (AZ: 05.06.04.1).

Competing interests: None declared.

References

- 1 Lee SH, Tseng SC. Amniotic membrane transplantation for persistent epithelial defects with ulceration. *Am J Ophthalmol* 1997;123:303-12.
- 2 Kruse FE, Rohrschneider K, Völcker HE. Multilayer amniotic membrane transplantation for reconstruction of deep corneal ulcers. *Ophthalmology* 1999;106:1504-10.
- 3 Gomes JA, Romano A, Santos MS, et al. Amniotic membrane use in ophthalmology. *Curr Opin Ophthalmol* 2005;16:233-40.
- 4 Resch M, Schlotzer-Schrehardt U, Hofmann-Rummelt C, et al. Adhesion structures of amniotic membranes integrated into human corneas. *Invest Ophthalmol Vis Sci* 2006;47:1853-61.
- 5 Li H, Niederkorn J, Neelam S, et al. Immunosuppressive factors secreted by human amniotic epithelial cells. *Invest Ophthalmol Vis Sci* 2005;46:900-7.
- 6 Cursiefen C, Chen L, Borges L, et al. VEGF-A stimulates lymphangiogenesis and hemangiogenesis in inflammatory neovascularization via macrophage recruitment. *J Clin Invest* 2004;113:1040-50.
- 7 Park WC, Tseng SCG. Modulation of acute inflammation and keratocyte death by suturing, blood, and amniotic membrane in PRK. *Invest Ophthalmol Vis Sci* 2000;41:2906-14.
- 8 Li W, He H, Kawakita T, et al. Amniotic membrane induces apoptosis of interferon-gamma activated macrophages in vitro. *Exp Eye Res* 2006;82:282-92.

Ocular myasthenia: a rare presentation with MuSK antibody and bilateral extraocular muscle atrophy

Atrophic extraocular muscles are rarely observed¹⁻³ and bilateral extraocular muscle (EOM) atrophy is extremely rare.³ It has been reported in chronic progressive external ophthalmoplegia (CPEO) related to mitochondrial myopathies,⁴ myotonic dystrophy type 1⁵ and in three other cases of myasthenia gravis (MG).⁴ This report is an atypical presentation of ocular MG with positive muscle-specific kinase (MuSK) antibodies and bilateral EOM atrophy.

Case report

A 49-year-old man presented with a 12-year history of slowly isolated progressive external ophthalmoplegia. His medical history was unremarkable and he did not take any drugs, smoke or drink alcohol. Examination revealed limitation of gaze in all directions to 3°-5° with bilateral ptosis and no exophthalmos. He had no other facial, bulbar or peripheral weakness. His acetylcholine receptor ϵ subunit-specific antibodies and γ subunit-specific antibodies were negative, but his anti-MuSK antibody level was increased at 32 693 pg/ml (normal values 249-264 pg/ml). Single-fibre electromyography revealed jitter and blocking that confirmed his diagnosis of MG. Complete blood count and routine chemistry laboratories were normal. Sedimentation rate, rapid plasma reagin, rheumatoid factor, antinuclear antibody, anti-Hu, anti-Lambert-Eaton myasthenic syndrome, anti-Yo and anti-MaTa antibodies were all negative. Thyroid testing and mitochondrial DNA rearrangement analysis for Kearns-Sayre syndrome were negative. MRI of the orbits revealed severe wasting of all extraocular muscles, except for the inferior oblique muscle (fig 1). Chest CT scan with contrast was normal. Even after 6 months of pyridostigmine and prednisone treatment, followed by thymectomy, his extraocular motility did not improve for 2½ years.

Comment

We have found no previously published cases of bilateral EOM atrophy in MuSK antibody-positive patients with myasthenia. The diagnosis of MG was confirmed by single-fibre electromyography to rule out the possibility of a false-positive increased level of anti-MuSK antibodies. Although >50% of patients with MG present with symptoms and signs related to EOM weakness,⁶ such as diplopia or ptosis, bilateral EOM atrophy is unusual. Unlike the three patients reported by Bau *et al*⁷ and Caress *et al*,⁸ who had anti-MuSK-positive ocular myasthenia with normal extraocular muscles on MRI, our patient had bilateral EOM atrophy. Unlike

Figure 1 On axial postcontrast T1-weighted MRI of the orbits, the rectus muscles seem to be "string-like pulleys". These muscles lost their normal fusiform shape and became thin ribbon-like bands.

the three patients with myasthenia with bilateral EOM atrophy reported by Okamoto *et al.*,⁴ who were positive for only acetylcholine-receptor antibodies, our patient was seronegative for these antibodies but positive for MuSK antibody. In ocular myasthenia, only 50% of patients are found to have anti-acetylcholine receptor antibodies,⁶ and of 38 patients with seronegative ocular MG, none show increased titres of the anti-MuSK antibody.^{9,10} Between 38% and 47% of patients with seronegative MG have increased anti-MuSK antibodies.^{9,10} Similar to the clinical course of patients in the study by Okamoto *et al.*,⁴ our patient had an approximately 15 year duration of limited ocular motility that did not improve with medication or thymectomy.

Diplopia and ptosis are well-recognised signs of MG, but bilateral EOM atrophy is exceedingly rare. Although ocular involvement is a less common presentation of the anti-MuSK MG syndrome, it may be reasonable to check for anti-MuSK antibody in any patient who presents with progressive external ophthalmoplegia.

Jane W Chan

Department of Neurology, University of Nevada, School of Medicine, Las Vegas, Nevada, USA

William W Orrison

Nevada Imaging Center, Las Vegas, Nevada, USA

Correspondence to: J W Chan, Department of Neurology, University of Nevada School of Medicine, 1707 W Charleston Blvd, Suite 220, Las Vegas, Nevada 89102, USA; worjun@aol.com

doi: 10.1136/bjo.2006.108498

Accepted 10 October 2006

Competing interests: None declared.

References

- Hupp SL, Williams JP, Curran JE. Computerized tomography in the diagnosis of the congenital fibrosis syndrome. *J Clin Neuroophthalmol* 1990;10:135-9.
- Horton JC, Tsai RK, Truwit CL, *et al.* Magnetic resonance imaging of superior oblique muscle atrophy in acquired trochlear nerve palsy. *Am J Ophthalmol* 1990;110:315-16.
- Hansman ML, Peyster RG, Heiman-Patterson T, *et al.* CT demonstration of extraocular muscle atrophy. *J Comput Assist Tomogr* 1988;12:49-51.
- Okamoto K, Ito J, Tokiguchi S, *et al.* Atrophy of bilateral extraocular muscles. CT and clinic features of seven patients. *Neuroophthalmol* 1996;16:286-8.
- T, Matsubara E, Nagano I, Shoji M, *et al.* Bilateral extraocular muscle atrophy in myotonic dystrophy type 1. *Neurology* 2004;63:759-60.
- Kaminski HJ. Myasthenia gravis. In: Katriji B, Kaminski JH, Preston DC, Ruff RL, Shapiro BE, eds. *Neuromuscular disorders in clinical practice*. Boston, MA: Butterworth-Heinemann, 2002:916-30.
- Bau V, Hanisch F, Hain B, *et al.* Ocular involvement in MuSK antibody-positive myasthenia gravis. *Klin Monatsbl Augenheilkd* 2006;223:81-3.
- Caress JB, Hunt CH, Batish SD. Anti-MuSK myasthenia gravis presenting with purely ocular findings. *Arch Neurol* 2005;62:1939.
- McConville J, Farrugia ME, Beeson D, *et al.* Detection and characterization of MuSK antibodies in seronegative myasthenia gravis. *Ann Neurol* 2004;55:580-4.
- Sanders DB, El-Salem K, Massey JM, *et al.* Clinical aspects of MuSK antibody positive seronegative MG. *Neurology* 2003;60:1978-80.

Botulinum toxin injection causing lateral rectus palsy

A 35-year-old woman with brow spasms, treated in the past with Botox, was referred

for increasing tightness in the brow. She reported having experienced an episode of difficulty focusing after a previous injection of Botox by another physician. The patient had no response to a subcutaneous injection consisting of 10 units of Botox above the eyebrows, nor to a subsequent injection of 20 units in the same location 2 months later. A third injection of 35 units of Botox resulted in improvement of her brow spasms, but only for 48 h. Accordingly, 2 months after the third injection, she was injected with 75 units of Botox, with 30 units injected in three separate locations above each eyebrow and 7.5 units injected just lateral to each lateral canthus. At 1 week after this injection, the patient had difficulty focusing, and shortly thereafter noted horizontal binocular double vision, worse on left gaze.

On examination 48 h after the onset of her symptoms, the patient had normal visual acuity of 20/20 OU, with normal colour vision and full visual fields. She had anisocoria, with the right pupil being 0.5 mm larger than the left, but both pupils reacted briskly to light stimulation and both dilated equally well after topical administration of a 10% cocaine solution. The right eye moved fully in all directions. The left eye had mild limitation of abduction, but otherwise moved fully. Duction measurements revealed 60° of abduction of the right eye versus 50° of the left eye. In primary position, the patient had an esotropia of 9 prism diopters (PD) at distance and 2 PD at near. The esotropia was incomitant, and increased on left gaze to 15 PD. Corneal and facial sensation were equal and normal bilaterally. Slit-lamp examination and funduscopy were normal. MRI of the brain and orbit were normal.

The patient was treated with a 6-PD Fresnel prism placed base out on her left spectacle lens for temporary symptom relief. One month later, her diplopia had resolved and she was able to remove the prism.

Comment

This is the first reported case of lateral rectus paresis after an injection of Botox into the lateral canthal region. Inferior oblique paresis is an uncommon adverse effect of Botox injection into the lower lid,¹ with a reported incidence of 1.7%.² The mechanism is postulated to be diffusion of the medication to the underlying inferior oblique muscle. Diffusion of botulinum toxin to surrounding muscles has also been reported after treatment of dystonia with large doses of the drug.³ We believe that, in our case, Botox spread from the lateral canthus to the left lateral rectus muscle, producing a transient paresis of the muscle characterised by an incomitant esotropia and a mild left abduction deficit. It is interesting that patients who experience diplopia after a periorbital injection of Botox tend to have recurrence of diplopia when they are reinjected, suggesting that they have some predisposition to this complication.² Our patient had an episode of difficulty focusing after a previous periorbital injection of Botox by another physician, which was similar to the prodrome that she experienced just before she developed diplopia after the injection we gave her. Fortunately, the diplopia was transient, resolving as the effects of the botulinum toxin dissipated. We believe that patients who experience diplopia or have difficulty focusing after an injection of botulinum toxin should be counselled as to the potential recurrence of such an effect with subsequent injections of the drug, regardless of the care taken to avoid these sequelae.

Celia S Chen

Neuro-Ophthalmology Unit, Wilmer Eye Institute, The Johns Hopkins Hospital, Baltimore, Maryland, USA

Neil R Miller

Wilmer Eye Institute, The Johns Hopkins Hospital, Baltimore, Maryland, USA

Correspondence to: Dr N R Miller, Maumenee 127, Wilmer Eye Institute, Johns Hopkins Hospital, 600 North Wolfe Street, Baltimore, MD 21287, USA; nrmiller@jhmi.edu

doi: 10.1136/bjo.2006.109926

Accepted 31 October 2006

Competing interests: None.

Informed patient consent was obtained for the publication of the persons details in this report.

References

- Aristodemou P, Watt L, Baldwin C, *et al.* Associated with the cosmetic use of botulinum toxin A for facial rejuvenation. *Ophthalm Plast Reconstr Surg* 2006;22:134-6.
- Wutthiphan S, Kowal L, O'Day J, *et al.* Diplopia following subcutaneous injections of botulinum A toxin for facial spasms. *J Pediatr Ophthalmol Strabismus* 1997;34:229-34.
- Lange DJ, Brin MF, Warner CL, *et al.* Distant effects of local injection of botulinum toxin. *Muscle Nerve* 1987;10:552-55.

Abducens palsy and Sjögren's syndrome induced by pegylated interferon therapy

Interferons (IFNs) and their pegylated forms (PEG-IFNs) are widely used in the treatment of viral hepatitis and some neoplasms. Although ophthalmic symptoms are common among their various side effects,^{1,2} abducens palsy is rarely observed. Here, we describe a case where abducens palsy developed during PEG-IFN therapy, and discuss the management.

Case report

A 65-year-old man had undergone removal of stage IV renal cell carcinoma 5 years ago. He had been managed with IFN α therapy (6×10^6 IU three times a week) for recurrence. Although the therapy suppressed tumour growth effectively, it had moderate side effects including influenza-like symptoms, erythema and depression. After 4 years of conventional IFN therapy, PEG-IFN α became available for the patient.

On the first day of PEG-IFN therapy (3×10^6 IU), the patient had the expected fever and malaise. The next day he developed diplopia, which worsened gradually. After 1 week, the patient was referred to ophthalmologists and was diagnosed as having right abducens palsy (fig 1, upper panel). IFN-induced retinopathy was absent. Diabetes mellitus was negative with fasting glucose (73 mg/dl) and haemoglobin A1c (5.5%). He did not have hypertension. MRI showed no evidence of metastasis or intracranial hypertension, but showed enhancement of the right abducens nerve (fig 2).

After 1 month of onset of diplopia, painful swelling of the right parotid gland was noted. In another 1 month, dry eye and dry mouth sensations developed. The result of Schirmer's test (3 mm) and fluorescent corneal staining suggested presumable Sjögren's syndrome (SS). Furthermore, serum examination showed increased SS A and SS B antibody levels (128