Medical Memoranda

A Case of Primary Amyloid Heart

Cardiac amyloidosis is being reported with increasing frequency, but is still a rare disease; the number reported probably does not much exceed 200 (Lindsay, 1946; Thomashow *et al.*, 1953; Pruitt *et al.*, 1953; Benson and Smith, 1956; Clark and Bennett, 1957; Chambers *et al.*, 1958), and far fewer have been diagnosed in life. The first symptoms have usually been those of heart failure, but pain has been rare and murmurs have been unusual; arrhythmias and various abnormalities in the form of the electrocardiogram have been common. We describe the case of a man who had unusual combinations of abnormal rhythms, Stokes-Adams attacks due to complete heart-block, a systolic murmur, and only late congestive failure.

CASE REPORT

A man aged 41 came under the care of one of us (A. E. de la T. M.) in 1953. Most of the early symptoms, dating from 1942, were related to the nervous system; in order of time they were a swollen painful knee, lightning pains in the legs, a perforating foot ulcer, right foot-drop, left leg weakness, numbness of the legs below the knees, and tingling, numbness, and weakness in the hands and wrists. Examination in 1953 revealed wasting and weakness of the arms and legs, claw deformity of the hands, peripheral sensory loss in the four limbs, and absence of all deep reflexes; thickened nerves could be palpated, specially the posterior auricular and supraclavicular. The blood Wassermann and Kahn reactions were negative. A biopsy of the nerve showed amyloid. The neurological aspects have been reported elsewhere (Chambers *et al.*, 1958).

The first symptoms of cardiac involvement appeared in December, 1957, when he was said to have lost consciousness and the pulse was impalpable; later it was very rapid.

When admitted to Wimborne Hospital in January, 1957, examination showed the following: apex beat not felt; cardiac dullness absent; first and second sounds only; B.P. 120/75 mm. Hg; electrocardiogram (unsteady owing to limb tremor) showed normal rhythm; rate 65; P-R interval 0.3 second; right axis deviation; left bundle-branch block; T inversion in leads II, III, and VF. Chest x-ray examination showed slight general cardiac enlargement. A second E.C.G. in February (Fig. 1) was like the first except for less inversion of T in the limb leads. In May he was admitted to Poole General Hospital; his legs were weaker and he had upper chest pain front and back. Signs were the same except that a systolic murmur was heard at the xiphisternum. Blood examination showed serum protein electrophoresis curve normal, serum albumin 4.3,

serum globulin 2.9, total 7.2 g./100 ml. A/G ratio to 1.48 to 1. An E.C.G. showed auricular flutter; the degree of A-V block was fairly high without digitalis (Fig. 2).

FIG. 2.—May 5, 1957. Auricular flutter, auricular rate 240, ventricular rate 45-75; broad Q waves in leads I and VL; left bundle-branch block.

He was not seen again until February, 1959, when he said he had fainted a number of times; the features of these faints suggested Stokes-Adams attacks. An E.C.G. (Fig. 3) showed auricular flutter and complete heart-block, ventricular rate 30 a minute; right bundle-branch block and axis changes. The fainting stopped after the prescription of ephedrine $\frac{1}{2}$ gr. (32 mg.) t.i.d. The apex beat was now palpable in the mid-axillary line; other clinical signs were as before; chest x-ray examination showed further cardiac enlargement and a small left basal pleural effusion. In July he was readmitted with a recurrence of the Stokes-Adams attacks; the electrocardiogram was the same as in February, except that the ventricular rate was higher at 40 a minute; signs of systemic congestive failure were still absent; the spleen and liver were palpable. With adrenaline injections and ephedrine by mouth the Stokes-Adams attacks again stopped. A chest x-ray film in November (Fig. 4) showed great cardiac enlargement and slight pulmonary hyperaemia, but no pleural effusion. On February 23, 1960, he was found dead in bed at home.

POST-MORTEM FINDINGS

Necropsy (Dr. T. A. J. Wickham) showed a dry scaly skin, scattered papules, and subcutaneous nodules on the trunk and neck; ulceration of toes of right foot; destruction of left metatarsus and firm induration of soft tissues; subcutaneous tissues, very little fat, some oedema. Heart: weight 28 oz. (794 g.), enlarged in all diameters, and unusually rigid, the chambers retaining their contours as if formalin-fixed; right auricle and interauricular septum much thickened and of a curious buff colour; left ventricle

FIG. 3.—February 19, 1959. Auricular flutter and complete heart-block; ventricular rate 30; right bundle-branch block and axis changes.

FIG. 4.—November, 1959. Chest x-ray film showing great cardiac enlargement, slight pulmonary hyperaemia, but no pleural effusion.

dilated and hypertrophied, showing two clearly defined zones on section—an outer of normal appearance and an inner buff-coloured which continued into the greatly thickened interventricular septum; valves normal apart from slight thickening of free margins; coronary arteries slightly thickened but patent. Lungs: bilateral pleural effusions of 1 litre; apical scars and emphysema; lower zones oedematous; tongue large, showing a buff-coloured zone on section. Liver: 83 oz. (1,872 g.), firm and congested, without microscopical evidence of amyloid. Spleen very firm, not enlarged. Pancreas, adrenals, and pituitary normal. Kidneys: early bilateral hydronephrosis and disorganization of parenchyma; ureters and bladder dilated. Prostate slightly enlarged and firm. Meninges thickened. Brain oedematous and congested. Nerves: brachial and sciatic plexuses appeared normal.

Histology.—Amyloid was demonstrable in varying amounts in all but a few of the sections. It was absent in the meninges, trachea, lungs, liver, seminal vesicles, and testes, but the blood-vessels transverse these tissues showed deposits on their walls. Most heavily affected were the tongue, heart, and kidneys; the heart (of which sections of right auricle, interventricular septum, and left lateral wall of left ventricle were examined) showed massive deposits of amyloid between the muscle fibres which were degenerate at the periphery of the deposits. Approximately 10% of the renal glomeruli were involved and traces of amyloid were present in the capsule and perirenal fat, in the form of amyloid "rings." Similar intermuscular deposits were to be seen in the prostate. In the spleen, lymph nodes, nerve plexuses, adrenals, and skeletal muscle the amyloid was confined to the supporting stroma.

COMMENT

The man whose history is described suffered from severe polyneuritis, the cause of which was unknown until a biopsy of nerve revealed amyloid degeneration. Fourteen years after the onset of the nerve lesion symptoms of cardiac involvement appeared, the first being syncope due to one of the abnormal rhythms with tachycardia. Later there were partial heart-block, T-wave inversion, bundle-branch block, auricular flutter, and complete heart-block with Stokes-Adams attacks. There was progressive cardiac enlargement, and at necropsy the heart weight was 28 oz. (794 g.), amyloid being a heavy material. Other unusual features were the systolic murmur and the late appearance of congestive failure.

Amyloid cardiopathy is a rare form of chronic heart disease and unless borne in mind may be overlooked, especially when there is no chronic sepsis. Amyloid disease should be thought of when cardiac enlargement and abnormal rhythm with conduction defect are present; when there is widespread T inversion unlike the usual ischaemic patterns; when there is involvement of other viscera and of the nervous system; and when there is no evidence of hypertensive, valvular, or ischaemic heart disease.

HAROLD COOKSON, M.D., B.Sc., F.R.C.P.,
Physician, Poole General Hospital.

A. E. DE LA T. MALLETT, D.S.C., M.D.,
Late Honorary Clinical Assistant, Poole General Hospital.

REFERENCES

- Benson, R., and Smith, J. F. (1956). *Brit. Heart J.*, **18**, 529.
Chambers, R. A., Medd, W. E., and Spencer, H. (1958). *Quart. J. Med.*, **27**, 207.
Clark, R. M., and Bennett, R. F. (1957). *Lab. Invest.*, **6**, 125.
Lindsay, S. (1946). *Amer. Heart J.*, **32**, 419.
Pruitt, R. D., Daugherty, G. W., and Edwards, J. E. (1953). *Circulation*, **8**, 769.
Thomashow, A. I., Angle, W. D., and Morrione, T. G. (1953). *Amer. Heart J.*, **46**, 895.