RUNT DISEASE*

BY

N. W. NISBET, F.R.C.S.Ed.

Associate Professor of Orthopaedic Surgery

AND

BARBARA F. HESLOP, M.D. (N.Z.)

Senior Research Officer

*Department of Surgery, University of Otago Medical School,
Dunedin, New Zealand*

[WITH SPECIAL PLATE]

Behaviour of Skin Homografts in Recovered Runts

If all the manifestations of runt disease are attributable—directly or indirectly—to the effects of a graft-versus-host reaction, then the continuance of the condition will obviously be dependent on the continued survival of the injected cells. Recovery from the disease in these circumstances might be expected (1) if the cells or their mitotic progeny were destroyed for any reason, or (2) if the cells were able to persist without immunological attack upon the host. In the first of these two situations the host would cease to be a chimera—that is, to contain cells of two genetically different strains—and the tolerance extended to any donor-strain skin homografts might likewise be expected to come to an end (Billingham and Brent, 1959). In the second situation chimerism and tolerance might persist, both the host and the donor-strain tissues remaining healthy—that is, runt disease might give way to the healthy tolerant state.

The behaviour of skin homografts on recovered runts might thus give some indication of the ways in which runt disease could be terminated other than by death of the host. Unfortunately, a sufficient number of observations of this nature is difficult to obtain for a variety of reasons:

(1) Few runts survive to an age when skin-grafting is practical without a high mortality (four to six weeks). In our own series 18 out of 89 runts survived this length of time.

(2) Even fewer acute runts recover (Billingham *et al.*, 1960). Recovery was seen in 12 of our 89 runts.

(3) It may be difficult to distinguish clinically between recovery from acute runt disease and the supervention of low-grade chronic runt disease (see example cited later).

(4) The presence of acute runt disease may be a difficult clinical diagnosis to make in retrospect, since the immunological runt shows few features which differentiate it from the natural runt seen from time to time among non-injected animals, especially in the less severe variants of the former. When in the throes of an intercurrent infection, even an animal which was previously healthy presents many of the stigmata of runt disease—for example, scruffy skin, temporarily retarded growth, hunched posture, and possibly snuffles and acute diarrhoea. Diagnostic difficulties arising from the lack of specificity of the clinical findings are no great problem in animals in which the progressive nature of the disease can be traced throughout the whole of the downward course, which started shortly after receiving homologous cells. On the other hand, an isolated early episode of sickness is much less easily classifiable in retrospect as a definite case of runt disease.

In our own series, out of 89 animals diagnosed as suffering from runt disease, only 18 survived to an age

suitable for skin-grafting (four to six weeks). Of these, four died within three weeks of grafting—that is, before any definite comment on tolerance was possible for the strain combination. Nine recovered and lost their grafts and two were chronic runts. In one instance the animal presented the classic picture of runt disease at the time of skin-grafting and for a few days afterwards. The condition was not progressive, and within two weeks the animal was noted to be improving clinically, the improvement being maintained over the next three months. Since it had been carrying its skin graft for 100 days after the first sign of clinical recovery, it was tempting to conclude that runt disease had given way to a healthy tolerant state. However, the animal weighed only half that of a healthy litter-mate control and there was retarded development of the facial bones.

Whether retarded growth alone is sufficient evidence for designating this animal a chronic runt as opposed to a recovered runt remains uncertain. By comparison with this case, two animals showed severe runt disease of several weeks' duration. Complete clinical recovery finally occurred, but in both instances the original graft remained perfectly healthy for some months. One of these animals was regrafted from the donor strain, and both grafts remained normal. Here there seems little doubt that runt disease was superseded by the healthy tolerant state (Fig. 7). Nevertheless, in an animal which recovers completely while still retaining its graft there must always remain an element of doubt concerning

FIG. 7.—Healthy skin graft from an inbred black rat on a recovered runt of the inbred Wistar strain.

the nature of the earlier illness. The presence of tolerance makes the diagnosis of runt disease seem more likely in the doubtful clinical situation, but it is evidence of a circumstantial rather than of a confirmatory nature. While the loss of tolerance probably precludes the presence of runt disease (Medawar, 1959b), it is by no means certain whether recovery from runt disease necessarily entails abrogation of the tolerance presumably acquired along with the disease. Oliner *et al.* (1961) note relapse after apparent recovery in two of their runt mice, and suggest that "grafted lymphoid tissue may lie dormant for lengthy periods of time and then for an unknown reason become reactivated and produce disease." Simonsen (1960) interprets recovery from runt disease as indicating acquisition of immunological tolerance by the grafted cells in respect of their host, and

*Substance of the second part of a paper read at the Annual Meeting, and Biennial Conference of the New Zealand Branch, of the British Medical Association, Auckland, 1961. The first part appeared in last week's issue.

N. W. NISBET AND BARBARA F. HESLOP: RUNT DISEASE

FIG. 1.—Lymph node virtually devoid of lymphocytes. There is some sinus proliferation, but many of the collections of cells are perivascular in arrangement. ($\times 60$.)

FIG. 1

FIG. 2.—Lymph-node cortex replaced by plasma cells which are seen to penetrate capsule and extend into surrounding fat. ($\times 50$.)

FIG. 2

FIG. 3

FIG. 3.—Lymph-node cortex largely replaced by histiocytes. ($\times 225$.)

FIG. 4

FIG. 4.—Histiocytes replacing lymph-node cortex. Nuclear debris is seen within one of the cells (arrow). ($\times 450$.)

FIG. 5.—Lymph-node cortex extensively replaced by pale reticulum cells. ($\times 225$.)

FIG. 6.—Lymph-node cortex in normal newborn rat. Relatively few lymphocytes are present, reticulum cells predominating. ($\times 207$.)

C. W. GRAHAM-STEWART: INJECTION TREATMENT OF HAEMORRHOIDS

FIG. 1.—Histological appearances after injection of pure almond oil. (H. and E. $\times 175$.)

FIG. 2.—Histological appearances after injection of 5% phenol in almond oil. (H. and E. $\times 175$.)

FIG. 3.—Radiograph of specimen showing extrusion of oil 12 hours after injection of (1 and 2) 3 ml. and (3 and 4) 1 ml. of neo-hydriol-almond oil. Sites of injection marked with silver wire.

FIG. 4.—Radiograph of specimen to show effects of pre-operative injection of neo-hydriol-almond oil. (1) The carcinoma. (2) The radio-opaque oil.

simultaneous "exhaustive sensitization" on the part of the host.

It would seem that the problem of recovery from runt disease may well be a complex one, and at present does not admit of any simple explanation.

Induction of Runt Disease in Healthy Tolerant Adults

It seems possible that if healthy animals already tolerant of skin from a homologous donor are injected with a further dose of immunologically competent cells from an animal genetically similar to the original donor, then a graft-versus-host reaction might ensue. Indeed, it is surprising that graft-versus-host reactions do not regularly occur in animals made tolerant by adult immunologically competent cells. Castermans (1958) succeeded in inducing a disease reminiscent of runt disease in a proportion of normal adult animals receiving large doses of homologous immunologically competent cells. On the other hand, Billingham and Brent (1959) were unable to produce sickness in healthy tolerant mice by further injections of normal adult spleen cells from the original donor strain.

In our own series an attempt was made to ascertain whether rats highly tolerant of skin from another strain, and presumably unable to react against subsequent cellular injections from the same donor, would develop runt disease (probably better called homologous disease in these circumstances) if reinjected with donor-strain cells sensitized against the recipients. A group of 10

Table showing Results of Injections of 10 Healthy Tolerant Adult Rats with Sensitized Cells Isogenic with the Original Tolerance-conferring Inoculum

Nature of First Injection	Interval between First and Second Injections (Days)	Nature of Second Cell Injection	Effect on Previously Tolerated Graft	Signs of Runt Disease
10 million spleen s.c.	171	142 million lymph-node i.p.	Graft rejected after second injection	None except for temporary episode of hunching at 15 days
10-12 million spleen s.c.	169	160 million lymph-node i.p.	None	None
10 million spleen s.c.	196	55 million lymph-node intracardiac	Graft rejected after second injection	"
10 million spleen s.c.	99	27 million lymph-node i.p.	None	"
10 million spleen s.c.	99	50-100 million lymph-node i.p.	"	Chronic runt disease. Died 78 days after second injection. Involution of lymph nodes
10 million spleen s.c.	99	95 million lymph-node i.p.	"	Acute runt disease. Died 17 days after second injection. Hyperplasia of lymph nodes
10 million spleen s.c.	99	120 million lymph-node i.p.	Graft rejected after second injection	None
10-12 million marrow i.v.	94	99 million lymph-node + 1,120 million spleen i.p.	None	"
10-12 million marrow i.v.	105	140 million lymph-node i.p.	"	Acute runt disease. Died 17 days after second injection. Lymph nodes normal to hyperplastic
10-12 million marrow i.v.	174	306 million lymph-node + 2,800 million spleen i.p.	Graft rejected after second injection	None

s.c. = subcutaneously; i.v. = intravenously; i.p. = intraperitoneally.

C

closed-colony Wistar rats, highly tolerant of skin from either inbred black or inbred hooded strains, were used for this purpose. Groups of four to eight donor-strain rats were sensitized against individual tolerant Wistar rats by two 1 x 1 sq. cm. skin grafts to the chest wall. Fourteen days later the inbred black or inbred hooded rats were killed and cell suspensions from the axillary lymph nodes were injected by either the intraperitoneal or the intracardiac route into the Wistar recipients, which were watched daily. Several of the rats also received large doses of spleen cells from the appropriate donor, the details being shown in the Table.

Six of the rats were quite unaffected, and these included two rats which had received over 2,000 million spleen cells in addition to the sensitized lymph-node cells. One rat developed hunching, together with mild pinkness of the ears, paws, and snout at 15 days, but this passed off and it remained healthy.

Two of the rats died at 17 days. The time interval here between the second cellular injection and death strongly suggested an immunological reaction. The lymph nodes were either normal in size or enlarged. One of these animals acquired pink ears, paws, and snout, but the other showed no definite alteration in colour, though at necropsy the blood and marrow were pale and watery. Apart from some pulmonary oedema, the spleen and other organs appeared normal. Both rats were highly tolerant of donor-strain skin grafts at the time of death.

One rat died of what appeared to be chronic runt disease 78 days after the second injection. It became hunched and developed sores around the mouth and nose. The ears, snout, and paws were definitely pink, and there was considerable epilation of the fur, with a resultant scruffy appearance. At necropsy the lymph nodes were involuted. This rat was likewise highly tolerant at the time of death. The spleen and other organs appeared normal apart from some pulmonary oedema.

The unaffected rats are healthy to date. Three of them have remained highly tolerant for over two years, and a second graft from the original donor strain has been healthy for many months. Following the second cellular injection no fewer than four of the rats rejected the donor-strain graft. This was interpreted as indicating that the animals were in fact partially rather than highly tolerant at the time of the injection. The second cellular injection possibly hastened a process which was already under way, though without chimera tests this suggestion remains entirely speculative. The relationship of the second injection to graft rejection could, of course, have been fortuitous, though this seems a little unlikely. It is of interest to note that Billingham and Brent (1959) observed a similar effect in a fully tolerant mouse reinjected with normal spleen cells from the original donor strain.

Macroscopic Appearances of Runt Disease at Necropsy

Apart from the variable evidence of terminal infection in the respiratory and alimentary systems and the loss of fat, the most striking abnormality has been in the lymphoreticular tissues. Billingham and Brent (1959) originally described severe involution of lymph nodes in mice, generally without increase in size of the spleen, and Russell (1960) presented essentially similar findings. Cole and Ellis (1958) recorded splenic atrophy in mice,

as did Porter (1960) in rabbits after an initial splenomegaly. On the other hand, Simonsen *et al.* (1958) found splenomegaly and hepatomegaly in mice to be a sufficiently constant feature of runt disease to warrant its use as an index of the presence of a graft-versus-host reaction. In rats Billingham *et al.* (1960) reported considerable enlargement of lymph nodes and spleen. Nisbet *et al.* (1960a), on the other hand, noted considerable variation in the size of lymph nodes and spleen in rats. It seems possible that the gross appearances of the lymphoid organs may vary at different stages of the disease. This has been shown to occur in chickens injected with homologous cells before hatching (Biggs and Payne, 1961a, b). Likewise, Simonsen and Jensen (1959) noted a peak of splenic enlargement at seven to ten days in some mouse strains. Gorer and Boyse (1959a) found splenic enlargement and variable lymph-node size early in the disease in mice, though shrinkage and atrophy of both were evident later. Howard (1961) noted an early increase in the size of the spleen, sometimes, but not invariably, followed by terminal atrophy.

In our own series the post-mortem appearances depended on whether the runt was red or pale. In red runts the blood was noticeably thick, and clotted so quickly that as a rule it could not be withdrawn through a hypodermic needle. The bone-marrow was a rich red colour and gelatinous in consistency. All the tissues, especially the muscles, were redder than normal. The lymph nodes and spleen were often enormously enlarged, soft, and brownish in colour. The size of the nodes throughout the body, however, often varied over a wide range, from being barely perceptible to the size of a large pea. This observation applied to lymph nodes in all sites.

Pale runts showed obviously thin and watery blood, which was easily withdrawn through a hypodermic needle. All the tissues appeared grossly anaemic, and the marrow was pale and fibrous, good smears being hard to make. The size of the lymph nodes and spleen was again variable. Often the lymph nodes were quite small and presented the involuted gross appearances

described by other observers (Woodruff and Sparrow, 1957; Billingham and Brent, 1959; Russell, 1960). This, however, was by no means invariable, and they were often enlarged. The size of the spleen was variable and could not always be correlated with the size of the lymph nodes. In both pale and red runts the thymus was usually normal in size and appearance.

The intestines tended to be gelatinous-looking, and on occasions the lumen of the gut was noticed to contain a dark material suggestive of altered blood. Petechial haemorrhages were sometimes seen in the wall of the stomach and in the skin. Other organs, including the liver, were usually normal, but sometimes the adrenals were enlarged. In severe cases of runt disease all the parenchymal organs presented a flabby consistency, which was regarded as being indicative of severe generalized systemic disease. In one animal the collagen, which appeared rather gelatinous, was tested for stretch by the method of Gustavson (1955). Runt collagen—at least in this animal—appeared to be abnormally stretchable, indicating abnormality of the hydrogen bonding and chemical composition. The central nervous system and joints showed no macroscopic abnormality.

Histopathology

Lymph Nodes

In animals receiving homologous cells during the neonatal period and also in adult F1 hybrids receiving parental-strain cells the most regularly reported change has been atrophy of the lymphoid tissues. Lymphoid atrophy need not necessarily be accompanied by a decrease in size of the nodes, though this has sometimes been recorded (Woodruff and Sparrow, 1957; Billingham and Brent, 1959; Porter, 1960; Russell, 1960). In spite of the disappearance of much of the lymphoid tissue, the lymph nodes are often enlarged (Kaplan and Rosston, 1959; Billingham *et al.*, 1960) as a result of the proliferation of other cell types. The nature of the latter cells has differed somewhat and they have been variously recorded as histiocytes (Gorer and Boyse, 1959a; Russell, 1960), reticulum cells and

histiocytes with stem cells and sometimes haemopoiesis in the medullary sinusoids (Billingham *et al.*, 1960), transitional and immature plasma cells (Porter, 1960), plasmablasts and reticulum cells (Howard, 1961), and stem cells with a variety of derivatives (Nisbet *et al.*, 1960a).

In our experience in rats the histological picture was extremely complex, and not necessarily uniform in any one animal, or even in any one node. For instance, it was not uncommon to find gross depletion of lymphoid tissue in one group of nodes while another group from the same animal showed this change to a much less noticeable extent. Again, moderate quantities of bronchial and ileal lymphoid tissue were sometimes seen in animals showing considerable lymphoid depletion of

FIG. 8.—Lymph node showing extensive depletion of lymphocytes and replacement by stem cells and reticulum cells. Some islands of normal lymphoid tissue remain (arrows). ($\times 20$.)

the main nodes. Probably analogous to the patchy nature of these changes was the occurrence of well-demarcated islands of normal tissue in a node which showed gross structural changes elsewhere (Fig. 8). Sometimes the abnormal areas in such a node tended to impart a moth-eaten appearance, the normal areas containing well-demarcated spaces or islands made up of abnormal cell types. In general, uniformity of the histological picture tended to be restricted to the more fulminating cases of runt disease. The area previously occupied by lymphoid tissue sometimes remained relatively empty (Special Plate, Fig. 1), but in other cases the node structure had been replaced by plasma cells and their precursors (Special Plate, Fig. 2), histiocytes, often containing nuclear debris (Special Plate, Figs. 3 and 4), or cells which were probably akin to the reticulum cells or activated reticulum cells (Special Plate, Fig. 5) of Marshall (1956). Pyroninophilic stem cells were almost invariably present irrespective of the type of cell predominating. Occasionally the sinuses contained an excess of iron-containing histiocytes, together with considerable numbers of mature plasma cells, but this appearance was not particularly frequent. Cellular proliferation originating in the sinuses was not generally a feature of the lymph nodes, but sometimes perivascular accumulations of cells gave an appearance superficially reminiscent of sinusoidal proliferation (Special Plate, Fig. 1). In many nodes, however, the initial site of proliferation could not be decided, since the normal node structure was entirely replaced by other cells, often with eventual penetration of the capsule. Very occasional nodes showed small and rather doubtful aggregations of haemopoietic tissue, but no large or very definite islands of haemopoietic tissue were seen within the nodes.

Bronchial and Intestinal Lymphoid Tissue

In our experience many cases of runt disease showed virtually no lymphoid tissue in these sites. However, this was not invariably the case, and sometimes small, or even moderately large, lymphoid nodules were identifiable, very occasionally with germinal centres. Replacement of lymphoid tissue by other cells did not seem to occur very often, though occasional collections of reticulum cells or pyronin-positive stem cells suggested that this may have happened. Plasma cells were not infrequent.

The occurrence in these sites of cells which were almost certainly of inflammatory origin made it difficult to assess the nature of the changes in the lymphoid tissues.

Spleen

Lymphoid atrophy in the spleen has been observed by Billingham and Brent (1959), Billingham *et al.* (1960), Kaplan and Rosston (1959), Porter (1960), and Howard (1961), and has usually been accompanied by infiltration with cells similar to those described in the lymph nodes. Biggs and Payne (1961a) give a detailed account of the histological changes seen in the spleen of chicks receiving homologous adult spleen cells, noting foci of reticulum cells together with aggregations of blast cells and granulocytes. Our own histological findings in the spleens of rats were similar, though, as in the lymph nodes, there was a considerable range of histological variation. The most regular change was a diminution in size of the lymphoid follicles, which had often completely disappeared. Where the Malpighian

bodies were identifiable they tended to be small and inconspicuous, and they were occasionally replaced by pale reticulum cells. In a minority of cases of runt disease the Malpighian bodies were not noticeably diminished in size, and occasionally showed well-defined perifollicular envelopes of Krumbhaar (1948). All the cases of runt disease showed patchy or diffuse aggregations of stem cells or reticulum cells either within the sinusoids or within the connective-tissue framework of the spleen. Histiocytic proliferation was less frequently evident, though sometimes the sinusoidal walls were noticeably thickened. The primitive cells often showed no tendency towards differentiation into mature cell types, but in a few cases collections of plasma cells were seen in the sinusoids, often in association with an excess of iron-containing histiocytes. The number of plasma cells seldom approached that recorded by Howard (1961) in mice, though these differences may be due to the fact that in the present study undifferentiated pyronin-positive cells were designated stem cells and not plasmablasts.

In all the runts showing the red clinical variant there was copious haemopoiesis in the spleen. In some of the cases reported by Nowell and Cole (1959) excessive extramedullary haemopoiesis was recorded within the splenic pulp. It would be interesting to know if these animals showed the clinical redness reported in the present series. Kaplan and Rosston (1959) similarly record increased haemopoiesis in those animals which received spleen cells as the original inoculum. In our own red runts which showed haemopoiesis in the spleen the quantity of haemopoietic tissue in any one section did not appear to be in excess of that seen in controls; but, taken in conjunction with the increase in splenic size and the hyperplasia of the marrow, it seems that there was an over-all increase in the production of blood cells. As in the case of the lymph nodes, it was interesting to observe that—at least in our strains—the histological appearances of the spleen at birth were virtually indistinguishable from those seen in the more acute runts dying at the age of 2 or 3 weeks. The possibility suggests itself that some of the appearances may have been the result of retardation of development. Neither fibrosis nor hyalinization was seen in our series, though either or both are noted by Billingham and Brent (1957), Woodruff and Sparrow (1957), Kaplan and Rosston (1959), Trentin (1959), and Porter (1960). Similarly, the spleens in our series were free of the necrotic lesions seen by Simonsen (1957), Isacson (1959), and Biggs and Payne (1961a) in chicks and by Woodruff and Sparrow (1957) in rats.

Thymus

Relatively few accounts are available of the histology of this organ in runt disease. Billingham *et al.* (1960) recorded thymic atrophy in rats. Porter (1960) found that the lymphocytes of the thymus tended to be spared in runt disease in rabbits, in contrast to the striking loss of lymphoid elements in other lymphoid structures. Kaplan and Rosston (1959) reported focal areas of necrosis at five days in F1 hybrid mice injected with parental-strain lymphoid cells. Simonsen (1957) observed expansion of the thymic medulla in chicks at the expense of the cortex, the appearances ultimately resembling those of the spleen, in which there was considerable infiltration by primitive reticulo-endothelial cells. In our own experience the thymus was normal in the majority of cases where thymic tissue was available

for examination. In view of the severity of some of the terminal infections, it was often surprising to find that thymic tissue was relatively easily identifiable, and that the extent of atrophy was a good deal less than might have been expected in the circumstances. The parathymic lymph nodes were often a source of histological confusion, as these, in common with the other lymph nodes, tended to show gross abnormality. Where the tissue removed from the thymic region contained only parathymic lymphoid tissue it was not always easy to ascertain whether this was thymus showing severe histological changes. Pyronin-stained sections were found to be particularly useful for a quick differentiation of thymus from lymph nodes, since the mast cells, whose granules stained a brilliant orange, were seen to be disposed throughout the lymph nodes. In the thymus they were fairly numerous, but were restricted to the fibrous trabeculae.

It was of interest to note that, despite the histological differences between the normal newborn and adult lymph nodes and spleen, the thymus of newborn rats did not differ noticeably from that of adult controls.

Skin

Billingham *et al.* (1960) describe a severe skin lesion in runt disease in inbred rats, with epithelial thickening and histiocytic infiltration of the dermis. Our own observations confirm the findings in rats, though in general the lesion seen by us was rather less severe than that recorded by these authors.

Liver

The livers of chicks injected with homologous cells have frequently been reported to show necrotic or infiltrative lesions (Simonsen, 1957; Biggs and Payne, 1961a), and mice have likewise been similarly involved (Billingham and Brent, 1959; Gorer and Boyse, 1959a; Siskind and Thomas, 1959; Howard, 1961). Porter (1960) noted focal necrosis in runt disease in rabbits. The rats in our own series were invariably devoid of lesions of this type. Occasionally there were minimal areas of centrilobular necrosis without associated inflammatory reaction. These lesions were sometimes associated with cellular vacuolation elsewhere within the lobules, and they were regarded as being non-specific and terminal lesions which were reasonably explained by the presence of severe infection in the host. Very occasional small islands of haemopoiesis were seen in a few of the red runts.

Respiratory and Alimentary Tracts

Inflammatory lesions of variable extent were evident in our own series. Plasma cells were fairly numerous in both sites. In many cases, however, the extent of the inflammatory lesions was surprisingly small considering the severity of the clinical symptomatology.

Other Tissues

In most cases the remaining tissues were normal, or at most presented only minor histological changes. In spite of the considerable antigenic potential of the nucleated cells of the host, no tissue showed any histological evidence of the type of homograft reaction seen in relation to homografts of skin (Medawar, 1944) and other tissues. In fact lymphocytes, which are the principal cell type in such a homograft reaction, were strikingly absent from the organs of animals with runt disease.

Difficulties in Interpretation of Histological Changes in Lymphoid Tissues

Because each of the following considerations contributes its own difficulties, the problem of interpreting the exact nature of the histological changes seen in the lymphoid tissues in runt disease is an enormous one.

(1) It is not certain which of the cells within the lymph nodes are of donor origin and which belong to the host. By using a chromosome marker Davies and Doak (1960) estimated a minority of the dividing cells in the spleens of runted mice to be of donor origin. Biggs and Payne (1961a), using a similar method in chicks, estimated 50% to be of donor origin. Even allowing that considerable mitotic activity of both donor and host cells is occurring, the exact disposition and behaviour of the two cell types remain uncertain.

(2) By the time the animal dies of runt disease, whatever the design of the experiment, the host should have become at least partially capable of mature response to a variety of antigens, not least among which may be the respiratory and gastro-intestinal organisms, the effects of which are so obvious clinically. Many of the histological appearances of the lymph nodes may merely reflect the response to a variety of antigens, not necessarily related to the original cellular inoculum.

(3) In the rat (Gillman and Gillman, 1952), and probably in other species, normal lymph nodes show a wide range of structural variation. Accordingly, unless a lesion is fairly well developed it may be difficult to decide whether it falls outside the range of normality.

(4) If the clonal selection theory of acquired immunity is correct (Burnet, 1959), then it seems likely that in a given lymphoid organ (a) only a proportion of the host cells will be capable of reacting against a given antigen, and (b) only a proportion of the donor cells will be capable of reacting against the host. This being so, then the distribution of lesions, whether caused by host or donor cell activity, might well be patchy.

(5) In rats—at least in our strains—the lymph nodes contain very few mature lymphocytes at birth, the majority of cells being much larger cells with immature nuclear structure and scanty, fairly pyroninophilic cytoplasm (Special Plate, Fig. 6). Similarly, the spleen shows histological appearances at birth which are very different from those seen in the adult animal, the pulp containing large numbers of stem cells and reticulum cells and the Malpighian bodies being inconspicuous. Thus histological comparisons between runts and normal adult controls may not be entirely valid, at least in those cases where runt disease is induced by injection of homologous cells during the neonatal period. Where the host is very young at the time of inoculation one wonders to what extent the histological appearances can be attributed to retarded development.

(6) In the case of runt disease induced during the neonatal period the introduction of adult cells into a less mature environment might conceivably exert a growth-promoting effect on the cellular inoculum. Tissue-culture experiments by Puck *et al.* (1958) have shown that embryonic calf serum is free from the mitotic inhibitors present in sera from older animals.

(7) It is very probable that the appearances of the lymphoid organs vary according to the stage of the disease (Isacson, 1959; Simonsen and Jensen, 1959; Porter, 1960; Biggs and Payne, 1961a, b). This adds

to the difficulty of comparing individual cases, particularly as the tempo of the disease is variable.

(8) The problem of interpreting some of the less severe changes seen in runt disease would appear to be of the same order as that presented by the notoriously difficult "doubtful lymph node" of human histopathology. Histological opinion may well vary for this reason alone.

Haematology

Simonsen (1957) recorded Coombs-positive haemolytic anaemia in chickens after chorioallantoic inoculation with adult homologous spleen cells. Since then immune haemolysis in runt disease has been described in rabbits by Porter (1960) and in mice by Oliner *et al.* (1961). Haemolytic anaemia, however, does not appear to be an invariable accompaniment of runt disease. Billingham *et al.* (1960) report hypoplastic dyschromic anaemia in rats with runt disease without any mention of abnormal haemolysis. In our own series the haematology of runt disease was not studied systematically. However, in those animals investigated it seemed that at least three different effects on the erythropoietic tissues were manifest:

(1) In a minority of the animals anaemia was accompanied by an increase in the iron-containing histiocytes in the spleen and lymph nodes, together with an associated increase in plasma cells in the sinusoids of the lymph nodes and spleen. While the Coombs test was not undertaken in these animals, there was presumptive evidence of haemolytic anaemia.

(2) In a far greater proportion—in fact, in the remainder of the pale runts—there was no increase in haemosiderin in the tissues. The red-cell count was definitely low, as might have been anticipated from the watery condition of the blood, and the marrow was hypoplastic. Polychromatophilia of the red cells was pronounced as compared with normal controls and there was considerable anisocytosis. Nucleated red cells were seldom increased in the peripheral blood, though Howell-Jolly bodies were not infrequent.

(3) In the red runts the blood was excessively thick and the red-cell count was higher than normal. At first it seemed possible that this was the result of haemoconcentration accompanying dehydration. However, the presence of copious haemopoiesis in the enormous spleens, together with the hyperplastic marrow, made this appear unlikely. It seems possible that this may well be a similar phenomenon to that reported by Nowell and Cole (1959) and Kaplan and Rosston (1959) in F1 hybrids, where the inoculum consisted of homologous spleen cells. The hyperplastic marrow reported by Porter (1960) in some of his runt rabbits may possibly also be akin to these findings.

The white-cell count varied over a considerable range, though this was somewhat less extensive than that recorded in rats by Billingham *et al.* (1960). On the other hand, Kaplan and Rosston (1959), Porter (1960), and Oliner *et al.* (1961) have reported a pronounced decrease in the level of the white blood cells. The appearances described by Billingham *et al.* (1960) as abnormal granulopoiesis were also evident in our own rats. Tissue sections of marrow occasionally showed localized collections of large cells with indented or lobed nuclei, and copious non-granular, faintly pyroninophilic cytoplasm. The identity of these cells was not entirely clear, but it seemed probable that they represented monocytoïd cells rather than abnormal granulocytes, especially in view of the number of similar cells seen elsewhere in the reticular tissues. It was felt that the lobed nucleus alone supplied insufficient grounds for the inclusion of these cells within the granular series. Gorer and Boyse (1959a) recorded collections of histiocytes within the sinusoids of the liver in F1

hybrid mice injected with parental-strain spleen, and noted the resemblance to malignant histiocytoma. Similar collections of cells were seen within a variety of vessels in our own series, though the exact identity of the cells was less clear.

Thrombocytopenia has been recorded by Porter (1960) and by Oliner *et al.* (1961).

Relationship to Autoimmune Disease in Man

While both runt disease in animals and autoimmune disease in man may be said to result from an immunological attack by host cells (or cells which are immunologically or genetically acceptable to the host) against normal body components, in the present state of our knowledge it is difficult to do more than point out certain similarities and differences between the two states. A number of pathological conditions in man have been classified as having an autoimmune component, notably haemolytic anaemia (Dacie, 1959), Hashimoto's disease (Roitt *et al.*, 1956), non-tuberculous Addison's disease (Anderson *et al.*, 1957), myasthenia gravis (Smithers, 1959), lupoid hepatitis (Mackay *et al.*, 1956), and disseminated lupus erythematosus and the so-called collagen diseases (Friou, 1958; Alexander and Duthie, 1958; Burnet, 1959). Inclusion of these conditions within the category of autoimmune diseases has rested mainly on the identification of antibodies which will react against the affected tissues, and to some extent on the histological presence of lymphoid aggregations and on the presence of abnormal globulin patterns in the serum. The response to corticosteroid therapy has also supplied confirmatory evidence of a circumstantial nature. While exact experimental counterparts have been difficult to produce, Oliner *et al.* (1961) point out that a number of the changes seen in acquired haemolytic anaemia in man, and also in such diseases as disseminated lupus erythematosus, are also present in runt disease in mice. They suggest that the disease seen in their animals may be the experimental analogue of these human conditions. One of the main stumbling-blocks, however, in comparisons of this kind lies in the nature of homograft immunity, especially as it affects the lymphoreticular tissues (Gorer, 1960). In the case of runt disease it is reasonably certain that the major immunological attack by the grafted cells is directed against the nuclear histocompatibility antigens of the host. While the nature of homograft immunity in runt disease is not yet elucidated, at least in other situations in the field of homotransplantation, the part played by humoral antibodies, as opposed to cell-bound antibodies, is open to question (Gorer, 1956, 1960; Medawar, 1959c). The type of immunological reaction responsible for runt disease in animals may not necessarily be related to that obtaining in the conditions at present suggested as being autoimmune in man. In addition, there may well be autoimmune conditions in man where the antibody is cell-bound and not easily demonstrable. Until the immunological basis of both the human and experimental conditions is more clearly defined comparisons are likely to remain difficult.

While the association is by no means proved, the analogy between experimental graft-versus-host reactions and lymphoid neoplasms in man, as suggested by Kaplan and Smithers (1959), seems more apt than does the analogy between experimental graft-versus-host reactions and those diseases in man which are at present often regarded as being autoimmune. They quote the following clinical evidence from patients with lymphomas in support

of their concept that these diseases may represent the human analogue of experimental homologous disease: (1) the multiple infections and poor immune reactions; (2) haemolytic anaemia, wasting, and lymphoid destruction; (3) the finding of autoimmune antibodies in the blood of these patients; (4) the similarities of expression of some of the known immunological diseases and of the malignant lymphomas; (5) the association of other known autoimmune disease with the subsequent development of lymphoid tumours in the organs or tissues involved; (6) the relation of the spleen and thymus to these diseases; and (7) the response of some of the "toxic" manifestations of the lymphoid neoplasms to treatment with corticosteroids.

Kaplan and Smithers (1959) suggest that in both conditions wasting, anaemia, and lymphoid depletion may be due to cyto-immunological reactions of lymphoid cells (tumour or homologous) against the usual haemopoietic cells of the host. The "forbidden clone" (Burnet, 1959) may come into being in the host tissues as a result of either neoplastic transformation of host cells or experimental artifice.

The histological appearances seen in lymph nodes in runt disease may frequently be reminiscent of those observed in some of the reticular neoplasms in man. For example, the appearances seen in Special Plate, Fig. 5, would almost certainly have been reported as reticulum-cell sarcoma had they occurred in the course of routine examination of human biopsy material. Gorer and Boyse (1959a) have likewise noted lesions resembling histiocytomas in some of their F1 hybrid mice injected with parental-strain lymphoid cells. In this connexion, however, it is important to remember that a limited number of morphological responses may result from a very wide variety of stimuli. Because of this it is hazardous to conclude that similar morphological end-results necessarily imply the operation of similar initial stimuli.

Some Unexplained Aspects of Runt Disease

In spite of widespread interest in the different manifestations of graft-versus-host reactions, an impressive array of questions remains unanswered. The exact relationship to tolerance is not entirely clear. While there is no doubt that tolerance can occur without runt disease, the possibility of runt disease occurring in the absence of tolerance cannot be wholly dismissed. The type of strain combination which yields a high degree of runting in the absence of tolerance will be familiar to most who have worked in this field, and the apparently narrow dividing-line between normality and runting in these circumstances does not admit of easy explanation.

The variability in response between litter mates exposed to apparently identical stimuli is always puzzling. While it seems likely that differences in immunological maturity may explain extreme differences in results, gross variations in response are nevertheless surprising in apparently similar members of an inbred strain. In our own series newborn rats, often from the same litter and injected with similar doses of homologous cells, tended to show extreme variability of response, which included sensitization, healthy tolerance, and runt disease. It seems that at least a proportion of these rats are more mature at birth than the results of Woodruff and Simpson (1955) would indicate.

The part played by genetic differences in influencing the readiness with which tolerance is obtainable is like-

wise largely a mystery. It remains to be seen whether the effects of strong and weak antigens result from quantitative or qualitative differences.

The manner in which the histopathological changes in runt disease are brought about is unexplained, as is the nature of the immunological reaction against lymphoid cells (Gorer, 1960). The conventional histological homograft reaction such as is seen in relation to skin grafts is not in evidence, though all the host tissues must be regarded as antigenic. In fact a number of host tissues appear completely to escape specific damage as a result of the graft-versus-host reaction.

The disappearance of the lymphocytes of both donor and host is unexplained. Since the lymphocyte is not only the most prominent cell in the homograft reaction to soft-tissue grafts (Medawar, 1944) but is also believed to be the cell responsible for the graft-versus-host immunological reaction (Burnet and Burnet, 1960), it is surprising to find a complete absence of graft lymphocytes as well as host lymphocytes in most cases of runt disease. Gorer and Boyse (1959b) believe that this phenomenon is explicable on a basis of allergic death of the graft cells.

The exact extent to which the changes seen in runt disease and allied conditions can be attributed to immunological attack on the host remains obscure. While there seems little doubt that an immunological reaction by the grafted cells does occur, nevertheless a variety of other factors, not necessarily immunological, may also contribute to the final outcome. These problems have recently been reviewed by Van Alten and Fennell (1959).

The state of our knowledge in this field has been aptly described by Billingham *et al.* (1960), who comment: "At present, however, it is probably fair to say that we are still like archeologists trying to determine from old relics on an ancient battleground exactly who the real contestants were, exactly how they fought, and what happened to them."

Summary

The demonstration that it was possible to induce immunological tolerance of homotransplantation antigens in experimental animals at first seemed to indicate that the solution of the homograft problem in man might be in sight. The available methods of inducing immunological tolerance in animals, however, led to a considerable mortality from runt disease, and any hope of applying these methods to man had of necessity to be abandoned. At about the same time a similar disease entity, secondary or homologous disease, was noted to occur as a complication of the treatment of irradiated animals with homologous lymphoid cells. In both situations it appeared that the pathogenesis was similar. The host was incapable of reacting against the homologous cells by reason either of immaturity or of radiation damage to the reticular tissues. The homologous inoculum, however, contained immunologically competent cells which were capable of reacting against the host. Both diseases were thus believed to be the outcome of graft-versus-host reactions. Though no exact human counterpart exists, runt disease may well be of interest in clinical medicine in so far as it represents a situation equivalent to autoimmunity in man—that is, cells regarded by the host as "self" are able to react against the host.

In the present paper the conditions necessary for the induction of runt disease are described, as are the clinical and pathological features. While there are as yet many aspects which admit of no ready explanation, runt disease promises to be an important tool in a wide variety of clinical and experimental fields.

We wish to acknowledge the assistance of our colleague, Dr. Irmgard Zeiss, during the initial stages of this study. We are indebted to Dr. W. I. McDonald, who undertook the neuropathology, and to Dr. B. McMahon for the haematology. The testing of runt collagen was carried out by Dr. T. C. Highton, and statistical analysis of results by Mr. G. F. S. Spears. The graphs were prepared by Miss L. Duncan, and the photographs by Mr. D. Weston. We are indebted to Mr. I. Houston, Miss L. Elmes, and Miss N. Richardson for technical and secretarial assistance.

REFERENCES

- Alexander, W. R. M., and Duthie, J. J. R. (1958). *Brit. med. J.*, **2**, 1565.
- Anderson, J. R., Goudie, R. B., Gray, K. G., and Timbury, G. C. (1957). *Lancet*, **1**, 1123.
- Anderson, N. F., Delorme, E. J., and Woodruff, M. F. A. (1960). *Transplant. Bull.*, **7**, 93.
- Barnes, D. W. H., Ford, C. E., Ilbery, P. L. T., Koller, P. C., and Loutit, J. F. (1957). *J. cell. comp. Physiol.*, **50**, Suppl. 1, 123.
- Berrian, J. H., and Jacobs, R. L. (1959). In *Biological Problems of Grafting*, ed. F. Albert, p. 131. Blackwell, Oxford.
- Biggs, P. M., and Payne, L. N. (1961a). *Immunology*, **4**, 24.
- (1961b). *Ibid.*, **4**, 38.
- Billingham, R. E., and Brent, L. (1957). *Transplant. Bull.*, **4**, 67.
- (1959). *Phil. Trans. B.*, **242**, 439.
- and Medawar, P. B. (1953). *Nature (Lond.)*, **172**, 603.
- (1956). *Phil. Trans. B.*, **239**, 357.
- Brown, J. B., Defendi, V., Silvers, W. K., and Steinmuller, D. (1960). *Ann. N.Y. Acad. Sci.*, **87**, 462.
- and Silvers, W. K. (1960). *J. Immunol.*, **85**, 14.
- Boyse, E. A. (1959). *Immunology*, **2**, 170.
- Burnet, F. M. (1959). *The Clonal Selection Theory of Acquired Immunity*. Cambridge Univ. Press.
- and Burnet, D. (1960). *Nature (Lond.)*, **188**, 376.
- and Fenner, F. (1949). *The Production of Antibodies*. MacMillan, Melbourne.
- Castermans, A. (1958). *Transplant. Bull.*, **5**, 381.
- Cole, L. J., and Ellis, M. E. (1958). *Science*, **128**, 32.
- Dacie, J. V. (1959). *Proc. roy. Soc. Med.*, **52**, 443.
- Danchakoff, V. (1916). *Amer. J. Anat.*, **20**, 255.
- Davies, A. J. S., and Doak, S. M. A. (1960). *Nature (Lond.)*, **187**, 610.
- Dempster, W. J. (1953). *Brit. J. Surg.*, **40**, 447.
- Eichwald, E. J., and Lustgraaf, E. C. (1958). *Ann. N.Y. Acad. Sci.*, **73**, 777.
- , Fuson, R. B., and Pfaff, J. P. (1960). *Ibid.*, **87**, 119.
- Finerty, J. C. (1952). *Physiol. Rev.*, **32**, 277.
- Fowler, R., and West, C. D. (1960). *Transplant. Bull.*, **26**, 133.
- Friou, G. J. (1958). *Ann. intern. Med.*, **49**, 866.
- Gillman, J., and Gillman, T. (1952). *Cancer*, **5**, 792.
- Gorer, P. A. (1956). *Advanc. Cancer Res.*, **4**, 149.
- (1960). In *Cellular Aspects of Immunity*, ed. G. E. W. Wolstenholme and M. O'Connor, p. 330. Churchill, London.
- and Boyse, E. A. (1959a). *Immunology*, **2**, 182.
- (1959b). In *Biological Problems of Grafting*, ed. F. Albert, p. 193. Blackwell, Oxford.
- Gowans, J. L. (1957). *Brit. J. exp. Path.*, **38**, 67.
- Gustavson, K. H. (1955). *Nature (Lond.)*, **175**, 70.
- Hicken, P., and Krohn, P. L. (1960). *Proc. roy. Soc. B*, **151**, 419.
- Hildemann, W. H. (1956). *Transplant. Bull.*, **3**, 144.
- Howard, J. G. (1961). *Brit. J. exp. Path.*, **42**, 72.
- Isacson, P. (1959). *Yale J. Biol. Med.*, **32**, 209.
- Kaplan, H. S., and Rosston, B. H. (1959). *Stanford med. Bull.*, **17**, 77.
- and Smithers, D. W. (1959). *Lancet*, **2**, 1.
- Krumbhaar, E. B. (1948). *Blood*, **3**, 953.
- Mackay, I. R., Taft, L. I., and Cowling, D. C. (1956). *Lancet*, **2**, 1323.
- Mannick, J. A., Lichte, H. L., Ashley, C. A., Thomas, E. D., and Ferrebee, J. W. (1959). *Surgery*, **46**, 821.
- Marshall, A. H. E. (1956). *An Outline of the Cytology and Pathology of the Reticular Tissue*. Oliver and Boyd, Edinburgh.
- Medawar, P. B. (1944). *J. Anat. (Lond.)*, **78**, 176.
- (1959a). In *Cellular and Humoral Aspects of the Hypersensitive States*, ed. H. S. Lawrence, p. 504. Cassell, London.
- (1959b). In *Biological Problems of Grafting*, ed. F. Albert, p. 188. Blackwell, Oxford.
- (1959c). *Ibid.*, p. 6.
- (1961). *Science*, **133**, 303.
- Mitchison, N. A. (1956). *Brit. J. exp. Path.*, **37**, 239.
- Murphy, J. B. (1916). *J. exp. Med.*, **24**, 1.
- Murray, J. E., Merrill, J. P., Dammin, G. J., Dealy, J. B., Walter, C. W., Brooke, M. S., and Wilson, R. E. (1960). *Surgery*, **48**, 272.
- Nisbet, N. W., Heslop, B. F., and Zeiss, I. M. (1960a). *Proc. Univ. Otago med. Sch.*, **38**, 11.
- (1960b). *Brit. J. exp. Path.*, **41**, 443.
- Nowell, P. C., and Cole, L. J. (1959). *Transplant. Bull.*, **6**, 435.
- Oliner, H., Schwartz, R., and Dameshek, W. (1961). *Blood*, **17**, 20.
- Owen, R. D. (1945). *Science*, **102**, 400.
- Porter, K. A. (1960). *Brit. J. Cancer*, **14**, 66.
- and Calne, R. Y. (1960). *Transplant. Bull.*, **26**, 458.
- Puck, T. T., Cieciura, S. J., and Robinson, A. (1958). *J. exp. Med.*, **108**, 945.
- Roitt, I. M., Doniach, D., Campbell, P. N., and Hudson, R. V. (1956). *Lancet*, **2**, 820.
- Russell, P. S. (1960). *Ann. N.Y. Acad. Sci.*, **87**, 445.
- Simonsen, M. (1953). *Acta path. microbiol. scand.*, **32**, 36.
- (1957). *Ibid.*, **40**, 480.
- (1960). *Ann. N.Y. Acad. Sci.*, **87**, 382.
- and Engelbreth-Holm, J., Jensen, E., and Poulsen, H. (1958). *Ibid.*, **73**, 834.
- and Jensen, E. (1959). In *Biological Problems of Grafting*, ed. F. Albert, p. 214. Blackwell, Oxford.
- Siskind, G. W., and Thomas, L. (1959). *Ibid.*, p. 176.
- Smithers, D. W. (1959). *J. Fac. Radiol. (Lond.)*, **10**, 3.
- Snell, G. D. (1957). *Annu. Rev. Microbiol.*, **11**, 439.
- Terasaki, P. I. (1959). *J. Embryol. exp. Morph.*, **7**, 409.
- Trentin, J. J. (1959). In *Biological Problems of Grafting*, ed. F. Albert, p. 207. Blackwell, Oxford.
- Van Alten, P. J., and Fennell, R. A. (1959). *J. Embryol. exp. Morph.*, **7**, 459.
- Van Bekkum, D. W., Vos, O., and Weyzen, W. W. (1959). In *Biological Problems of Grafting*, ed. F. Albert, p. 292. Blackwell, Oxford.
- Vos, O., de Vries, M. J., Collenteur, J. C., and Van Bekkum, D. W. (1959). *J. nat. Cancer Inst.*, **23**, 53.
- Woodruff, M. F. A. (1961). *J. roy. Coll. Surg. Edinb.*, **6**, 75.
- and Simpson, L. O. (1955). *Brit. J. exp. Path.*, **36**, 494.
- and Sparrow, M. (1957). *Transplant. Bull.*, **4**, 157.
- Zeiss, I. M., Nisbet, N. W., and Heslop, B. F. (1960). *Brit. J. exp. Path.*, **41**, 345.
- Zukoski, C. F., Jackson, B. T., Egdahl, R. H., and Hume, D. M. (1960). *Fed. Proc.*, **19**, 358.

INJECTION TREATMENT OF HAEMORRHOIDS*

BY

C. W. GRAHAM-STEWART, M.B., F.R.C.S.
St. Thomas's Hospital, London

[WITH SPECIAL PLATE]

The changes in the tissues which are caused by the injection of phenol have been described by Dukes (Anderson and Dukes, 1924). I have no knowledge of this work being repeated since Dukes's experiments, but in the meantime a drastic change has occurred in the nature of the fluid injected in the treatment of piles. In Dukes's cases either 10% or 20% phenol in glycerin and water was used, whereas now 5% phenol in almond oil is the sclerosant most often employed in Great Britain. It should be appreciated that the fundamental difference between the two solutions is that the glycerin-and-water solution is miscible with water, and thence with tissue fluid, whereas the oily solution is not.

The histological changes caused by 10% or 20% phenol in glycerin and water in Dukes's series are now compared, in the small series here reported, with the changes caused by 5% phenol in almond oil, which are in turn compared with the effects of the injection of pure almond oil.

In three cases of carcinoma of the rectum 2 or 3 ml. of 5% phenol in almond oil and pure almond oil have been injected at different sites at 6, 9, and 12 days before abdomino-perineal resection of the rectum. After

*From a section of a thesis submitted for the M.S. degree of London University.