nausea, vomiting, or diarrhoea was observed. This is of particular importance because the tetracycline group of antibiotics, when taken orally, may cause diarrhoea.

It is worth noting that 14 ears had bulging tympanic membranes yet no myringotomy was found necessary during the entire series. Furthermore, no defect of hearing was detected in these ears when reviewed some months after treatment was completed.

Staph. aureus was isolated from the discharge of 70% of those cases presenting with a perforated tympanic membrane. This agrees with the experience of Morrison (1961), who found that the pyogenic staphylococcus was responsible for 66% of the discharging ears in his series. Of the 30 ears in which the offending organism was isolated, five recovered in spite of the fact that the organisms were resistant, by *in vitro* studies, to the antibiotic given. Treatment was continued in spite of the adverse bacteriological report, as the clinical response of the patient was regarded as adequate. This raises the question of the need to give antibiotics in every case of acute suppurative otitis media. Fry (1958) believes that most cases of otitis media will recover without antibiotic therapy. An alternative explanation is that *in vitro* studies may not give conclusive evidence of the sensitivity of the infecting organism in the human body.

The response of the patients, judged by the return of the temperature to normal and by the disappearance of pain and discharge, was somewhat slower to phenethicillin than to penicillin G. However, all cases treated were afebrile and free from otorrhoea and earache after five days' treatment.

The impression gained from observing the response of these patients to phenethicillin is a most favourable one. It would appear to be the treatment of choice in acute otitis media in childhood because it includes the advantages of adequate blood levels obtained by oral administration, with none of the side-effects on the intestinal tract common to other oral antibiotics. Penicillin G would appear to be preferable to phenethicillin in acute suppurative otitis media with mastoiditis because it produces a slightly quicker response.

Summary

A therapeutic trial was conducted to compare the response of children with acute suppurative otitis media to treatment with phenethicillin and with penicillin G.

All cases responded adequately except for two who had indrawn tympanic membranes, but both had normal hearing after treatment was completed.

The response to phenethicillin in the dosage given would appear to be slower than to parenteral penicillin G, but it has the advantage over penicillin G in that oral administration gives satisfactory blood levels without side-effects on the intestinal tract.

I am indebted to Mr. D. H. Craig and to Mr. R. S. McCrae for permission to carry out this trial on their patients, and to Mr. McCrae for much help in conducting the trial and in the preparation of the paper. My thanks are also due to Sister Lavelle for her co-operation in the care of these young patients. The phenethicillin was kindly supplied by Beecham Research Laboratories Ltd.

BIBLIOGRAPHY

- Asay, L. D., and Hartman, G. L. (1956). *J. Pediat.*, **49**, 565.
Daggett, W. I. (1942). *J. Laryng.*, **57**, 427.
Friedman, I. (1957). *Proc. roy. Soc. Med.*, **50**, 406.
Fry, J. (1958). *Brit. med. J.*, **2**, 883.

- Garrod, L. P. (1945). *Ibid.*, **1**, 107.
— (1952). *Lancet*, **1**, 400.
— (1960). *Brit. med. J.*, **1**, 527.
Hodgkin, K. (1957). *Lancet*, **2**, 514.
Johnston, C. M. (1960). *Brit. med. J.*, **1**, 1493.
Knudsen, E. T., and Rolinson, G. N. (1959). *Lancet*, **2**, 1105.
Lewis, R. S., Gray, J. D., and Hewlett, A. B. (1952). *J. Laryng.*, **66**, 142.
Mawson, S., Gray, J. D., and Hewlett, A. B. (1953). *Brit. med. J.*, **1**, 817.
Morrison, A. W. (1961). *Ibid.*, **2**, 8.
Walker, S. H. (1954). *J. Pediat.*, **44**, 50.
Young, A., and Hall, I. S. (1948). *J. Laryng.*, **62**, 551.

DIAGNOSIS OF AMOEBIC COLITIS ON ROUTINE BIOPSIES FROM RECTUM AND SIGMOID COLON

BY

T. A. McALLISTER, M.B., Ch.B.

Senior House Officer (now Registrar), University Department
of Pathology, Western Infirmary, Glasgow

[WITH SPECIAL PLATE]

In the United Kingdom idiopathic ulcerative colitis is one of the commonest causes of recurrent diarrhoea with the passage of blood and mucus in the stools. It can be firmly diagnosed only after exclusion of other conditions such as carcinoma of the colon and the specific dysenteries. Clinicians and bacteriologists are only too well aware of the prevalence of the bacillary dysenteries, and accordingly thorough bacteriological examination is the rule in all suspected cases. Amoebic dysentery is relatively rare, and consequently there is always a danger that the exacting procedures needed for its diagnosis, requiring the co-operation of clinician, nursing staff, and bacteriologist, will not be carried out efficiently. The present paper shows that in a general hospital the routine parasitological investigation of patients with this syndrome is sometimes inadequate to distinguish cases of amoebiasis from non-specific ulceration. The following three cases demonstrate that the histopathologist can on occasion identify or strongly suspect intestinal amoebiasis by examining fixed biopsy specimens from the rectum or sigmoid colon in patients thought to have idiopathic ulcerative colitis.

Case Reports

Case 1

A man aged 53 presented in May, 1960, with a two-months history of severe diarrhoea accompanied by rectal pain and the passage of blood and mucus. He had a poor appetite and had lost 2 stones (12.7 kg.) in weight. Neither he nor his wife had ever travelled outside the West of Scotland; at a later date he recalled that they had both drunk from a horse-trough at Lanark while walking in the country in October, 1959.

He was pale, thin (weight 111 lb. or 50.4 kg.), and slightly febrile, and apart from slight tenderness over the sigmoid colon there was no other physical abnormality. Haematological examination revealed: Hb 11 g. per 100 ml.; W.B.C. 26,700 per c.mm. (neutrophil polymorphs 82%); E.S.R. 62 mm. The stools were unformed and contained blood, mucus, and pus, but repeated bacteriological examination failed to demonstrate any pathogenic intestinal bacteria. Sigmoidoscopy revealed an oedematous mucosa with small bleeding ulcers, but biopsy on two occasions showed a normal mucosa. Barium-enema examination suggested ulcerative colitis of the sigmoid colon, and accordingly treatment with penicillin, succinylsulphathiazole,

and retention enemata of hydrocortisone hemisuccinate was started. After one week prednisolone by mouth was added, but, following an initial improvement, his condition deteriorated and corticosteroid therapy was tailed off. One month later the findings were: weight 97 lb. (44 kg.); Hb 9.5 g. per 100 ml.; W.B.C. 27,500 per c.mm. Radiologically, the whole of the large bowel now showed severe ulcerative colitis, and preparations were made for total colectomy. Fortunately, further sigmoidoscopy and biopsy were carried out, and structures with the morphology of entamoebae and containing ingested erythrocytes were identified (Special Plate, Figs. 1 and 2). Bacteriological examination of further fresh warm stools confirmed the presence of many motile *Entamoebae histolyticae*.

Intensive treatment with successive courses of penicillin, sulphaguanidine, emetine bismuth iodide, diiodohydroxyquinoline, and carbarsone produced a rapid return to health. He was last seen in March, 1961, when he was symptom-free and weighed 124 lb. (56.2 kg.).

In early 1960 the patient's wife suffered from transient bouts of diarrhoea with blood in the stools. However, these subsided spontaneously, and when stools were examined in October, 1960, there was no evidence of amoebae or cysts.

This case presents two other facets of interest: (a) it is only the second reported case of indigenous amoebic dysentery in Scotland (Conway and Watt, 1961); and (b) it supports the suggestion that local and systemic treatment with corticosteroids may cause unmasking of latent amoebic infection or exacerbation of a mild phase (Mody, 1959).

Case 2

This patient, a man aged 34, who had been in the Forces in India and Burma in 1945, was first seen in October, 1960, as an out-patient with a six-months history of bouts of epigastric pain and diarrhoea characterized by the presence of mucus and blood. Each bout lasted four to five days and there were remissions of one to two weeks. He had no history of dysentery while abroad, but in 1955 he had a pericolic abscess drained. Unfortunately, no details of this are available. He had also had a haemorrhoidectomy in 1959. Examination revealed tenderness in the left iliac fossa and a granular proctitis. Blood examination showed: Hb 13.6 g. per 100 ml.; P.C.V. 44%; E.S.R. 3 mm. The appearances on sigmoidoscopy were typical of ulcerative colitis in the quiescent phase.

Histological examination of a biopsy specimen showed no evidence of ulceration, but an increase of lymphocytes, plasma cells, and eosinophils in the lamina propria. The outstanding feature, however, was the presence in the mucinous surface layer of many large basophilic mononuclear cells containing ingested erythrocytes—that is, the morphological appearances of vegetative *E. histolyticae* (Special Plate, Fig. 3). He was admitted to hospital, and the relevant findings were: E.S.R. 7 mm.; eosinophils 720 per c.mm.; Wassermann reaction negative; no occult blood in faeces. Repeated examination of the faeces failed to reveal amoebae, but in view of the histopathological diagnosis he was given a course of emetine followed by diiodohydroxyquinoline. A further sigmoidoscopy two months later showed a normal mucosa to the recto-sigmoid junction. Since the completion of treatment the patient has remained symptom-free and is well.

Case 3

A male Nigerian student aged 29, who had been in the United Kingdom only one month, presented with epigastric and rectal pain accompanied by diarrhoea containing blood and mucus, all apparently of one month's duration. Clinical examination revealed no abnormality, and rectal examination confirmed the nature of the stool. No intestinal pathogens and no evidence of amoebae were seen on routine bacteriological examination of faeces. Sigmoidoscopy showed a friable, oedematous, and haemorrhagic mucosa,

and a biopsy specimen revealed widespread infiltration of the lamina propria by inflammatory cells with numerous well-formed crypt abscesses. In addition, the mucosal surface exhibited many structures which, from experience of the earlier cases, were recognized as *E. histolyticae*. Their identity was confirmed by examination of a fresh stool on a warm stage. A course of emetine was started, but within ten days severe pain in the right iliac fossa necessitated removal of an acutely inflamed appendix, in which, however, amoebae were not detected. Recovery from this operation was uneventful, and when seen two months later the patient was well, sigmoidoscopy then revealed no abnormality.

Recently we have seen a South African girl aged 13 in whom very scanty bodies similar to those described above were present on the mucosal surface of a rectal biopsy. She had previously been fully investigated for amoebic dysentery in another hospital with negative results, and consequently was under treatment for ulcerative colitis. In view of the histological finding the faeces were again most thoroughly examined for amoebae, but without success. Nevertheless, as a further safeguard she was given a therapeutic trial with emetine. Her clinical condition improved markedly, but after only three weeks she returned to South Africa. No further follow-up studies are as yet available.

Discussion

Amoebic dysentery in a person who has never been outside the United Kingdom is a clinical curiosity—in all about 40 cases are recorded—yet the incidence of symptomless carriers or cyst-passers (Manson-Bahr, 1943, 1954) in the population is relatively high. These cyst-passers can be divided into two groups: (a) convalescent cases of amoebic dysentery; and (b) healthy, symptom-free individuals who have never had a clinical amoebic infection. Estimates of the latter group vary according to the methods used, and range between 1.6% (Morton *et al.*, 1951), 2.7% (Dobell, 1921), and 4.7% (Matthews and Smith, 1919) for persons who had never left the United Kingdom, and 8.8% in R.A.F. personnel recently returned from overseas. Adams (1960) estimates the proportion of indigenous symptomless carriers as "about 5%." The variety of figures quoted is probably related to the different home backgrounds and standards of hygiene in each group (Morton *et al.*, 1951) and to the intensity of the search for cysts. Both Dobell (1921) and Matthews and Smith (1919) examined only one specimen of stool, and the former has shown that this will reveal only one-third of infections. Thus the incidence of symptomless carriers in the general population, which includes increasing numbers of immigrants, is probably about 10%, which is the figure quoted by Craig (1944) for the United States. It is interesting to note that Morton discovered evidence of amoebic colitis in 26 of 144 symptomless carriers examined sigmoidoscopically.

The various methods used for the detection and recognition of vegetative and cystic forms of *E. histolyticae* were discussed and evaluated by Faust *et al.* (1939) and need not be discussed here. In view of the high incidence of symptom-free cyst-excretors, the demonstration of cysts in a patient with colitis does not establish the diagnosis; only the identification, by an experienced observer, of the vegetative amoeba can make the diagnosis certain. In some instances even the most careful examination may fail to demonstrate amoebae, and a therapeutic trial with amoebicidal drugs may then be necessary. In the United Kingdom failure to

establish the diagnosis is likely to arise where the condition is not suspected, and either the stools are not examined for amoebae or insufficient care is taken over the collection and handling of specimens. This was so in Case 1. The diagnosis in each of the cases was first suggested by the finding of large rounded mononuclear cells, containing ingested erythrocytes, lying in the tissues, and in two cases in the surface exudate. These cells had the morphology of vegetative *E. histolyticae*, and in spite of their large numbers could conceivably be confused with large macrophages showing erythrophagocytosis. Thus the discovery of these cells in a routine rectal biopsy does not furnish a conclusive diagnosis, but it at least indicates the need for a rigorous examination of the freshly passed stool, and, should this fail to reveal amoebae, a therapeutic trial with anti-amoebic drugs similar to that suggested by Paulley (1961) for cases of liver abscess of uncertain aetiology.

It is known that amoebic dysentery may masquerade as idiopathic ulcerative colitis; for example, in addition to the three cases now reported, Paulley mentioned that he had seen six, and Passarelli (1960) described *E. histolyticae* in five of 18 cases diagnosed as ulcerative colitis. Manson-Bahr (1943) reported the catastrophic results of surgery in the 1933 Chicago outbreak of amoebic dysentery, and Guthrie (1956) recorded the development of fatal amoebiasis cutis after surgery in an unsuspected case. Medical treatment with corticosteroids appears to be dangerous.

In an attempt to avoid further instances of delay in diagnosis the following recommendations are made:

(a) Rigorous bacteriological examination of all cases of ulcerative colitis, including microscopical examination of the freshly passed stool; examination of samples two to three hours old is of no value.

(b) An extensive search of biopsy material from cases of "ulcerative colitis" for morphologically identifiable vegetative *E. histolyticae* in the mucosa and submucosa, and particularly in the surface exudate. Recognition of amoebae is greatly facilitated by the use of the periodic-acid-Schiff method, by which they are coloured bright red.

(c) If only (b) above is positive, or if there remains doubt about the diagnosis, a therapeutic trial with amoebicidal drugs should be instituted. The toxic effects of emetine, for example, have been much exaggerated, and in practice the drug can be given in therapeutic doses without concern (Adams, 1960). The small risk of toxic effects is far outweighed by the possible benefits. If administration of these drugs resulted in preventing even one unnecessary colectomy per annum in a general hospital it would be worth while.

Summary

Three cases of amoebic dysentery initially thought to be examples of ulcerative colitis are described and discussed. In each case the diagnosis was indicated by histological examination of fixed biopsy material from the rectum or sigmoid colon. The importance of differentiating between amoebic and ulcerative colitis is stressed, and to facilitate this the following recommendations are made: (a) more rigorous protozoological examination of the stools in all cases of ulcerative colitis to exclude amoebic infection; (b) thorough search of the tissues and exudate of routine rectal and sigmoid biopsies for vegetative *E. histolyticae*; and (c) more frequent use of a therapeutic trial with amoebicidal drugs where indicated.

I am indebted to Dr. A. J. Watson, of the Department of Pathology, University of Durham, for assistance in the

preparation of this paper. I am grateful also to Professors E. J. Wayne and C. F. W. (now Sir Charles) Illingworth and Messrs. A. D. Roy and D. H. Clark for permission to publish clinical details, and to Mr. G. Kerr for photography.

REFERENCES

- Adams, A. R. D. (1960). *Brit. med. J.*, **1**, 956.
 Conway, H., and Watt, D. A. L. (1961). In the press.
 Craig, C. F. (1944). *The Etiology, Diagnosis and Treatment of Amebiasis*. Williams and Wilkins, Baltimore.
 Dobell, C. (1921). *Spec. Rep. Ser. med. Res. Coun. (Lond.)*, No. 59.
 Faust, E. C., Sawitz, W., Tobie, J., Odom, V., Peres, C., and Lincicome, D. R. (1939). *J. Parasit.*, **25**, 241.
 Guthrie, W. (1956). Report to the Association of Clinical Pathologists, Edinburgh.
 Manson-Bahr, P. H. (1943). *The Dysenteric Disorders*, 2nd ed. Cassell, London.
 — (1954). *Manson's Tropical Diseases*, 14th ed. Cassell, London.
 Matthews, J. R., and Smith, A. M. (1919). *Ann. trop. Med. Parasit.*, **12**, 349, 361; **13**, 91.
 Mody, V. R. (1959). *Brit. med. J.*, **2**, 1399.
 Morton, T. C., Neal, R. A., and Sage, M. (1951). *Lancet*, **1**, 766.
 Passarelli, N. (1960). *Arch. bras. Med.*, **50**, 55.
 Paulley, J. W. (1961). *Brit. med. J.*, **1**, 462.

BIOPSY OF THE RECTUM AS AN AID TO THE DIAGNOSIS OF AMYLOIDOSIS

BY

P. H. FENTEM, M.Sc., M.B., Ch.B.
University Demonstrator in Pathology

L. A. TURNBERG, M.B., Ch.B., M.R.C.P.
Senior House Officer in Medicine

AND

K. G. WORMSLEY, M.D., B.S., B.Sc., M.R.C.P.
Senior Medical Registrar
Manchester Royal Infirmary

[WITH SPECIAL PLATE]

The examination of biopsy material is the most conclusive way of establishing a diagnosis of amyloidosis (Rukavina *et al.*, 1956). The primary form of amyloidosis so often affects the alimentary tract (Symmers, 1956; Thingstad, 1951) that biopsy of some part of the intestine might be expected to produce tissue containing the characteristic material, while in secondary amyloidosis the high incidence of intestinal involvement (in 55% of cases according to Dahlin, 1949) has been re-emphasized by Gafni and Sohar (1960), who used rectal biopsy to establish the diagnosis in 25 cases.

From the clinical material of a general hospital it has been apparent that, in patients suspected of having amyloidosis, useful information can be acquired by rectal biopsy. The purpose of this paper is to report the cases in which we have used this procedure to confirm the diagnosis of amyloid disease.

Method

A modified Truelove-Salt suction-biopsy instrument (Truelove *et al.*, 1955) was used to obtain biopsy specimens of the rectal mucosa. No special preparation of the bowel was required before instrumentation, but the presence of haemorrhagic disease was excluded by determining the platelet count and prothrombin time.

T. A. McALLISTER: DIAGNOSIS OF AMOEBIC COLITIS

FIG. 1.—Case 1. Vegetative *Entamoeba histolytica* containing ingested erythrocytes. These lie near surface of small ulcer. Note polymorphs in surrounding tissues. (H. and E. $\times 445$.)

FIG. 2.—Case 1. Vegetative *Entamoeba histolytica* in surface exudate. (P.A.S. $\times 1,005$.)

P. H. FENTEM *ET AL.*: DIAGNOSIS
OF AMYLOIDOSIS

FIG. 3.—Case 2. Vegetative *Entamoeba histolytica* in surface exudate. (H. and E. $\times 445$.)

Histological section of tissue obtained from Case 1, showing rectal submucosa with amyloid material in walls of small arterioles and venules stained by congo red (indicated by arrows). ($\times 120$.)