part, together with the progressive and scientific development of the barbiturates and other anaesthetic and analgesic agents, in the recent remarkable advance in the methods of producing anaesthesia, muscular quiescence, and adjustment of the blood pressure, to meet the needs of surgery. It is obvious, I think, that these improvements, all dependent on a scientifically based medicinal therapeutics, together with the now rapidly improving methods of establishing and physiologically controlling an artificial circulation, are playing essential parts in the dramatically successful extensions of surgery to the heart and its great vessels, and to the other thoracic organs.

Conclusion

I have been making a rather hasty survey, in which has been noted the recent, fundamental, and still progressive change from a medicinal therapeutics which still derived its sanction from a largely irrational empiricism to one which endeavours to follow ever more closely, step by step, the likewise recent development of an aetiology based upon experimental science. The association will certainly continue, and no limit can be rationally predicted to what may further be achieved under its guidance. I do not think that we can doubt the magnitude of the benefit which it has already brought, and will continue to bring, to medicine in its service to the needs of humanity. In its practical applications, however, it will inevitably raise, and indeed may be assumed to be raising already, new problems for medical practice which are likely to create a peculiar difficulty for unspecialized, general practice.

Pharmacology will certainly acquire a growing importance in any effective scheme of medical education. I imagine, however, that its educational aim will have to be increasingly directed to general principles of pharmacological action, in relation to a scientifically warranted therapeutics. For it is hardly to be expected that pharmacopoeias, textbooks, or even annually revised lectures, will continue to keep pace with *materia medica* subject to continual innovation and replacement, dependent for sources of supply, and, to a large and growing extent, for the researches producing the new remedial agents, on large-scale pharmaceutical industry. To enable the optimum benefit for the patient to be obtained from such new therapeutic resources the practitioner will surely need to have a continuing access to some expert and completely independent source of information and advice, on the relative merits and disadvantages of the remedies presented for his choice in so embarrassing and bewildering an abundance, as well as access to the means for a scientific control of the application of many of them. It is one among the needs which we might surely expect a national health authority to be considering, if it were indeed concerned, not merely to settle the terms on which a medical service of some kind can be provided, but to ensure the continuous and progressive improvement of that service, in step with the relevant advances of scientific knowledge and methods, as these become available for its use.

When the All-India Institute of Mental Health was inaugurated at Bangalore in 1954, the Health Minister, RAJKUMARI AMRIT KAUR, emphasized the inadequate facilities for treating mental patients in India. Training facilities, too, were equally poor. The object of the Institute is to train psychiatrists, nurses, and other staff for mental hospitals and to conduct research into problems of mental health and disease. It also intends to conduct surveys in selected urban and rural areas on the incidence of mental abnormality and the biological factors that accompany mental illness. The Institute is divided up into fifteen departments; these include departments of clinical psychiatry, social psychiatry, neurology, radiology, and statistics. In addition to a number of distinguished Indian medical men on the staff, the Institute has, as medical consultant, Dr. W. Mayer-Gross, of Great Britain. Its handbook, recently published, gives details of the courses of study available to postgraduate students.

SUDDEN LOSS OF MEMORY

BY

ALEXANDER KENNEDY, M.D., F.R.C.P.
F.B.Ps.S.

*Professor of Psychological Medicine, University of
Edinburgh*

AND

JOSEPH NEVILLE, M.B., M.R.C.P., D.P.M.
D.C.H.

*Senior Registrar, Joint Department of Psychological
Medicine, Royal Victoria Infirmary and King's College,
Newcastle upon Tyne*

The individual who is found wandering or who presents himself at a police station unaware of his identity, his past history, or his whereabouts is a familiar figure in the lay mind which has been reinforced by his frequent appearance in fiction and on the screen. Sudden amnesia is well known in a variety of contexts as occurring after accidents involving head injury, after epileptic fits, in boxers who have been knocked out, after alcoholic intoxication, after sudden personal disasters or bereavements, after stage displays of hypnotism, in battle-exhausted soldiers, in criminals facing arrest, in absconding treasurers and reluctant bridegrooms. Further, it can be induced in normal subjects under hypnosis, and has occurred from time to time in almost epidemic form, the report of one case in the press being the signal for the appearance of several apparently sincere imitators.

Although general opinion thus recognizes that the aetiological factors responsible for an apparently identical clinical feature, a period of amnesia, may be either organic or psychogenic, there is scanty reference in the medical literature to this dual approach to the study of pathological forgetting. Owing to the transient nature of the amnesia, a relatively small proportion of those seen at police stations, casualty clearing stations, hospital accident departments, and doctors' surgeries are referred for detailed psychiatric examination or are, in fact, examined before time and circumstance have either led to recovery or greatly modified the clinical picture.

The object of the present study was to examine the aetiological factors in two series of instances (in all, 74 cases) of abrupt failure of memory with or without loss of personal identity, with a view to establishing the relative importance of organic, psychoneurotic, and volitional factors. As the result of this investigation general conclusions have emerged which seem to us to have practical significance for diagnosis and treatment. In this article the attempt is made to extract from the detailed case material the evidence on which these conclusions may be based.

Previous Investigations

Freud (1904) stated that the primary function of amnesia is that of defence against the admission into conscious thought of traumatic elements involved in a neurotic conflict. An entirely dissimilar view was held by Mott (1919), who, basing his conclusions on patients from the first world war, implicated the factor of organic commotio cerebri resulting from shell-shock. Parfitt and Gall (1944), from their experience of Service men who had not been exposed to front-line hazards, described abrupt loss of memory as a hysterical or malingered evasion of the consequences of

wrongdoing. The hysterical symptom was thus in their view the result of an act of conscious evasion, subsequently forgotten.

The majority of cases of sudden loss of memory reported in the literature are regarded as hysterical. Kanzer (1940), describing a series of 71 cases of "hysterical amnesia" admitted in 1937 to the psychiatric division of the Bellevue Hospital, claimed that the patient who took refuge in loss of memory displayed his basic inadequacy by adopting a childish and ineffective method of dissociating his conscious mental life from the problems demanding solution. Parfitt and Gall also stressed the global psychic immaturity and exhibitionistic dependency of most of their patients, and made a distinction between hysterical amnesia and the "defensive" amnesia found in the neurotic personality. These authors' suggestions echo Gillespie's (1937) division of amnesia into simple malingering and "affective" amnesia; the latter term would appear to be synonymous with Parfitt and Gall's "defensive" memory loss. Several workers have observed that somatic manifestations of tension and anxiety are often exhibited by the amnesic patient. In terms of the theory of the unconscious this could be interpreted as an incomplete conversion with anxiety reaching the conscious level.

Wilson, Rupp, and Wilson (1950) described evidence of perturbation and anxiety in a small number of their series, and Adatto (1949) was impressed with the frequent coexistence of anxiety with amnesia. Kanzer's patients, most of whom were basically reacting against financial and interpersonal difficulties, were often worried by headache, and this symptom was interpreted by him simply as indicating the presence of emotional tension. Abrupt loss of memory, then, is not always a fully effective screen against disabling anxiety. Again, sudden loss of memory tends to be of short duration so that the patient may soon be confronted again with his original difficulties. This is in contrast to the persistent and sometimes impregnable neurotic defences employed by the persistent hysteric, who finds a solution of his conflict in some more permanent disability.

A further point of interest is the tendency of functional amnesia, in the opinion of several observers, to occur in association with demonstrable organic disease of the central nervous system (Kennedy, 1940, 1949; Stengel, 1941, 1943). In some of Kanzer's patients the periods of amnesia followed head injuries and epileptic attacks. Wilson *et al.*, in a series of 59 cases of amnesia, described 15 suffering from organic nervous disease. In this group alcoholic intoxication was a common precipitant of amnesia. In a statistically well-controlled study, Berrington, Liddell, and Foulds (1956) demonstrated that the incidence of previous head injury was significantly greater in patients presenting with amnesia or fugue than in other psychiatric patients. Williams (1947) incriminated epilepsy as an important cause of amnesic episodes in air-crews while flying, and indicated that psychoneurotic factors might contribute towards the production of epileptic amnesia. Williams's work bore out the suggestion, made by Lennox (1943), that cerebral dysrhythmia was an important cause of attacks of loss of memory in patients who, in many cases, had never had convulsions. Despite the attractive psychodynamic interpretations that could be elaborated around individual cases, Abeles and Schilder (1935) pointed to the appearance of short-term amnesic attacks in illnesses as different as cerebral arteriosclerosis and carbon monoxide poisoning, and Buzzard (1897) was probably the first to draw attention to its frequency in disseminated sclerosis.

The plea of sudden loss of memory in criminal actions is quite frequently made, though less often sustained, and Baker (1901) was a pioneer in delineating epileptic automatic behaviour as a valid defence. This view was first given the support of electroencephalography in this country by Grey Walter. Hopwood and Snell (1933) indicated the hysterical or malingered nature of amnesic states in criminals, pointing out that recollection waxed as the need for repression waned. Authorities such as Leitch (1948)

and Davidson (1952) have attempted to distinguish clearly between malingered, hysterical, toxic, and organic amnesic states, but other workers have found themselves forced to admit that clear-cut differentiation is rarely possible (Wilson *et al.*, 1950). Siegal (1951) and Rudolf (1947) have produced evidence that similar transient memory loss can occur in both psychoneurotic and organic disorders.

Theoretical Considerations

In daily life an almost continuous inward process takes place by which a selection is made of the information offered to the senses, in such a way that priority in the field of attention is given to those stimuli which are of immediate interest and use to the individual. That this can occur presupposes the rejection from consciousness or the relegation to marginal awareness of material inconsistent with the preoccupations of the subject at the moment. A man who is walking on a girder high above the ground, for instance, will do so more efficiently and safely if he pays close attention to stimuli from his immediate path and rejects distant appeals to his attention. This mechanism has survival value. At a higher level of nervous integration, the attention required to deal mentally with an abstract problem can be achieved only by the rejection of outside stimuli from consciousness for the time being. This is comparable to the way that the music of a bad gramophone record can be appreciated with an effort by concentrating on its meaning and thus discriminating against the accompanying noise. There is abundant evidence of such a discriminating mechanism at lower levels of nervous activity. The principle necessary for such a system—that is, that one part of the nervous system may react positively or negatively as the result of activity in a distant part—is well established (Sherrington, 1947).

It has been suggested that the clinical manifestations of hysteria may be an exaggeration of normal protective mechanisms (Kennedy, 1940). In time of war, for instance, the soldier must learn to blind himself to some of the realities and dangers of his situation in battle if his efficiency is not to be reduced by anxiety. Under overwhelming or long-continued stress this mechanism of rejection of the not-immediately-useful or the unpleasant may be exaggerated to a pathological extent. Apart from the almost experimental situations of war a process of rejecting sensory impressions, percepts, and memories which are painful or which are inconsistent with the individual's preconceived beliefs or attitudes is a commonplace in the genesis of psychiatric symptoms, and, under the name of "repression," is an essential part of psycho-analytic theory.

Mechanisms of a similar kind have been observed clinically and experimentally at different levels of nervous activity and of human behaviour, and according to the point of view have been called "negative induction" (Pavlov, 1928), "extinction" (Bender, 1952), "non-realization" (Kennedy, 1940), "selective attention" (Mackay, 1952), "discrimination" (McCulloch, 1948), or "selective inattention" (Sullivan, 1956). It is certain that without some such mechanism for excluding from attention stimuli that might detract from efficiency in dealing with any matter in hand it would be impossible for thought and action to be directed consistently towards predetermined ends. Since the capacity to direct the attention towards a goal is essential for normal survival and adaptation, if such a mechanism exists, it might be expected that it would exhibit disorder not only when the individual was under psychological stress but when the nervous apparatus was the subject of disease.

From the psychological point of view, where stress is sufficiently in excess of the individual's constitutional powers of adaptation to give rise to a pathological and intolerable degree of tension, inward flight may be achieved (Kennedy, 1940) either by diverting the attention from the tension-producing subject (flight into activity) or by reducing the amount of attention paid to stimuli generally (flight into non-realization). The latter mental mechanism results in the extinction from the individual's consciousness of the

offending subject, together with a greater or lesser amount of psychologically related material. The result is a loss of awareness of the intolerable memory or situation, which either may be strictly circumscribed to the offending "complex" or may spread to involve the whole of his conscious mental activity, resulting in hysterical stupor or coma. An examination of patients with loss of memory might therefore be expected to throw some light on the conditions, nervous and psychological, which favour or prevent this gross pathological exaggeration or caricature of a normal process which must be in operation at every moment of our waking lives.

Method and Results

Two independent investigations were carried out, one by each of us, giving a total of 74 patients studied in detail.

Series 1 (J. N.).—During a period of three years 21 patients presented themselves with amnesia at the casualty and outpatient departments of the Charing Cross and Fulham Hospitals, at the Royal Salop Infirmary, and at the Royal Victoria Infirmary, Newcastle upon Tyne. All were examined while amnesia was present and a full history was obtained after recovery.

Series 2 (A. K.).—During a period of 15 months 53 persons presented themselves at a London police station or were referred to the Maudsley Hospital. All were at the time unaware of their identity and could not, or would not, tell whence they had come. All were examined for the first time while fully amnesic and the majority were either admitted to hospital for investigation or seen on several occasions as out-patients. In 51 it was possible to take a full history from the patient after recovery from the amnesia, while in two patients who proved to have a gross underlying organic loss of memory the history of the illness was obtained from relatives only.

Investigation of Aetiology

In all the cases it was possible either to arrive at a firm diagnosis of organic nervous disease or to come to a definite conclusion regarding the immediate psychological precipitants. Simple situations relating to marriage, bigamy, debt, financial irregularities, in which there was an element of either fear or conflict, accounted for most of the amnesias. There was a group of patients with a history of previous hysterical illness and of practice in the use of the advantage-of-illness motive. Similar factors were brought to light in some of the patients with demonstrable organic disease, and the amnesia was thus the means of bringing to light the fact that the patient's general adaptation to life had been deteriorating for some time. The following are illustrative of special types:

1. A man of 32 who had walked blindly for several miles before jumping into a pond and who had been rescued by a passer-by. He was calm and helpful but could give no account of himself. Under hypnosis his condition became one of severe agitated depression. There was a history of two previous attacks of depression. His amnesic state was evidently an escape from his psychotic depression, and the idea of suicide had evidently influenced his behaviour in the fugue.

2. An Australian company director who had persuaded others to put up money for a large venture. On doubts being expressed at a meeting about the soundness of his ideas he became abusive and walked out into the street. Picked up as an unknown man, he had been found to be suffering from general paralysis, and for this he was treated in a mental hospital. After improving considerably he escaped, and, in spite of complete amnesia for the period of his illness, obtained work and prospered. His organic illness came to light when he sought advice in this country about the part of his life of which he had no memory. His history, given after the amnesia had responded to hypnosis, was fully confirmed.

3. A young man who was picked up in an exhausted condition by the police. On recovery from the amnesia it transpired that the patient feared that he had murdered his sister. He had in fact attacked her, but she was not seriously harmed. His previous history and the E.E.G. allowed of a confident diagnosis of psychomotor epilepsy. The epileptic automatism had been

followed by a functional one due to the situation created by his attack of violent behaviour, of which there had been previous examples.

4. A man of 28 with a good work record up to four years previously had been sent to prison for a series of burglaries. He was wanted by the police and had been sleeping in houses which he entered at night in search for food and money. He walked into a police station in an amnesic state. After some discussion his memory returned and he said that he would like to tell the police what he had been doing, serve his sentence, and get a job. On examination, however, there was an early hemiparkinsonism. There was a history suggestive of an attack of encephalitis seven years before. Four years ago he had left a good job and taken to night work because of inversion of his sleep rhythm. The decline in his adaptation to life could clearly be traced to the organic disease.

5. A 17-year-old girl of low average intelligence was brought to hospital by the police, having been found "sleeping rough." At first she exhibited a complete loss of memory extending to ignorance of her identity, but apart from a patchy amnesia for the previous two days her memory returned after a bath and a meal. She had run away to London after a quarrel with her stepmother. To make a living without revealing her whereabouts to her family, and since she "felt that the end of the world had come and things couldn't be worse," she began to accost men in one of the royal parks. Promiscuity seemed initially to gratify her, but after she had been ill-treated and robbed by a gang of men she became severely depressed. During the two days' amnesic period she had evidently been in a state of clouded consciousness, making no effort to obtain food or shelter. In this case a combination of amnesia with a fugue state seems to have acted as a screen against realization of what she had done and against her intolerable depressive thoughts. In this case no useful purpose would have been served by attempting to break down this defence, and, after her memory had recovered in other respects, the matter was not pressed.

The aetiology of the amnesia as ascertained in the 74 cases is summarized in Table I.

TABLE I.—*Psychiatric Aetiology in 74 Patients with Sudden Amnesia*

	Series		Total	%
	1	2		
Gross organic disease of C.N.S. . .	8	19	27	37
Major epilepsy	5	4	9	12
Hysterical amnesia without organic disease	10	25	35	47
(a) With previous history of hysteria	6	16	22	30
(b) No previous neurosis	4	9	13	17
Psychopathic personality or very bad employment record	8	7	15	20
Feeble-minded or illiterate	0	2	2	3
Admitted malingering	0	5	5	7
Fugue associated with suicidal attempt	3	3	6	8
Functional psychosis present	3	4	7	10
Depressive state	2	3	5	7
Other	1	1	2	3
Fearing arrest or disciplinary action	3	9	12	16
Senile	0	2	2	3
Under age 12	3	0	3	4

Organic disease was found in 27 patients. The distribution is shown in Table II.

TABLE II.—*Varieties of Organic Nervous Disease Found in 27 Out of 74 Patients with Sudden Amnesia*

	Series		Total	%
	1	2		
Idiopathic epilepsy	1	4	5	19
Head injury	3	1	4	15
Chronic encephalitis lethargica	1	3	4	15
Rheumatic chorea	1	0	1	4
Disseminated sclerosis	0	2	2	7
Increased intracranial pressure	0	2	2	7
Cerebral syphilis	0	2	2	7
Presenile dementia	1	3	4	15
Hypertensive encephalopathy	1	0	1	4
Freidreich's ataxia	0	1	1	4
Diffuse sclerosis	0	1	1	4
Total	8	19	27	

In some of the patients it was not possible to assign a single aetiology, and organic nervous disease existed in the presence of a clear psychogenic precipitating factor. The overlap of aetiologies is shown in Table III.

TABLE III.—*Psychogenic, Organic, Psychotic, and Malingered Amnesia in 74 Cases Showing Overlapping Causes*

	Series		Total	%
	1	2		
Convincing psychogenesis discovered but no evidence of organic disease	10	16	26	24
Psychogenesis elicited and organic disease present	6	19	25	23
Organic disease or dementia present	8	21	29	27
Organic disease present but no evidence of psychogenic factors	2	8	10	9
Malingering	0	5	5	5
Psychosis, excluding depression	3	4	7	7
Endogenous depression with hysterical fugue	2	2	4	4
Total	31	75		

The extent to which organic and psychogenic factors contributed throughout the series is shown in the following analysis of 61 patients with sudden amnesia (malingerers and psychotics excluded): psychogenic, 43%; dual aetiology, 41%; organic, 16%.

The sex incidence is shown in Table IV.

TABLE IV

	Series		Total	%
	1	2		
Males	16	36	52	70
Females	5	17	22	30
Total	21	53	74	100

Treatment

Amnesia is a condition calling for symptomatic treatment if only to enable a history to be taken from the patient. There is a strong tendency to spontaneous recovery, although this may be much delayed if nothing is done. The method chosen in the two series depended on three factors: (1) The kind of treatment possible in a police station, casualty department, or general hospital ward, where absence of distracting noise is difficult to ensure. (2) The state of the amnesia and the attitude of the patient to medical aid. The presence of mental confusion, intoxication, or aggressiveness, for instance, made hypnosis or narcotic abreaction impossible. (3) The approach used was that which the individual investigator preferred and felt he could use with success. Thus in Series 1 (J. N.) suggestion with a vehicle was used in 4 out of 21 cases, while it was not used at all in Series 2 (A. K.), where 31 out of 53 patients were treated by means of hypnosis only. It was shown that recovery could be brought about by six main methods.

(i) *Therapeutic Anamnesis*.—Establishing contact with the patient and causing memory to return by encouraging him to extend his history until he realized that he could recall details about himself normally.

(ii) *Hypnosis*.—Amnesic patients are easy to hypnotize if the induction is rapid and amnesic somnambulism rather than light hypnosis is the aim. Sometimes induction is almost instantaneous. Under hypnosis the patient is made to relive known incidents in his past and to produce a sort of running commentary. During this there is usually considerable emotional abreaction with deepening of the hypnosis. When the emotion has subsided the patient is kept talking and told to wake up, when he discovers that he is recounting past events and that his amnesia is at an end.

(iii) *Narcotic Abreaction*.—A great deal of the effect of using intravenous barbiturates is due to suggestion, especially if the patient has the impression that he has been given a "truth drug." There is usually a considerable abreaction, and the patient may relive some event related to his amnesia and realize its significance when the effects of the narcotic have passed. This aspect is increased by the use of methamphetamine.

(iv) *Suggestion, Direct or with Vehicle*.—An impressive approach to the patient may induce recovery almost at once, and the giving of an injection, exposure to flashes of light from a photic stimulator, or the use of faradism to the scalp may act as vehicles for suggestion.

(v) *Direct Discussion*.—This was reserved for shallow hysterics and also for malingerers wanted by the police and "on the run." The process consisted in letting the patient know that loss of memory was a method of evasion fully understood by both doctors and police officers, and that the best course would be to avoid introducing medical and psychiatric complications into a simple matter of law. Where the patient had already been picked up by the police he was then questioned by them *de novo* after a rest, there being no admixture of police and psychiatric procedure.

(vi) *Medical Measures*.—Where the amnesia is associated with a physical disorder affecting the nervous system it is necessary to apply as early as possible any method likely to improve the level of consciousness which carries no undue risk to the patient. The single case of hypertensive encephalopathy responded to the intravenous injection of 50 ml. of hypertonic dextrose alone, while one patient with raised intracranial pressure responded to intravenous sucrose. In the other it was deemed inadvisable to use this method, and the patient recovered his memory under light hypnosis attempted on the principle that if the amnesia can have a dual aetiology it might respond to either physical or psychological treatment.

The mode of recovery of the 74 cases is summarized in Table V.

TABLE V.—*Progress and Response to Immediate Treatment of Amnesia*

	Series		Total
	1	2	
Spontaneous recovery, immediate or after medical measures	0	3	3
Recovered during therapeutic interview	14	12	26
" under hypnosis	0	31	31
Suggestion with vehicle	4	0	4
Recovered after narcotic abreaction	1	2	3
" spontaneous convulsion	2	0	2
Malingering, dismissed with warning	0	3	3
Organic amnesia, no appreciable improvement	0	2	2
Total	21	53	74

Clinical Findings

The majority of the patients were aware of their loss of memory and either complained of this or allowed it to be inferred. In many, personal identity was retained with ignorance of address and locale. The greater number exhibited an attitude of incurious detachment to their disability, although others were distressed or perplexed. It was not possible to separate features of the amnesia itself which indicated an organic or functional cause, and even when the patient's capacity for attention was low the state of consciousness was of small diagnostic value. The discovery of organic nervous disease depended, therefore, on physical signs or on the detailed neuropsychiatric assessment made after memory was restored. In two patients with gross degenerative cerebral disease there was very severe underlying organic loss of memory, yet at the first impression the clinical picture was not unlike that seen in the exhausted patient with a functional amnesia.

A finding common to both series was that the patients were highly suggestible, so that the level and content of the amnesia was readily influenced by the nature of the investigator's questions and by his attitude as interpreted by the patient. This suggestibility in recent hysteria is in contrast to the resistance to suggestion characteristic of patients with functional disabilities which have become well established. As experience grew it became easier to relieve the loss of memory in organic as well as in functional amnesia by direct suggestion. In the second series it was noteworthy that a very high proportion of the patients were

easy to hypnotize, irrespective of whether or not organic disease was present. It was easier to induce a deep somnambulistic state than light hypnosis, as if the patients welcomed the further escape from present problems that it permitted, and rapid induction seemed to work well. When the patient was induced to relive the period before the amnesia, hypnosis usually became deeper and then lighter as violent emotional storms made their appearance, and it was in this way that memory and orientation were usually regained.

The malingerers, who were all confirmed by admission on their part, were hardly suggestible at all on interview, but, surprisingly, two of them were easily hypnotized. It may be that they were quite glad to take the opportunity of having their problems brought to light, and one insisted on going to the police and reporting a number of burglaries. It may be significant that none of this group had committed anything more than petty offences, and it is hardly likely that a malingerer who was concealing a serious criminal offence would have co-operated so well. It is not suggested that admission of malingering excludes a diagnosis of hysteria.

Among the patients with organic disease epilepsy was the condition most often found. The fact that gross hysterical states, including amnesia, appear to be particularly common in association with epilepsy, disseminated sclerosis, chronic encephalitis lethargica, and after head injury would suggest that the destruction of tissue in these disorders in some way specially predisposes the patients to hysterical "conversion" or lowers the threshold at which stress will bring it about, and the present series would bear out this impression.

Subsequent Treatment

In every case a patient presenting with sudden loss of memory was discovered, after this symptom was relieved, to have serious organic nervous disease, to be malingering with the object of covering an offence, or to have an intolerable depressive or anxiety state from which temporary escape was being secured by a process of inward flight which exceeded the mere repression of a painful affect and extended the repressed material to include a loss of memory, identity, or orientation, or a combination of these. Where the aetiology was predominantly psychogenic the underlying problem, when brought to light, was evident enough and consisted mainly of banal difficulties such as conflicts over marriage, hire-purchase overcommitment, quarrels with parents, extramarital pregnancy, or misuse of funds. In the patients with organic disease it was evident that some of them had been able to deal with their daily problems quite adequately until handicapped by a disability of which they themselves were unaware. When finally breakdown had occurred it had done so after almost trivial stress. This was evident also in the depressives, who were often quite unaware of their underlying mood change. Some simple folk, with little comprehension of abstract ideas, when suffering from depression are not able to understand what has happened to them and can express their feelings only in terms of somatic symptoms. Of these the amnesics are a special variety, and, since the prognosis of the underlying depression is usually good, its early recognition is most important.

A general impression of the whole group was that in presenting themselves with an amnesia for their own difficulties these patients were flying a distress signal and throwing themselves on the mercy of the community or of the hospital to which they came, and that they hoped in an unclear and unconscious way that someone could apply to their problems a patience and imagination which they themselves did not possess. In many it was difficult to obtain their co-operation in reviewing and making decisions about their difficulties. In this primitive reaction to stress they expected magic, and, having got it in the form of a sudden relief of amnesia, they preferred to remain dependent on the "magician" rather than to resume, even with help, responsibility for their own lives.

Conclusions

Investigation of two series of cases of sudden amnesia indicates that organic and psychological factors may both take part in the production of a clinical state which is fairly uniform in its manifestations.

The high proportion of cases of organic nervous disease in both series and in other series in the literature suggests that its presence may predispose to primitive mental mechanisms of escape.

Amnesic episodes are frequent incidents in the course of the degenerative hysteria seen in psychopathically inadequate individuals, or in those suffering from chronic organic brain disease. When the amnesia is relieved in these patients, relatively little anxiety is present and there is conspicuous pathological detachment—the *belle indifférence* which Janet (1907) described in just such patients.

It would seem that episodes of amnesia are particularly apt to occur in cases of organic brain disease in which disturbances of consciousness, of attention, and of memory are present. Even in the presence of gross organic disease a complete and convincing psychogenic causation can often be found. Neither aetiology excludes the other.

Examination of the precipitating factors suggests that the immediate purpose of the amnesia is to protect the individual from mental pain or tension which is intolerable to him.

Flight may be from any kind of unpleasant emotional state or "mental pain," and of these endogenous depression is one. The fugue in which a suicidal attempt or gesture occurs may well be due to a depressive state masked by the amnesia.

Amnesia of functional type rarely persists longer than a few hours or days. It is a transitional state, the immediate psychological purpose of which is flight. It passes on either to recovery or to the development of some form of pathological adjustment, such as a hysterical disability.

Where there is a good previous personality, treatment of the amnesia often reveals a severe anxiety and a pressing personal problem with which the patient needs urgent help. In this case the prognosis is usually very good.

Malingering amnesia is not usually difficult to detect, but hysterical amnesia frequently resembles malingering. The clinical picture is determined to some extent by the patient's idea of an amnesic episode, as popularly understood. The patient, if pressed, may actually admit he is malingering when in fact there is strong evidence of an organic or psychogenic causation. A confession cannot therefore be accepted as certain evidence of malingering.

Epilepsy, whether latent or manifest as convulsions, is a common background to amnesia. Chronic encephalitis, disseminated sclerosis, and the effects of brain trauma appear especially to predispose to amnesia. After treatment of hysterical amnesia search should always be made for a residual organic memory defect and other organic deficits.

The method of treatment is not important, as recovery occurs usually within a few days. For patients arriving at a casualty department, hypnosis, suggestion during injection of an intravenous barbiturate, and simple therapeutic anamnesis appear to do equally well. Since the patient is highly suggestible the danger of producing iatrogenic symptoms or a false psychogenesis must always be borne in mind.

The conflicts and stresses found in the patients with previously good personalities were mainly traceable to simple domestic, social, or financial difficulties. It would appear from war experience that if the tension is sufficient, anyone can develop an amnesia, but organically vulnerable and psychopathic individuals often become amnesic on trivial stress. In degenerative hysteria with character disorder the amnesia may be imitative or designed simply to draw attention to the patient or his maladjustment to life. Organic brain disease may first come to medical notice in this way.

The study of sudden amnesia indicates that it is not possible to draw any sharp dividing-line between organic and functional disabilities and that it is not always possible to distinguish clinically the causes of pathological forgetting.

Summary

The aetiology of sudden loss of memory has been investigated in two series of cases, totalling 74 patients who were submitted to a full neurological and psychiatric examination.

In both series a large proportion suffered from undetected gross organic nervous disease. In the majority, whether organic disease was present or not, a simple psychogenic cause was found and the amnesia could be demonstrated to be a psychological escape mechanism. There is evidence that organic nervous disease may predispose to this type of psychological mechanism. Sudden amnesia is a temporary state that responds well to simple psychotherapy, which can be given, if necessary, in a casualty department. It is essential in such cases to seek the causes of the patient's underlying maladjustment not only in his life and circumstances but in his central nervous system as well.

REFERENCES

- Abeles, M., and Schilder, P. (1935). *Arch. Neurol. Psychiat. (Chicago)*, **34**, 587.
 Adatto, C. P. (1949). *Ibid.*, **62**, 82.
 Baker, J. (1931). *J. ment. Sci.*, **47**, 260.
 Bender, M. B. (1952). *Disorders in Perception*. Thomas, Springfield, Illinois.
 Berrington, W. P., Liddell, D. W., and Foulds, G. A. (1956). *J. ment. Sci.*, **102**, 280.
 Buzzard, T. (1897). *Lancet*, **1**, 1.
 Davidson, H. A. (1952). *Forensic Psychiatry*, p. 160. New York.
 Freud, S. (1904). In *Collected Papers*, **1**, 267. London, 1924.
 Gillespie, R. D. (1937). *Arch. Neurol. Psychiat. (Chicago)*, **37**, 748.
 Herman, M. (1938). *Psychiat. Quart.*, **12**, 738.
 Hopwood, J. S., and Snell, H. K. (1933). *J. ment. Sci.*, **79**, 27.
 Janet, P. (1907). *The Major Symptoms of Hysteria*. London.
 Kanzer, M. (1940). *Amer. J. Psychiat.*, **96**, 711.
 Kennedy, A. (1940). *J. ment. Sci.*, **86**, 988.
 — (1949). *Modern Practice in Psychological Medicine*, edited by J. R. Rees, chapter 19. Butterworth.
 Leitch, A. (1948). *Med. Press*, **219**, 459.
 Lennox, W. G. (1943). *Amer. J. Psychiat.*, **99**, 732.
 McCulloch, W. S. (1948). *Ann. N.Y. Acad. Sci.*, **50**, 259.
 Mackay, D. M. (1952). *Proc. Arist. Soc., Suppl.*, **74**.
 Mott, F. W. (1919). *War Neuroses and Shell Shock*. London.
 Parfitt, D. N., and Gail, C. M. C. (1944). *J. ment. Sci.*, **90**, 511.
 Pavlov, I. P. (1928). *Lectures on Conditioned Reflexes*. Internat. Publisher, N.Y., 1941.
 Rudolf, G. de M. (1947). *J. ment. Sci.*, **93**, 342.
 Sherrington, C. (1947). *Integrative Action of the Nervous System*. Cambridge Univ. Press, Cambridge.
 Siegal, L. J. (1951). *A.M.A. Arch. Neurol. Psychiat.*, **66**, 700.
 Stengel, E. (1941). *J. ment. Sci.*, **87**, 572.
 — (1943). *Ibid.*, **89**, 224.
 Sullivan, H. S. (1956). *The Interpersonal Theory of Psychiatry*. Tavistock Publications, London.
 Williams, D. (1947). In *Psychological Disorders in Flying Personnel of R.A.F. 1939-45*. H.M.S.O., London.
 Wilson, G., Rupp, C., and Wilson, W. W. (1950). *Amer. J. Psychiat.*, **106**, 481.

An urgent appeal to governments, scientists, and industry to intensify research into the global problem of insect resistance has been made by the Technical Conference on Insect Resistance called together by the World Health Organization in Geneva recently. According to a statement issued by W.H.O., in 1946 only two kinds of insects of public health importance, the house-fly and certain kinds of mosquitoes, were resistant to D.D.T. To-day they number 38, among them several species of malaria-transmitting mosquitoes, plague-carrying fleas, and typhus-bearing lice. Countries all over the world have spent huge sums of money to control the great insect-borne pestilences. Faced with the possibility of failure, they have no practical solution for the control of resistant insects except emergency switching from one insecticide to another. But scientists are running out of effective insecticides which are at the same time economic and non-toxic to man, although chemical firms are testing as many as 50,000 compounds a year for their insecticidal properties. The world may soon face an emergency because of the dynamic nature of the resistance problem as opposed to the more static nature of many other scientific and technical problems.

POLYPOSIS OF GASTRO-INTESTINAL TRACT: THE PEUTZ SYNDROME

BY

DAVID BAILEY, M.Chir., F.R.C.S.

First Assistant, Surgical Unit, University College Hospital, London

The association of multiple adenomata of the gastro-intestinal tract with melanin pigmentation of the oral mucosa, lips, and face was first described by Peutz (1921), but his description attracted little attention at the time. The subject was fully reviewed by Jeghers *et al.* (1949), and since then many cases have been reported. The syndrome is, however, still not widely known, and the significance of the characteristic melanin pigmentation may not be recognized even if it is noted on clinical examination, and experience has shown that it may be missed altogether if not especially sought.

In this paper four cases of gastro-intestinal polyposis are described, in two of which oral and facial pigmentation was certainly present, but it was not noted in the other two. My purpose is, firstly, to draw attention again to this important association, which may prove a useful diagnostic aid, and, secondly, to show that there is without doubt a considerable risk of malignant change developing in these polyps and to discuss the problems of treatment posed by this tendency.

Case 1

A woman aged 35 who had suffered from recurrent attacks of abdominal pain for 17 months was admitted to University College Hospital in June, 1955. The pain was ill localized, but appeared to begin in the epigastrium. The attacks lasted one to four hours and occurred at intervals varying from a few days to five or six weeks. The pain, which was intermittent, with intervals of complete freedom between the spasms, had been more severe recently. During the attacks, which were not related to food, she felt sick but did not vomit. Usually diarrhoea followed subsidence of the pain, and on several occasions she had passed a little fresh blood in the stool. She also suffered from mitral stenosis and atrial fibrillation, and the first attack of pain occurred about six weeks after a successful mitral valvulotomy had been carried out at another hospital.

She had no children and no brothers or sisters. She could not recall that her parents (now dead) had ever suffered from any serious abdominal symptoms.

On examination she was seen to be a thin, healthy-looking woman. Atrial fibrillation at 72 a minute was controlled with digoxin. A tender, mobile, soft sausage-shaped mass was felt in the left iliac fossa. Rectal examination was negative. Sigmoidoscopy showed no abnormality. There was melanin pigmentation of the face, lips, hands, and feet, consisting of numerous dark-brown and black discrete macules, which were particularly numerous on the lower lip, around the medial side of the eyes, on the back of the hands and fingers, and on the dorsum of the feet. Scattered over the face were several other lighter-coloured spots resembling freckles. Mucosal pigmentation was present on the buccal surface of the lips and cheeks, consisting of several dark blue-black patches varying from 1 to 5 mm. in diameter. (N.B.: this pigmentation was not recognized at the early examinations, being noted only later when it was specially looked for.)

X-ray Examinations.—Barium-meal and follow-through examination carried out elsewhere shortly after the first attack of pain showed the signs of an entero-enteric intussusception of the upper jejunum. Three subsequent barium follow-through examinations were carried out: the first two