have a shorter duration of stay if they are admitted to a mental hospital.

To demonstrate the relationship between physique and examination results larger samples are needed, because correlations between physique and intellectual performance are usually of a lower order than those between physique and the temperamental and other qualities involved in severe mental illness. In the article¹ to which Dr. Finlay refers only 19 Leeds students are reported who gained first-class honours, together with 14 who were "bad failures" academically. Even if real differences should later be established, there are again differences in manner of selection at the two universities to be considered. Whether or not the small band of brilliant Lf students is a phenomenon peculiar to Oxford it is difficult to say, but the following conclusion from the original Oxford report² should not be forgotten: "Mesomorphs were most consistent in attaining a good level of performance with first- or second-class honours."—I am, etc.,

Oxford.

R. W. PARNELL.

REFERENCES

- ¹ Finlay, S. E., *Univ. Leeds med. J.*, 1956, 5, 76.
- ² Parnell, R. W., *British Medical Journal*, 1954, 2, 491.

Toxoplasmosis Simulating Glandular Fever

SIR,—Dr. Mary Bateman (*Journal*, July 20, p. 141) has brought a useful reminder to your readers that postnatally acquired toxoplasmosis may give rise to a clinical picture resembling infectious mononucleosis. Dr. J. C. Siin¹ was probably the first to make this observation, and therefore we owe to him the knowledge that toxoplasmosis is among the possible causes of a "glandular fever syndrome"; Dr. I. A. B. Cathie² subsequently confirmed this. Siin described seven patients presenting with lymphadenopathy, lassitude, fever, and atypical lymphocytes in the peripheral blood. In each case the Paul-Bunnell test was negative; serological investigation demonstrated that they had acquired toxoplasmosis.

May I take this opportunity of appealing to medical writers to break away from the present fashion of saying "simulate" when "resemble" is meant? The former word, as with "feign," implies active pretending.—I am, etc.,

London, W.1.

REGINALD LIGHTWOOD.

REFERENCES

- ¹ Siin, J. C., *J. Amer. med. Ass.*, 1951, 147, 1641.
- ² Cathie, I. A. B., *Lancet*, 1954, 1, 813.

Habitual Abortion Due to Insufficiency of the Internal Cervical Os

SIR,—Abortion and premature labour may start in three different ways: (1) by bleeding associated with placental separation, (2) by painful uterine contractions, or (3) by rupture of the foetal membranes. In those cases in which the first evidence of impending abortion is premature rupture of the membranes, it is tempting to assume, as do Mr. V. B. Green-Armytage and Professor J. C. McClure Browne (*Journal*, July 20, p. 128)—and many before them—that an incompetent cervix is the cause. There is no reliable evidence to support this view, and, indeed, a controlled study of cases of abortion and premature labour reported in 1956¹ showed that patients with radiologically demonstrated cervical deficiency are not more likely to commence labour in this way than in any other. Moreover, those women who have a normal upper cervix according to hystero-graphy are just as likely to begin the evacuation of their uteri by rupturing the membranes as are those with a "funnel"-shaped cervix.

Out of a total of 306 cases of premature labour and abortion studied, 104 (34%) gave a history of rupturing the membranes as the first sign that the uterus was about to evacuate itself. In cases where a probable cause for the mishap could be determined clinically (for example, foetal abnormality, twins, hydramnios, hypertension, toxæmia, etc.) premature rupture of the membranes occurred in 42 out of 174 (24%).

Of the remaining 132 cases in which the cause of abortion or premature labour was not obvious, 81 were later subjected to hystero-graphy, and the findings were as follows:

Hystero-graphic Appearances	No. of Cases	No. Commencing Abortion or Labour by Rupturing the Membranes
Cervical incompetence	38	12 (32%)
Cornual malformation	19	10 (53%)
Cervical incompetence and cornual malformation	12	5 (42%)
Normal uterus	12	5 (42%)
Total	81	32 (40%)

Whatever be the cause of premature rupture of the membranes, it is one which operates indiscriminately throughout the various groups of cases of recurrent abortion. Furthermore, if a sufficiently large number of cases are investigated it will be found that many women will carry a subsequent pregnancy to term despite a history of previous abortions, despite a radiological demonstration of an incompetent cervix, and despite not receiving any treatment—medical or surgical.

It has to be recognized too that a cervix which is anatomically normal between pregnancies may be functionally incompetent during pregnancy. Its efficiency as a functional sphincter may even vary from pregnancy to pregnancy in the same woman. Even so, the finding of a dilated cervix and a bulging bag of membranes during pregnancy is not necessarily significant and is not incompatible with the continuance of pregnancy to term.

In making these comments it is not intended to decry the value of the Shirodkar type of operation in certain cases of recurrent abortion but to emphasize the difficulty both in deciding which cases are suitable for it and in assessing the results. If the diagnostic criteria suggested by Mr. Green-Armytage and Professor McClure Browne are accepted, then many unnecessary purse-string operations will be carried out and many will receive undeserved credit for the birth of a live baby.

Our study led us to conclude that, although an incompetent cervix is a significant cause of abortion and of premature labour, its reliable diagnosis as the operative factor in any individual case is almost impossible. The nearest approach to the diagnosis is a well-authenticated history of several abortions occurring *after* the 14th week, the foetus being alive at birth and the process of expulsion being characterized by *painless* dilatation of the cervix, associated with a hystero-graphic evidence of a funnel-shaped cervix between pregnancies. A history of the membranes rupturing before the onset of expulsive uterine contractions is without significance in this context. Even if these criteria are accepted, the Shirodkar operation is often empirical and its results indeterminate.

It may be added that our findings support the statement made by Mr. Green-Armytage and Professor McClure Browne to the effect that an incompetent cervix, although it may be caused by previous obstetrical or surgical injury, is often found in women who have never had any interference with the internal os. In such it presumably represents either a developmental anatomical error or a disturbance of function.—I am, etc.,

Liverpool.

J. K. WILSON.

REFERENCE

- ¹ Jeffcoate, T. N. A., and Wilson, J. K., *N.Y. St. J. Med.*, 1956, 56, 680.

Increasing Muscle Strength

SIR,—Your leading article on increasing muscle strength (*Journal*, July 20, p. 150) has not given a correct impression of the discussion at the Royal Society of Medicine, and one cannot agree with your general appraisal of the problems involved. It is true that weight-lifters have made advances in their techniques for developing crude strength and that these techniques have been used in clinical practice. But the applications are few, and it is wrong to assume that the