Chart of a typical case.

No similar detailed survey was done in the women's medical ward, but the general impression was that there were no significant differences.

The treatment to which the majority of patients responded was 2 tablets of A.P.C. administered three times a day, a sedative cough mixture given three times a day, light diet, and plenty of fluids. The complications—six cases of bronchitis and one of pneumonia—were treated with antibiotics in the accepted manner. A post-influenzal depression was a fairly constant feature for which no treatment was given.

In a number of these cases there was an associated mild heat exhaustion. However, probably they were mostly cases which had been on a minimum salt intake before the onset of the influenza, and subsequently the fever and excessive sweating were sufficient to create a mild salt deficiency. It is notable that the number of patients admitted with a diagnosis of heat exhaustion was far lower during the influenza epidemic than in previous years, although the climate was not dissimilar to that of other years. Constipation proved to be a troublesome complaint in many cases, again presumably due to fluid loss, a state of affairs not uncommon in cases of mild heat exhaustion.

The Singapore report on Asian influenza noted a high incidence of sore throat. This was not found in the present series, and in fact there was little if anything to distinguish this epidemic of Asian influenza from the influenza seen in other years. Mutation of the causative virus is not an improbable cause, and further reports as the epidemic spreads westwards may possibly confirm this.

D. W. A. MCCREADIE, L.R.C.P.&S.Ed., L.R.F.P.S.Glas.,
Senior Medical Officer, Health Department, Kuwait.

Fatal Pulmonary Embolism in Pregnancy

The following case is worth reporting because of the unusual site of primary thrombosis unassociated with trauma or infection.

CASE REPORT

A booked maternity patient, a para 1 aged 29, was admitted to the antenatal ward complaining of breathlessness on exertion for the past three days. By her dates she was 29 weeks pregnant, and her antenatal course had up till then been completely uneventful. There was nothing of note in her family, past medical, or menstrual history. In 1947 she had a spontaneous term delivery at her home of a male child weighing 7 lb. 8 oz. (3.4 kg.). The antenatal period and puerperium were uneventful.

She booked at the antenatal clinic when she was 16 weeks pregnant. No abnormality of the head, neck, chest, or abdomen was found, her blood pressure was 125/70, and there was no sugar or albumin in the urine. A routine chest x-ray examination at the time showed nothing of note, and her haemoglobin was 12.2 g. per 100 ml., group O rhesus positive. She was referred to the county council clinic until the thirty-sixth week of pregnancy, where she attended at regular intervals.

Three days before admission she became slightly breathless when walking up a flight of stairs while at work, but on

resting for a few moments recovered completely. She was sent home to rest, and was perfectly well until the day before admission, when her breathlessness became more noticeable, the slightest exertion causing discomfort. She was quite comfortable at rest, did not complain of chest pain, and had no other symptoms.

Examination on admission showed a pale young woman lying in bed, in no apparent distress. The temperature and pulse were normal and the respiration rate was 20. No abnormalities were noted in the head, neck, chest, cardiovascular system, or abdomen. The blood pressure was 130/80. The uterus was enlarged to correspond with a 30 weeks' pregnancy, a single foetus was present, and the foetal heart was heard. There was no oedema, nor were there varicosities of the lower extremities. Vaginal and speculum examination revealed no abnormality. The urine contained no sugar or albumin.

A full blood count and a chest x-ray examination were ordered. A physician's opinion was obtained, and although he asked for an electrocardiogram he felt that the cause of the breathlessness was more likely to be ascertained by the blood examination. The haemoglobin was found to be 9.1 g. per 100 ml. and white cells 16,000 per c.mm.; no abnormalities were noted in a blood film. The chest x-ray examination showed nothing of note.

On the following day, while the electrocardiogram was being taken, the patient became cyanosed. Oxygen was given, but her pulse became extremely fast and of poor volume, and she collapsed and died soon afterwards.

As the foetal heart was still audible, a post-mortem caesarean section was performed and a living male infant was delivered; it died, however, an hour after delivery.

A post-mortem examination was performed by the pathologist appointed by the coroner, and following is a relevant extract from the report. *Lungs*: There was minimal congestion at the bases of both lungs posteriorly. Numerous small ante-mortem clots were present in the pulmonary arteries, but there was no evidence of infarction. *Heart and pericardium*. The heart was of normal shape, size, and development. The pulmonary artery was blocked by a large ante-mortem clot. *Pelvic organs*: The veins on either side of the vagina were thrombosed and clots could be traced into the left internal iliac veins. There was no evidence of trauma or infection. The cause of death was given as pulmonary embolus secondary to thrombosis of the perivaginal plexus of veins.

COMMENT

Damage to the intima of blood vessels by trauma or infection, slowing of the blood stream, and a natural or acquired tendency due to changes in the plasma and cellular elements of the blood are all factors which promote clotting of blood in the vascular system. In obstetric practice a large proportion of cases occur as a complication of cardiovascular and pulmonary disease, and thrombosis of varicose veins of the extremities accounts for a number of cases. Ophuls and Dobson (1932) found that 52% of their cases of pulmonary embolus were associated with cardiac disease. In cases of induced abortion, trauma or sepsis may also give rise to pulmonary emboli.

This case is instructive in that none of the specific conditions mentioned above was operative, and the cause for the thrombosis at the primary site remains unexplained. Although pulmonary embolus is one of the first conditions to be considered in the differential diagnosis of the sudden onset of breathlessness, absence of symptoms or physical signs indicating a primary site made a positive diagnosis difficult.

I thank Professor J. C. McC. Browne, of the Institute of Obstetrics and Gynaecology at Hammersmith Hospital, under whose care the patient was admitted, for permission to publish this case.

M. B. WINGATE, M.B., F.R.C.S.Ed.,
Late House-surgeon to the Obstetric Department,
Hammersmith Hospital, London, W.12.

REFERENCE

Ophuls, W., and Dobson, L. (1932). *Arch. Path. (Chicago)*, 14, 215.