REFERENCES

- Allen, C. L. (1899). *J. nerv. ment. Dis.*, 26, 471.
 Bucy, P. C. (1935). *Arch. Neurol. Psychiat.*, 33, 30.
 — (1949). *The Precentral Motor Cortex*, p. 29. Urbana, Illinois.
 Crockett, D. J. (1956). *Lancet*, 2, 1179.
 Drinker, C. K., and Yoffey, J. M. (1941). *Lymphatics, Lymph, and Lymphoid Tissue*. Cambridge, Mass.
 Eccles, M. V. (1956). *Brit. J. phys. Med.*, 19, 5.
 Gilbert, A., and Garnier, M. (1897). *C.R. Soc. Biol. (Paris)*, 4, 553.
 Hare, H. A. (1898). *J. nerv. ment. Dis.*, 25, 189.
 Landis, E. M. (1934). *Physiol. Rev.*, 14, 404.
 Luhan, J. A. (1936). *Arch. Neurol. Psychiat.*, 36, 42.
 McMaster, P. D. (1937). *J. exp. Med.*, 65, 347.
 Parhon, C., and Goldstein, M. (1899). *Roumanie med.*, No. 3. Abstract in *Rev. neurol.*, 1902, 10, 865.
 Raymond, F., and Courtellemont, V. (1904). *Rev. neurol.*, 12, 397.
 Stead, E. A., and Warren, J. V. (1944). *J. clin. Invest.*, 23, 283.
 Taylor, G. W., Kinmonth, J. B., Rollinson, E., Rotblat, J., and Francis, G. E. (1957). *Brit. med. J.*, 1, 133.
 Weiss, S., and Ellis, L. B. (1931). *J. clin. Invest.*, 9, 17.

TRICUSPID ATRESIA

BY

A. L. GWEЕ, L.M.S., M.R.C.P.

Senior Registrar, Department of Clinical Medicine,
University of Malaya, Singapore

AND

E. S. MONTEIRO, C.B.E., M.D., F.R.C.P.

F.R.F.P.S., D.C.H.

Professor of Clinical Medicine, University of Malaya,
Singapore

Among congenital heart diseases, tricuspid atresia is by comparison a rare complaint (White, 1951) and by itself is incompatible with life. Gibson and Clifton (1938) reported one case in a series of 105 congenital cardiac cases, Abbott (1936) had 16 out of 1,000 cases, and we had only one case out of 106 cases (Gwee, 1956, unpublished). However, compensatory mechanisms in the form of other coexisting congenital cardiac malformations sometimes enable a patient to live for many years. Taussig (1947) pointed out that the method of compensation may be either a communication of the two atrial chambers by an interatrial septal defect, the vestigial right ventricle receiving its blood supply through a ventricular septal defect, or persistent ductus arteriosus with interatrial defect and non-functioning right ventricle. White (1951) stated that tricuspid atresia is almost always attended by compensatory septal defects. Blackford and Hoppe (1931) regarded it as a functionally two-chambered heart.

Quite often the coexisting deformities produce no pathognomonic clinical signs, and in fact some cases may even be silent. Their presence is nevertheless suspected from the fact that the circulation must be able to complete its circuit to be compatible with life. The diagnosis is being made with increasing frequency in life, and in more and more cases knowledge of the coexisting defects is obtained pre-mortem by means of special investigations, of which angiocardiology and cardiac catheterization are probably the most informative.

Clinically, the patient is a cyanotic child with polycythaemia and gross clubbing of fingers and toes, and, cursorily inspected, may well pass for a case of Fallot's tetralogy (Alexander and White, 1947). A closer examination, however, may reveal in some cases a left ventricular apical impulse to suggest the true nature of the anomaly. Electrocardiography typically, though not necessarily, shows a left axis deviation and suggests left hypertrophy (Rihl, Terplan, and Weiss, 1929; Burchell, 1949). With the clinical appearance of Fallot's tetralogy, this finding is almost diagnostic of tricuspid atresia

(Astley, Oldham, and Parsons, 1953)—one of the few occasions when the electrocardiograph is of value in the elucidation of cardiac malformations. Radioscopy usually shows a large left ventricle, and the appearance may resemble that of truncus arteriosus, which, however, produced more cyanosis and is seldom seen beyond infancy. Also, the plateau-like appearance in truncus arteriosus is usually more marked, and the vascular pedicle remains the same size on rotation during radioscopy (Taussig, 1947). The diagnostic triad therefore consists of clubbing with cyanosis, left axis deviation in the electrocardiogram, and a characteristic appearance on radioscopy (Taussig, 1936). However, the shape may vary considerably (Astley *et al.*, 1953), and the pulmonary vascularization may be normal or markedly oligoemic. An exact diagnosis of all the coexisting anomalies may be impossible even after very exhaustive studies.

The salient differences of these cyanotic types of congenital heart disease are shown in the Table.

The life expectancy is usually short, although Abbott's (1936) series of 16 cases included a case of Hedinger's (1915) surviving up to 56 years. The review of Edwards and Burchell (1949) shows that the longevity of the patient depends to a large extent on the presence or absence of a transposition of great vessels. In their review, 45 cases were classified into four groups, thus: (1) No transposition of great vessels, (a) pulmonary

Signs and Symptoms

	Cyanosis	Clubbing	Polycythaemia	Stunting of Growth	E.C.G.	Pul. Vasc.	Death
Fallot's tetralogy	++	++	++	++	R.V.H.	Sparse	Often reach adult life
Eisenmenger's complex	Later	Later	++	May be little	R.V.H.	Normal	" "
Tricuspid atresia	++	++	++	++	L.V.H.	Sparse	Usually infancy and childhood
Truncus arteriosus	++++	++	++	++	L.V.H. or normal	"	" "

atresia, closed ventricular septum; (b) subpulmonary stenosis. (2) Transposition of great vessels, (a) pulmonary and subpulmonary stenosis; (b) no pulmonary or subpulmonary stenosis.

Of these, 28 cases belong to group 1b with subpulmonary stenosis and 4 to group 1a with pulmonary atresia. The life expectancy of groups 1a and 1b is stated to be quite short, the oldest being four years (Bellet and Stewart, 1933) and the causes of death being cardiac failure, bronchopneumonia, intracranial venous thrombosis, brain abscess, and bacterial endocarditis, in that order of frequency (Edwards and Burchell, 1949). The presence or absence of persistent ductus arteriosus and the adequacy of the pulmonary circulation do not seem to influence the length of survival.

Case Report

A male Chinese aged 20 was first seen on September 26, 1955, with a history that he had been breathless and blue as far back as he remembered, and that at about 5 years of age his fingers and toes began to get bulbous at the tips. Three weeks before admission he was short of breath and febrile. His legs and body became swollen.

On examination he was very cyanosed, with gross clubbing of fingers and toes. He was on the small side, height 4 ft. 8 in. (142 cm.), with no adult secondary sex

characteristics. His blood pressure was 134/102. The neck veins were engorged and pulsatile, the pulse was 90 a minute and regular. The apex beat was felt at the fifth left interspace, 3 in. (7.5 cm.) from the midline with a left ventricular impulse, and on auscultation both heart sounds were loud, with a rough systolic murmur, maximal at the left fourth interspace near the sternal margin. No thrill was palpable. The liver was enlarged two fingerbreadths below the costal margin and pulsatile.

Investigation showed polycythaemia of 7 millions with haemoglobin of 20.4 g. per 100 ml. E.C.G. showed left axis deviation. Screening and radiography confirmed the presence of a prominent left ventricle with poor pulmonary vascularization. A clinical diagnosis of tricuspid atresia was made, and angiocardiology subsequently showed direct passage of dye into the left auricle, causing it to opacify

Necropsy specimen showing probe in interventricular septal defect and a small vestigial right ventricle to the left of it, indicated by arrow.

early. The aorta was in the normal position, but the pulmonary artery was not seen. The case was regarded as one of tricuspid atresia with interatrial septal defect and no transposition of great vessels.

He was digitalized and kept resting in bed, with improvement, but cardiac failure occurred again in January, 1956, and this time the course was progressively downhill until he died suddenly on April 12, 1956.

Post-mortem examination showed an enlarged heart, mainly left ventricle. The tricuspid valve was completely atretic, and its position was indicated by a dimple on the right

auricle. A large interatrial septal defect 1.5 cm. in diameter, representing the persistence of septum secundum, connected the two auricles. The right ventricle was only a small cavity, measuring 1 by 0.5 cm., opening into the left ventricle through a small interventricular septal defect about 1 cm. in diameter (see Fig.). A normal pulmonary artery with valvular stenosis (valve opening 0.5 cm. in diameter) arose from the vestigial right ventricle. Ductus arteriosus was completely obliterated. The lung showed congestive changes with an infarct in the right lower lobe. Evidence of subacute bacterial endocarditis was found on the pulmonary valve cusps, which were grossly deformed but three in number. All other organs were normal except for a large thymus, which could possibly account for the observed deformity of the dye column across the mediastinum during angiocardiology.

Summary

A case of tricuspid atresia with pulmonary stenosis and without transposition of great vessels, surviving to the age of 20, is reported.

The special investigations and the post-mortem findings are correlated.

We are grateful to Dr. T. Balasingham for his necropsy report.

REFERENCES

- Abbott, Maude E. (1936). *Atlas of Congenital Cardiac Disease*. American Heart Association, New York.
 Alexander, F., and White, P. D. (1947). *Ann. intern. Med.*, 27, 64.
 Astley, R., Oldham, J. S., and Parsons, C. (1953). *Brit. Heart J.*, 15, 287.
 Bellet, S., and Stewart, H. L. (1933). *Amer. J. Dis. Child.*, 45, 1247.
 Blackford, L. M., and Hoppe, L. D. (1931). *Ibid.*, 41, 1111.
 Burchell, H. B. (1949). *Med. Clin. N. Amer.*, 33, 1157.
 Edwards, J. E., and Burchell, H. B. (1949). *Ibid.*, 33, 1177.
 Gibson, S., and Clifton, W. M. (1938). *Amer. J. Dis. Child.*, 55, 761.
 Hedinger, E. (1915). *Zbl. allg. Path. path. Anat.*, 26, 529. Quoted by Abbott (1936).
 Rühl, J., Terplan, K., and Weiss, F. (1929). *Med. Klin.*, 25, 1543.
 Taussig, Helen B. (1936). *Bull. Johns Hopk. Hosp.*, 59, 435.
 — (1947). *Congenital Malformations of the Heart*, Chap. 4, p. 79. Commonwealth Fund, New York.
 White, P. D. (1951). *Heart Disease*, 4th ed. Macmillan, New York.

COLD CONIZATION WITH SPENCER'S TRACHELOTOME FOR CERVICAL EROSION

A REVIEW OF 35 CASES

BY

R. G. EMERSON, M.B., B.A.O., D.Obst.R.C.O.G.

Major, R.A.M.C.; Late Gynaecological House-Surgeon,
Royal Salop Infirmary

Read (1955), referring to the problem of the unhealthy cervix, states: "It would appear obvious that in the case of all cervical lesions, before treatment is undertaken, the presence of malignant change must be excluded. Reliance on clinical appearance alone, except in minimal lesions, can be dangerous, and recent work has shown that even the clinically innocent-looking cervix can be the site of early malignant metaplasia."

Until the present day, cervical erosions have been treated in the out-patient departments of most centres in Great Britain by means of the electrocautery, while only in clinically suspicious cases have the patients been admitted to hospital for cervical biopsy as a prelude to cauterization. As a result cases are by no means unknown where an erosion has been cauterized on several occasions without healing taking place, followed by a belated biopsy examination showing the presence of a carcinoma of the cervix. Then, again, even in those cases which heal after cauterization it is certain that many instances of carcinoma-in-situ will be overlooked, with the result that invasive carcinoma may occur at a later date. It would therefore seem better that these patients should all have biopsies carried out as a preliminary to any form of treatment. More acceptable still would be a procedure, such as conization, where diagnosis and treatment are combined in one operation.

Historical

Conization of the cervix, as either a diagnostic or a curative measure, is not a new technique. As long ago as 1815 Lisfranc (1834), in the treatment of a case of suspected cervical carcinoma, is reported to have removed a wedge-shaped portion of the cervix which extended from the vaginal margin up as far as the internal os. Forty-five years later Hughier, of Paris, Marion Sims, and Thomas Emmett were all performing similar operations (Emmett, 1874). By 1916 Arnold Sturmdorf had devised an operation, the basis of which was conization followed by inversion of flaps of vaginal mucous membrane over the bare areas. In 1928 Mortimer Hyams introduced the electroconizer, the purpose of which was the removal of the endocervical mucosa. This instrument had an electrode only $\frac{1}{4}$ in. (3 mm.) wide, and was suitable for removing only a very limited portion of the cervix. The instrument was later improved upon by