After the two-year analysis reported here had been completed the first report of the Joint Committee of the Medical Research Council and the Nuffield Foundation was published (1957). In this carefully controlled trial prednisone comes out rather better than cortisone. In America, Black *et al.* (1955) conclude: "Prednisone has been found to be an effective antirheumatic agent in patients with rheumatoid arthritis whose response to other steroids was unsatisfactory. Its antirheumatic potency still appears to be three to four times that of cortisone or hydrocortisone. The return of rheumatic symptoms after 4 to 20 weeks of maintenance therapy may be controlled by a slight increase in the dosage."

The patients' assessment of their own improvement is often viewed with some scepticism, and the well-recognized response of many patients to a placebo justifies this. At the same time, in the everyday practice of medicine subjective improvement is only too often the sole criterion of successful therapy. The history of the treatment of rheumatoid arthritis is so beset with extravagant claims for dubious remedies that a healthy scepticism is desirable; but, like the tendency it counteracts, it too can be overdone. The conventional grades of functional status do not tell the whole story. We have been especially impressed by the fact that the improvement of some of our patients consists not so much in doing more than they did previously as in doing what they have always had to do more quickly, more comfortably, and more expeditiously. There is unfortunately no easy test for this kind of improvement. Without labouring the point, we believe that a report of subjective improvement supplemented by confirmatory answers to experienced questioning is of value.

Three fundamental questions, we believe, must be applied to any new drug used in rheumatoid arthritis: (1) Does it produce substantial relief in a high proportion of patients? (2) Is such relief maintained, and if so for how long? (3) Does the benefit obtained outweigh the risks and complications? From our experience with prednisone the answer to the first question is yes. In regard to the second, we believe that relief has been satisfactorily maintained in most patients so far, but it must be admitted that this belief depends in part on what our patients tell us, which may be subject to many fallacies. The answer to the third question at present is yes; further experience, however, could well cause revision of this opinion. If it be accepted that the newer steroids will help most patients for a time at least, the physician's problem is to know which patients it is justifiable to subject to the hyperadrenal state, however mild, with the risk of the attendant complications. We believe it justified in the cases of the wage-earner on the point of incapacity, the housewife unable to look after a young family, and the patient who is deteriorating rapidly despite all other measures. Whether steroids should be given to patients with milder disease or less arduous commitments cannot be decided until a great deal more experience throughout the world has been gained, and in fact may not need to be decided if research should produce a better and safer remedy.

Summary

The practice of a rheumatism clinic, relating to the treatment of 167 patients with rheumatoid arthritis seen over a two-year period, is reviewed.

Prednisone in a dosage of 15 mg. daily was given to 15 of these patients when first seen; by the end of two years 47 patients were receiving the drug, the increase being accounted for by the failure of other drugs.

The commonest complication of prednisone therapy has been dyspepsia, which occurred in 11 patients, two of whom developed peptic ulcers.

Improvement of functional status occurred in about 50% of patients given prednisone; subjective improvement was reported by 90%.

The value of, and indications for, prednisone in rheumatoid arthritis are discussed.

REFERENCES

- Black, R. L., Bollet, A. J., Segal, S., and Bunim, J. J. (1955). Presented before the First International Conference on Clinical and Metabolic Effects of Meticorten and Meucortelone, New York City, May 31 and June 1, 1955.
Bunim, J. J., Pechet, M. M., and Bollet, A. J. (1955). *J. Amer. med. Ass.*, 157, 311.
Copeman, W. S. C., Savage, O., Bishop, P. M. F., Dodds, E. C., Kellie, A. E., Stewart, J. W., Glyn, J. H. H., Henly, A. A., and Tweed, J. M. (1952). *Brit. med. J.*, 1, 397.
— Dodds, E. C., Glyn, J. H. H., and Fearnley, M. E. (1954). *Ibid.*, 1, 1108.
Joint Committee of Medical Research Council and Nuffield Foundation (1957). *Ibid.*, 2, 199.
Nabarro, J. D. N., Stewart, J. S., and Walker, G. (1955). *Lancet*, 2, 993.

SEA-SNAKE BITE

A SURVEY OF FISHING VILLAGES IN NORTH-WEST MALAYA

BY

H. A. REID, M.B., M.R.C.P.Ed., D.T.M.&H.

Specialist Physician

AND

K. J. LIM, M.B., M.R.C.P.Ed.

Formerly Medical Registrar

General Hospital, Penang, Malaya

In their vast habitat round the shores of the Indian and western Pacific Oceans sea-snakes abound. As they are all venomous and many are highly poisonous, it is fortunate they are not aggressive. Fishermen handle them daily, but there are few records of their being bitten. However, Smith (1926) maintained that casualties were not infrequent, the paucity of reports being due to the incidents occurring in remote localities. Responsible medical authorities rarely observe such cases.

As part of research on sea-snakes, their bites, and poisoning (Reid, 1956) we planned a survey of fishing villages in north-west Malaya to try to find out the incidence of bites, how victims were bitten, and their fate, together with any information of relevance or interest.

Area Surveyed

Fig. 1 indicates the recorded world habitat of sea-snakes (excepting one species, *Pelamis platurus* (Linn.), which extends to western America and eastern Africa). Inset is Malaya. In Fig. 2 the 17 villages visited are shown. They constitute most of the fishing communities of Penang Island and the adjoining 80 miles of coastline of the States of Kedah and Perlis. It was impracticable to include three small villages and the city of George Town, as fishing folk there are too scattered.

TABLE I.—Numbers of Sea-snake Bites per Village with Approximate Populations

Name of Village	Approximate Population	No. of Visits	No. of Sea-snake Bites			
			Fatal	Severe	Trivial	Total
1. Telok Bahang ..	2,000	3	2	—	—	2
2. Tanjung Bungah ..	2,500	2	2	4	4	10
3. Tanjung Tokong ..	5,000	2	2	1	2	5
4. Batu Maung ..	600	4	—	1	3	4
5. Damar Laut ..	1,100	2	—	2	—	2
6. Telok Kumbar ..	3,200	10	4	6	9	19
7. Gertak Sanggul ..	700	2	—	2	2	4
8. Pulau Betong ..	500	3	—	—	2	2
9. Jalan Bahru ..	800	4	3	4	7	14
10. Sungei Pinang ..	1,700	2	—	7	2	9
11. Panteh Acheh ..	800	2	2	3	2	7
12. Kuala Muda ..	3,000	6	11	5	4	20
13. Tanjung Dawai ..	1,700	2	2	—	1	3
14. Sungei Yen ..	4,200	2	2	3	—	5
15. Kuala Kedah ..	7,300	3	6	5	6	17
16. Sanglang ..	1,800	1	3	2	6	11
17. Kuala Perlis ..	3,200	3	2	3	5	10
	40,100	53	41	48	55	144

FIG. 1.—Recorded world habitat of sea-snakes (dotted area), with inset of Malaya.

No reliable population statistics exist for these villages. Approximate figures in Table I were obtained by the Health Department, Federation of Malaya, and include all ages, sexes, and races. Populations per village vary from 500 to 7,300, the average being about 2,500. In many of the larger villages a minority are fishing folk. Malays and Chinese are roughly equal in total numbers, although individual villages varied.

Methods

We visited the villages with a member of the Health Department known to the fishermen concerned. A bag containing a variety of species of dead sea-snakes and coastal water-snakes was opened in the appropriate coffee-shop (see Fig. 3). Every fishing village has one or more coffee-shops where fishermen congregate in leisure hours. The snakes always attracted a large audience, with whom we talked. If possible, we saw Chinese separately from

Malays, as this established better confidence. The trend of questions was purposely from generalities on sea-snakes to particulars of actual bites. A form on general information was personally completed in the coffee shop, or shortly afterwards if we thought its production would lessen co-operation. The reverse side was used for details of individual bites, these being obtained from victims or close witnesses in fatal cases. The survey was started in February, 1955, and during the ensuing 14 months 53 visits were made to 17 villages. The number per village varied from 1 to 10 according to accessibility and the response.

In the tables expressing our results, cases are divided into fatal, severe, and trivial. They are judged as severe if unequivocal

paresis was described and/or at least two days' work was lost owing to the bite. These criteria are not altogether satisfactory, because several victims stayed off work for two or more days, although they did not appear to have severe poisoning as judged by the symptoms related. Also, a few victims returned to work within 48 hours, but their cases might be judged severe—for example, having "bulbar" symptoms and trismus—rather than trivial, although they had no paresis. However, the classification has the merits of simplicity and clarity. The majority of trivial cases had no symptoms of poisoning.

Answers to all the questions were not always available, particularly in fatal cases where the witnesses did not know all the facts and in the less recent incidents where the victim had forgotten some of the details. Consequently several tables do not comprise the total 144 cases. All 144 were in our opinion unequivocal instances of sea-snake bite (see under "Symptoms").

Results

A total of 144 sea-snake bites was disclosed, the number per village being listed in Table I. Two victims were bitten twice, one three times, and another on four occasions. The response was remarkably uniform. The villagers talked freely about sea-snakes, and for the most part were remarkably well informed. But immediately the topic of bites was approached the attitude changed and all knowledge of victims in that district would be denied. Eventually, cases in a neighbouring village would be admitted. Later, Chinese would recall Malay victims in their own village (if we were talking with Chinese, and vice versa), then instances amongst their own race, and finally themselves. This great reluctance to talk about bites is mainly due to superstitions. It is the chief reason for failure to realize the frequency of bites among fishing folk. Also, victims may forget having been bitten, especially if trivial or no symptoms of poisoning ensued, which seems to be a common experience. This is borne out by Table II, in which more cases are conceded each year, reaching a maximum in 1955. Moreover, there is a disproportionate increase of cases with trivial or no symptoms of poisoning in recent years. It is unlikely that the sea-snakes have altered their toxicity or habits in the space of twenty-odd years. Reluctance to talk is more likely

FIG. 2.—The 17 fishing villages visited in the survey.

TABLE II.—Years in Which Sea-snake Bites Occurred. There is a Steady Increase in Recent Years, Particularly of Trivial Cases

Year	Sea-snake Bites			
	Fatal	Severe	Trivial	Total
1945 or before ..	16	14	12	42
1946-50 ..	10	15	6	31
1951 ..	—	2	2	4
1952 ..	3	—	2	5
1953 ..	—	5	4	9
1954 ..	4	5	4	13
1955 ..	8	7	25	40
	41	48	55	144

to affect recent rather than remote cases. We therefore conclude that a significant number of victims forget the incident.

A third factor reducing the reliability of our figures in estimating the incidence is the number of people with whom we talked in each village. It was impracticable to talk with more than about a dozen villagers at each visit. Many victims may have existed in a village, unknown to the people whom we interviewed. To counteract this we visited the headman of each community on our first visit. On leaving we asked the headman and any other people with whom we had talked to spread the news around the village that we were interested in sea-snakes and their bites for research pur-

FIG. 3.—Collection of sea- and water-snakes in a fishing village coffee-shop always attracted an audience and stimulated discussions. The sea-snake being held up is an *Enhydryna schistosa*.

poses and that we would like to talk personally to as many victims as possible. We repeatedly reassured them that this was our sole purpose, that we did not intend to take blood from any victims or suggest their admission to hospital (we discovered that these were common fears even though they had been bitten years previously). On subsequent visits we followed up as many cases and as much partial information as possible. In the smaller villages we think we saw a representative number of the inhabitants, but in the few larger ones it is probable a significant number of victims who would have been willing to admit their bites were not seen.

Sex, Age, and Race.—In Table III details of sex, age, and race are given. As would be expected, about 90% are males. The age incidence also is merely a reflection of the

TABLE III.—Distribution of Sea-snake Bites According to Sex, Age, and Race

	Sea-snake Bites			
	Fatal	Severe	Trivial	Total
Male	34	45	49	128
Female	7	3	6	16
Age in years:				
5-9	1	2	1	4
10-19	9	4	4	17
20-29	9	10	11	30
30-39	11	14	14	39
40-49	3	10	9	22
50 onwards ..	5	6	9	20
Not known ..				12
Malay	22	20	26	68
Chinese	17	26	28	71
Other	2	2	1	5

groups at risk. The racial incidence and mortality are approximately equal for Malays and Chinese with the exception of the relatively large number of Malay children bitten compared with Chinese. Of 14 victims aged 14 years or less, 12 were Malays (including all seven fatal cases). The difference is probably due to Malay children wading and playing in river-mouths, a habit less favoured by Chinese.

Circumstances of the Bite.—Details of the activity leading to the bite, and location of the victim at the time, are recorded elsewhere (Reid, 1956, Table I). Handling nets accounted for nearly half the bites, the sea-snakes escaping their scrutiny. Further particulars about the nets and fishing methods are mentioned below under "General Aspects." One-third were bitten washing in a river-mouth or wading, usually to or from their boat. They probably trod on the snake—a matter of bad luck considering how often fishing folk have to pursue such activities. River-mouths, particularly those with fishing villages, are favourite haunts of sea-snakes. Five victims were bitten while swimming in a river-mouth. We were careful to explain this as meaning "feet off the river-bed." One victim felt the snake coiling round his forearm and biting his hand. Trivial poisoning followed. Two severe bites were in the thigh and forearm respectively, one swimmer was fatally bitten in the hand, and in another fatality the site of the bite was not known. Presumably these five were unlucky enough to hit an on-coming snake.

Of 52 incidents out at sea, only 10 were fatal, whereas 22 of the 50 bitten near the shore or in a river died. However, we make no conclusions from these figures regarding the respective danger of these locations. There was no evidence that the season or weather influenced the frequency or mortality of bites. One-third were bitten at night, but only 10 were engaged in activities such as sorting fish or nets, where lack of light might be a contributory factor to the bite. Twelve victims were bitten whilst engaged in similar activities during daylight. One victim aged 16 had established a reputation for playing with sea-snakes. He was fatally bitten picking up a *Hydrophis* 30 cm. long.

Identification of the Sea-snake.—On asking victims or witnesses (of fatal cases) to identify the snake from the collection presented at the survey visit, many were unable to do so. Table IV lists 65 incidents in which the species was

TABLE IV.—Offending Sea-snake Species Identified by Victims or Witnesses (in Fatal Cases) from Collection Presented at Survey. Species Frequency and Mortality Correspond Well with Experimental Findings

Sea-snake Species	Sea-snake Bites			
	Fatal	Severe	Trivial	Total
<i>Enhydryna schistosa</i> (Daudin) ..	7	11	6	24
<i>Hydrophis cyanocinctus</i> (Daudin) ..	4	—	5	10
" <i>spiralis</i> (Shaw) ..	2	3	3	8
" <i>klossi</i> (Boulenger) ..	1	5	10	16
<i>Kerilia jerdoni</i> (Gray) ..	—	—	2	2
<i>Lapemis hardwickii</i> (Gray) ..	1	1	—	2
<i>Microcephalophis gracilis</i> (Shaw) ..	1	1	1	3
	16	22	27	65

identified. The findings should be regarded with due reserve, but it is interesting to note the remarkably close correspondence between the frequency of alleged offending species and its occurrence in the area as judged by 600 specimens recently collected (Reid, 1956). Even more arresting is the accuracy with which the fatalities correspond to experimental findings. Thus *Enhydryna schistosa* and *Hydrophis cyanocinctus* have a relatively high mortality compared with *Hydrophis klossi*. Experimentally the venom yield of *H. klossi* is so small that it is unlikely to kill adult men. In two instances an *E. schistosa* was described as clinging like a limpet, having to be pulled off. One victim died, but the other recovered after severe poisoning.

Bite Marks.—Among 69 victims, search found no marks in three instances (one fatal, one severe, one trivial), and in one severe case only a "graze" was seen. One puncture mark was noted in four cases, two marks in 33 cases, three in 8 cases, four in 7 cases, and multiple marks in 13 cases. There was no correlation between the number of marks seen or the distance between them and the degree of poisoning. All bites were painless after the initial prick, and no swelling, prolonged oozing, or local reaction was observed. On

22 occasions teeth or fangs were extracted by a hair or needle from the site of the bite. In four instances only one fang was withdrawn, in eleven instances two fangs were withdrawn, in three instances three teeth, in one instance four, and in three cases it was claimed that 12, 18, and 23 teeth were withdrawn. Attempts were unsuccessful in nine instances. Fishermen regard this manoeuvre as diagnostic and therapeutic, but there was no correlation between the number extracted and the fate of the victim or the species of the alleged offending sea-snake.

Site of the Bite.—Table V lists the site of the bite in 112 cases. As is usual in snake bite, two-thirds were bitten in the lower limb because one way or another the victim treads on the snake. Acton and Knowles (1914) maintained that

TABLE V.—*Site of Sea-snake Bite. The Low Mortality in Bites on Fingers or Toes is Noteworthy*

Site of Bite	Sea-snake Bites			
	Fatal	Severe	Trivial	Total
Finger	—	6	12	18
Hand	5	1	7	13
Forearm	1	1	—	2
Arm	—	1	1	2
Toe	1	10	3	14
Foot	10	10	14	34
Leg	5	9	4	18
Thigh	5	4	2	11
	27	42	43	112

bites of elapid and hydrophine snakes, by reason of their short, fixed fangs, were usually ineffective unless a good grip was obtained, such as on a finger or toe. The findings in the survey point to the opposite conclusion. Thus, of 32 victims bitten on a finger or toe, only one died, whilst of 80 bitten on the hand, foot, or more proximally, 26 died.

Latent Period and Death Time.—The latent period between the bite and the onset of symptoms varied from five minutes to eight hours. The duration of this period and of the death time (time from the bite to death) is sometimes regarded as being proportional to the dose of venom injected. The higher the dose the shorter these periods become (intravenous injection, which would greatly shorten the latent period, probably never occurs in sea-snake bite, as the fangs are too short). However, our findings do not support this alleged dose/latent period/death time relationship. Although the average latent period in 13 fatal cases was 1.4 hours compared with 1.8 hours in 28 severe cases, the variation in both categories was up to eight hours. No correlation was found between the latent period and death time in 11 fatal cases, as follows :

Latent Period	Death Time	Latent Period	Death Time
1 hour ..	6, 36, 42 hours	3 hours ..	25 hours
1 ..	4, 4, 21 ..	4 ..	21 ..
2 hours ..	24, 25 ..	8 ..	20 ..

Similarly, no correlation was found between the latent period and the recovery time in either severe or trivial cases. One victim had no symptoms until eight hours after the bite, yet died 12 hours after they started. The duration of the latent period is therefore not reliable in prognosis.

Symptoms

Table VI gives the approximate percentage of symptoms occurring in 125 cases. The clinical paradox of trismus and flaccid limb weakness is an outstanding feature of sea-snake poisoning, and the characteristic triad in severe or fatal cases is trismus, flaccid body paresis, and "haemoglobinuria." Among fatalities the proportion with red or black urine was only 35%, but many died without urine being passed or its colour noted. One-quarter of the severe cases also had red or black urine. For some time this appeared to be another paradox, because, experimentally, sea-snake venoms are only feebly haemolytic. Theoretically, haemoglobinuria should indicate an enormous dose of venom, inevitably fatal. However, recent clinical research

TABLE VI.—*Percentage Incidence of Symptoms in 125 Cases. Of the 55 Trivial Cases 32 Had No Symptoms of Poisoning At All*

Symptom	Percentage with Symptom		
	Fatal Cases	Severe Cases	Trivial Cases
No. of cases	22	48	55
Paresis	90	70	6
Trismus	85	60	7
Breathing change ..	70	25	4
Dysphagia	60	35	—
Ptosis	55	21	2
Cyanosis	55	6	—
General stiffness ..	50	65	4
Increased sweating ..	50	15	—
General coldness ..	45	35	22
Thirst	40	15	6
Red or black urine ..	35	25	—
Dysphasia	35	17	2
Numbness	20	4	4
Dry throat	15	15	6
Vomiting	10	—	—
Cheek swelling	10	2	—
Convulsions	10	—	—
Face palsy	10	—	—
Difficult cough	5	4	—
Unconscious	—	8	2
Drowsy	—	—	2

has shown this pigment to be myoglobin, not haemoglobin, thus resolving a number of enigmas. At the time of the survey we did not realize the significance of another clinical observation—namely, the severe muscle pains and tenderness which our hospital patients have suffered. This important symptom does not therefore feature on the list.

From the survey four features would appear to have bad prognostic significance—increased sweating, difficulty in swallowing, ptosis, and breathing change. Coupled with breathing changes are, of course, other signs of respiratory insufficiency such as cyanosis and convulsions. These are terminal. Increased sweating may also occur in respiratory failure, but the symptom as referred to here is an early one. It may be noted before any paresis is evident.

Among 44 severe cases, 24 victims returned to work within a week, 12 within one to two weeks, and 8 within one to three months. If paresis was mild, complete recovery occurred within a few days, but if it was moderate or severe it usually took several weeks. There may be total paralysis as related dramatically by one fisherman who six hours after the bite was unable to speak, swallow, or move at all (including his eyelids). He was presumed dead and had the unpleasant experience of hearing his coffin being brought into the room and his funeral discussed. However, someone felt his pulse, so he escaped being buried alive. Full recovery took three months.

Of the 55 bites classified as trivial, only 23 produced symptoms of poisoning, the commonest being a feeling of general coldness. Thirty-two, or almost one-quarter of the total, had no symptoms of poisoning at all. These were unequivocal sea-snake bites. In each instance either the sea-snake was seen or the bite took place out at sea, where water-snakes do not occur, and the characteristic dot-like marks were painless after the initial prick.

Treatment

About half the victims had a tourniquet applied, but many ligatures were of unsuitable material or were applied too close to the bite for them to be effective in delaying absorption. In the absence of effective antivenene, immediate amputation is the only first-aid measure likely to help the patient (Reid, 1956). In the survey, however, nobody knew of any victim trying it. Amputation is only practicable in finger or toe bites. Table V shows that if it had been universally practised 31 victims would needlessly have lost a finger or toe and one victim's life might have been saved. Fishermen were unanimous in saying they would rather risk death than lose a finger or toe. It is doubtful if propaganda would persuade them otherwise even if pursued with missionary zeal. Only seven had excised the site of the bite

(one fatal, from the arm; two severe, from the foot; four trivial, one from a toe and three from a finger). Judging from the victims' descriptions and the six scars seen, it is unlikely that the excisions were sufficiently radical to affect the issue. There were no instances of incision and suction being attempted.

Thereafter the procedure followed depended on the victim's race. If Chinese, herbal brandy made from a variety of herbs in rice wine was almost invariably taken. It was also customary to avoid smoking, talking, and eating for 24 hours after the bite (a rational prophylactic procedure considering the possibility of "bulbar palsy," inhalation of vomit, etc.). Malays, on the other hand, would usually consult a *bomor* (medicine-man), who relied mainly on incantations. Malays and Chinese usually disparaged the other's methods and alleged a resulting high mortality, but the survey figures did not support either advocate. Only 18 of the total 144 sought orthodox medical help in hospital or dispensary. Despite this small proportion, many said they would come to hospital if they knew a "special injection" were available for sea-snake bite. This opinion is supported by the fact that, although a specific antivenene is not yet available, the numbers of sea-snake victims admitted to hospitals in north-west Malaya steadily increase each year.

General Aspects and Folklore

Except in one village, fishermen said they encountered sea-snakes every time they fished. They were more numerous in January-March, specially during or after stormy weather. No general change in the frequency or habits of sea-snakes was known to have occurred during the villagers' lifetime. The number of snakes varied according to the method of fishing, from nil up to a total of 100 or more each day. The first action on hauling in the catch is to spot sea-snakes, pick them up quickly by the tail, and throw them overboard. This responsibility usually rests on one member of the crew, who becomes adept. Malays often used a stick rather than fingers for lifting the snakes. The sea-snakes are never killed. Line fishing is not common in the area surveyed, but an average of 20 sea-snakes per 300-hook line was quoted by two fishermen using this method in shallow water. Usually they cut off the line concerned, but one victim was bitten trying to disengage a sea-snake and thereby save the hook. He had trivial symptoms following the bite, but thereafter conformed to custom, cutting the lines with sea-snakes.

As would be expected, sea-snakes were very rare in entangling structures (gill nets). They were not numerous in engulfing structures (drive-in nets), and some entrapping structures (palisade traps, lift nets, falling nets). But a gape net known by the Chinese as *ch't cheh* invariably caught 10-20 sea-snakes each lift. A conical net with the mouth held open by two poles stuck into the sea-bed and the tail raised by floats is used in shallow water and lifted two or more times daily according to the tide. However, although this net invariably traps large numbers of sea-snakes, only 4 of the 50 victims bitten whilst handling nets were using this method. The other net in which sea-snakes are numerous is a small drag seine called a *kesa* by the Chinese. It also is used in shallow water, being dragged along by two or three fishermen who are in the sea. Twenty-seven of the 50 were bitten handling a *kesa*, 16 (6 fatal, 5 serious, 5 trivial) being in the sea at the time and 11 (7 serious, 4 trivial) in the boat pulling in or sorting the net.

All species of sea-snake presented at the survey visit were recognized at one or more villages. Much the commonest was *E. schistosa*, and thereafter in descending frequency, *Hydrophis spiralis*, *H. cyanocinctus*, *Lapemis hardwickii*, *H. klossi*, and *Serilia jerdoni*. In contrast, *Hydrophis melanostoma* Günther, *Microcephalophis gracilis*, and *Praescutata viperina* (Schmidt) were rarely recognized. The harmless non-fanged water-snake *Acrochordus granulatus* (Schneider) was universally mistaken for a sea-snake and regarded as highly poisonous. As it is often caught with sea-snakes and has tapering bars, slightly flattened tail, and

narrow ventral scales the mistake is understandable. Generally speaking, *E. schistosa* was considered the most dangerous. All fishermen agreed that sea-snakes never attacked human beings spontaneously. The only exception was one man who claimed to have been chased by an *E. schistosa* whilst in a fish-trap. But others present laughingly disbelieved him. Several fishermen recounted sea-snakes swimming between their vest and body: all would be well so long as they kept still. As one fisherman put it, if sea-snakes were aggressive there would be no more fishing.

According to Herre (1942) sea-snakes are inclined to be aggressive during the breeding season, but this was not confirmed by the fishermen. However, they appeared to know little or nothing about the breeding habits, season, etc., of sea-snakes. It was agreed that sea-snake bite was commoner in some river-mouths. Lowered salinity, making the snakes more aggressive, and available food increasing their frequency, were two reasons put forward, but the majority admitted ignorance of the cause. Several species were known to swim up river, particularly *E. schistosa*, and one fisherman thought the latter accounted for the frequency of river-mouth bites. In captivity it is certainly the most aggressive species of those mentioned above (Reid, 1956).

In contrast to *A. granulatus*, water-snakes such as *Cerberus rhynchops* (Schneider) were distinguished by all as harmless. Everyone knew their round tapering tails, and many mentioned the differences in the shape of the head and the belly—the broad scales being quite different from those of sea-snakes. These water-snakes were common in river-mouths and shores. Children played with them in every village, many being adept at catching them. People were bitten quite often. The majority admitted they could not distinguish such bite marks from those of a sea-snake (our cases of bites with no symptoms of poisoning were not due to water-snakes for the reasons already given). Two fishermen said the water-snake marks were semicircular and more teeth were left in. One of us has been frequently bitten by specimens of *C. rhynchops* in captivity, and teeth have been left in on almost every occasion. It is a back-fanged snake which presumably ejects venom only on swallowing.

All fishermen were well acquainted with the commoner symptoms of sea-snake poisoning. It was recognized that many victims developed few or none of these symptoms, but this was invariably regarded as being due to the treatment received (a comforting outlook often shared by the medical profession in relation to snake bite). Deaths were attributed to disregard of advice or delay in receiving treatment. Land-snake bites were much less common in these villages. Offered the alternative of a cobra or a sea-snake bite, preference for the former was unanimous. Similarly, they laughed when asked whether they would rather be stung by catfish, sting-ray, or jellyfish in preference to a sea-snake bite. Anybody who has had the extremely painful and unpleasant experience of these fish stings will realize the significance of this laughter. Fear of sea-snakes was admitted by all—"more than anything else," as one fisherman put it. Another philosophically likened them to war—both the inevitable lot of mankind. Night-time, and the consequent difficulty in seeing the snakes, was particularly feared (although analysis of the survey figures did not suggest this is an important factor in sea-snake bite). These fears are prevalent despite the vaunted confidence in their respective treatments. There is no doubt that if these fishing folk knew a specific antivenene were available much of their fear would be abolished.

Superstitions and Myths

The reluctance to admit to specific instances of bites is in great contrast to the freedom with which they discuss general aspects of sea-snakes. One reason for this reluctance is suspicion. Deaths due to sea-snake bite are often notified as "fever," etc., in order to avoid necropsy or delay in burial (Reid, 1956): many suspected we intended to make trouble on this account. But the most important factor was superstition. Most Malays admitted frankly to these super-

stitutions, whereas many Chinese scoffed at them. However, we received the impression that the Chinese were in reality just as deeply under their influence as the Malays. Four common superstitions are: (1) Talk about sea-snake bites offends the sea-snakes and may change their timidity to aggression. (2) If a victim talks about his bite he will be bitten again. (3) If a pregnant woman hears of a sea-snake bite the victim will die. (4) If faecal droppings of a house lizard fall on a victim he will die.

The first three discourage the victim from broadcasting his misfortune, whilst the fourth superstition precludes his being treated in his home. In practice, the victim tells his best friend to fetch the *bomor* (medicine-man). If all goes well, no one else in the village is informed of the incident. But, as one *bomor* rather cynically admitted to us, if the victim is not satisfied and sends his friend for a second *bomor*, the first one spreads the news so that a pregnant woman will hear of it. The discredit of the inevitable death will then fall on the second *bomor*.

Other common beliefs include: (5) The presence of a *bomor* in a village attracts sea-snakes and causes more bites. (6) If fresh lime is taken shortly after a bite the symptoms will recur and probably be fatal. (7) Drinking blood of a sea-snake or eating its tail gives protection against poisoning. A few Chinese fishermen admitted doing this, and others sometimes ate them for general medicinal effect. None ate sea-snakes as ordinary food, although one Chinese said his father had been fond of them. Malays believed eating sea-snakes would cause shedding of their own skin. (8) Both Chinese and Malays had heard of the *ular belerang*, a sea-snake so poisonous that, if it bites the rudder, all in the boat will die. A few claimed to have seen this snake, always far out in deep water of more than 20 fathoms. They described it as about 30 cm. long, 2 cm. thick, usually a striking yellow colour with a longitudinal black stripe (*belerang* is Malay for sulphur). The descriptions suggest a young *P. platurus*. Experimentally its venom is as toxic as that of *E. schistosa*, perhaps more so.

(9) Originally, sea-snakes were harmless and pythons were the most poisonous snakes in the world. The pythons used to live in a certain well where they feasted on the fish. One day, in the absence of the pythons, a fisherman caught all the fish and took them away. A crow watching from a near-by tree broke the news to the returning pythons, pointing out the footprints of the marauder. The poisonous pythons were so incensed that they bit one of the footprints and the fisherman promptly died. At his funeral feast the villagers ate the fish he had caught. The crow then jeered at the hungry pythons: though their poison was so potent it could kill a man by biting his footprint, it was not powerful enough to recover the fish. Mortified with shame at this revelation, the pythons swam far out to sea and vomited up nearly all their poison. Returning, they brought up the rest of their venom near the shore and thus became non-poisonous. The *ular belerang* inhabiting the deeper seas swallowed most of the pythons' venom, while coastal sea-snakes (the majority) acquired the rest. This is why the *ular belerang* is now the most poisonous snake in the world.

Discussion

A survey such as this can only indicate trends. We would emphasize that the results by themselves can form no basis for generalizations. The individual figures expressed in the various tables must all be regarded with due reserve.

We obtained records of 102 sea-snake bites occurring in the 10-year post-war period. Because of the reluctance of the fishermen to disclose bites, the number forgotten, and the relatively small proportion interviewed in the larger villages, we consider this is a gross underestimate of the true incidence in the area surveyed. Of these 102 bites, 40 occurred in 1955. In this year also, 30 victims were admitted to hospital from Kuala Perlis village, representing 1% of the population at risk (in 1956 admissions rose to 51). Our

1955 survey total for this village was only six, of whom five were admitted to hospital. Thus we have survey or hospital records totalling 65 sea-snake bites in north-west Malayan fishing villages during 1955. Kuala Perlis and roughly half the villages surveyed are situated at a river-mouth, and in them the number of bites was consistently higher than in the other villages. Several river-mouth villages recounted many more bites than in Kuala Perlis. We consider a yearly incidence of $\frac{1}{4}$ –1% of the population at risk in river-mouth villages and $\frac{1}{4}$ – $\frac{1}{2}$ % in the rest would be a conservative estimate. If this is correct, there is a total of at least 150 victims of sea-snake bite each year in the villages surveyed. These contain about one-tenth of the Malayan fishing population.

On the east coast of Malaya the incidence may well be similar, as during 1955–6 30 victims (with 4 deaths) of sea-snake bite were admitted to Kota Bharu Hospital from scattered fishing communities of about 10,000 population (Dr. Thelma M. Ward, personal communication). It is known that some victims do not come to hospital or dispensary. If we assume this rough approximation to be reasonable, is it representative of the general incidence amongst the Asian fishing community? The two species *E. schistosa* and *H. cyanocinctus* abound virtually throughout the area shown in Fig. 1. Presumably they may bite fishing folk there as often as those in the north-west Malaya. On the other hand, different fishing methods may greatly decrease the risks. Also, regional changes in habits, toxicity, etc., of the sea-snakes may occur. On the 1951 Galathea Expedition, H. Volsoe (personal communication) noted that specimens taken in the Malacca Straits were far more aggressive than those captured off the Indian coast.

As already recorded (Reid, 1956), the mortality in the survey is not regarded as representative of sea-snake bite in general. In our opinion the true figure lies between 25% and 3%, being nearer 3%. If the potential danger of *E. schistosa* to man as judged by venom yield and toxicity is about the same as that of the Indian cobra and the Australian tiger-snake the mortality of sea-snake bite is low. Fairley (1929) quoted the mortality of tiger-snake casualties as 40%. The relatively low mortality from sea-snake bite may be due to most victims being bitten by species much less toxic than *E. schistosa*. One of us believes it is unusual for poisonous snakes to inject much venom when, as in human subjects, the bite is defensive (Reid, 1956). However, it must be admitted that mortality rates, indeed all statistics in snake bite, are highly suspect. Apart from the mortality—which is negligible compared with such "killers" as tuberculosis—fear of sea-snakes is an important feature in the lives of Asian fishermen. We consider that on this account alone it is well worth while undertaking research to improve the treatment of sea-snake bite and poisoning. Unlike land-snakes, an abundant source of sea-snakes is constantly available, and adequate venom could be obtained if a sea-snake station were to be set up as proposed recently (Reid, 1956). The basic toxicology of sea-snake poisoning could then be elucidated and rational treatment, including an effective polyspecific antivenene, should ensue. We are confident such measures would be welcomed by the fishing folk and would greatly increase their security and happiness.

Summary

A survey of fishing villages in north-west Malaya indicates a high incidence of sea-snake bites. It is estimated that a total of 150 victims are bitten each year in the villages surveyed. The latter contain about one-tenth of the fishing population of Malaya, itself a minute area amongst the world habitat of sea-snakes.

Relevant findings from the 144 sea-snake bites recounted in the survey are presented. Some of the more pertinent or interesting items of general knowledge, folklore, and superstitions amongst the fishermen are described.

The mortality of sea-snake bite appears to be low, but fear of them is universal and a potent source of worry. Effective treatment, including a polyspecific antivenene, would be a great boon to Asian fishing folk.

We wish to thank the State Surgeon, Kedah and Perlis, the Chief Medical Officer, Penang and Province Wellesley, the Director of Fisheries, Federation of Malaya, and many members of their departments for helpful co-operation in the survey; and Mr. C. J. Sundram for the maps.

REFERENCES

- Acton, H. W., and Knowles, R. (1914). *Indian J. med. Res.*, 2, 46.
 Fairley, N. H. (1929). *Med. J. Aust.*, 1, 296.
 Herre, A. W. C. T. (1942). *Copela*, p. 7.
 Reid, H. A. (1956). *Trans. roy. Soc. trop. Med. Hyg.*, 50, 517.
 Smith, M. A. (1926). *Monograph of the Sea Snakes*. London.

PROPHYLACTIC USE OF OXYTETRACYCLINE FOR EXACERBATIONS OF CHRONIC BRONCHITIS

BY

P. C. ELMES, B.M., M.R.C.P.

C. M. FLETCHER, C.B.E., M.D., F.R.C.P.

AND

A. A. C. DUTTON, B.M., B.Ch.

From the Departments of Medicine and Bacteriology,
 Postgraduate Medical School of London

Investigations both here (Elmes *et al.*, 1953) and elsewhere (Mulder *et al.*, 1952; Stuart-Harris *et al.*, 1952; May, 1953) have shown that exacerbations of chronic bronchitis are usually associated with proliferation in the sputum of one or more of a few species of bacteria which have become recognized as pathogens. Of these, *Haemophilus influenzae* is the organism most often found. *Streptococcus pneumoniae* is the next most frequent, while beta-haemolytic streptococci, *Staphylococcus pyogenes*, and *Klebsiella friedländeri* are occasionally found. Exacerbations are usually sequels of upper or lower respiratory infections by a virus or follow exposure to cold or to smoky fog. Subsequent bacterial invasion is thought to prolong exacerbations, to increase their severity, and to play an important part in causing permanent damage to the lungs (Reid, 1954).

Since wide-spectrum antibacterial drugs are effective against these organisms they should be able, given early enough, to prevent bacterial multiplication and its consequences. There is evidence (McVay and Sprunt, 1953; Helm *et al.*, 1956; May and Oswald, 1956; Edwards *et al.*, 1957) that the tetracycline group of drugs given continuously over long periods may control persistent bronchial infection and reduce the frequency and severity of acute exacerbations. These drugs are, however, too expensive for continuous use among the vast numbers of patients with chronic bronchitis in this country, most of whom have only one or occasionally two exacerbations each winter, between which they are able to continue at work. The trial described in this paper was to determine whether a short course of oxytetracycline given at the first sign of the onset of an exacerbation could reduce its duration, and thus provide an economic method of reducing loss of working time.

Clinical Method

The criteria for selection of patients for the trial were that they should be under the age of 65, be in regular employment, have had a productive winter cough for not less than three years, during which time they had had at least two illnesses with purulent sputum, causing loss of time from work. The patients had no other disabling disease. Twenty-one of them were attending a bronchitis clinic; other cases were referred by local general practitioners at our request. Of 125 cases so referred, 67 were selected (16 failed to attend and 42 were unsuitable in one way or another). Eighty-eight patients were therefore admitted to the trial at various times between November, 1954, and November, 1955. Special clinics were held on Saturday mornings to reduce loss of time from work. Each patient was seen at approximately monthly intervals in the winter and at two to three-monthly intervals in the summer.

It was decided to make the trial a "double blind," by allocating oxytetracycline tablets to some patients and indistinguishable dummy tablets (containing lactose) to the others, neither doctors nor patients knowing which was which. This was achieved by allocating oxytetracycline and control tablets in equal proportions to the serial numbers 1 to 99, using Fisher and Yates's (1948) table of random numbers. The patients were each given a serial number in the order in which they entered the trial and were prescribed tablets of the same number. The key list of numbers was held by the hospital's chief pharmacist, who dispensed 28 oxytetracycline or control tablets according to the serial number of each patient. Table I shows the distribution of

TABLE I.—Comparison between the Oxytetracycline and Control Groups before the Trial

	Oxytetracycline	Control
Males	37	37
Females	5	9
Mean age and range	54.6 (31-65)	53.5 (28-65)
Mean dyspnoea grade and range*	2.289 (1-4)	2.283 (1-4)
Time lost in days during 18 months before trial	76.6 (0-560)	71.24 (0-547)

* Grades ranging from 1 (normal) to 5 (breathless on least movement) (Fletcher, 1952). The figure used was the grade of dyspnoea noted by the patient when he was feeling well.

patients in the oxytetracycline and control groups according to age, sex, degree of respiratory disability, and amount of time lost from work during the 18 months preceding their admission to the trial. The two groups were similar except for a small preponderance of females in the control group.

On their first attendance a full clinical history was taken from the patients and the results of examination of their cardiovascular and respiratory systems were recorded. They were then given the following instructions:

"If you have a head cold, increased difficulty in breathing, increased cough and phlegm, or any other chest symptoms which you think may indicate the beginning of a chest illness, take one tablet after breakfast, dinner, tea, and supper—that is, four tablets in the day. Continue with the tablets for at least five days at first. If you are then quite well again, stop the tablets. If not, continue taking four daily until they are finished. If you develop looseness of the bowels or diarrhoea while you are taking the tablets take three tablespoonfuls of the medicine† daily. If diarrhoea continues stop the tablets. If, in spite of taking the tablets, your chest gets worse, consult your doctor. Please put a specimen of your phlegm into one of the containers: (a) before you start taking the tablets, (b) when you have been taking the tablets for five days, (c) five days after you stop the tablets. The specimens should be brought to the bacteriology department of the hospital on the same day either by yourself or a friend. Please report to the clinic as soon as you are well again."

On subsequent visits to the clinic details of the onset, symptoms, and duration of any chest illnesses were recorded, further supplies of tablets were given, and further clinical examinations were performed if necessary. Many patients

†Chalk and opium mixture (N.F.).