Figure 2 Ulcer in the plantar aspect of 2nd and 5th toes (A). Longitudinal incisions were made on the dorsum of the foot with corrugated drains inserted (B).

damage to the digital neurovascular bundle. The infected bones are excised and all septic tissue carefully debrided. The wound is washed, and the skin closed with interrupted sutures over a corrugated drain. The foot is dressed lightly with gelonnet, wool and crepe bandage.

Postoperatively, the patient should continue antibiotics and remain on bed-rest for 5 days, before being discharged with special well-padded surgical footwear.

DISCUSSION

This technique saves the toe and heals rapidly. The conventional ray amputation often involves the sacrifice of healthy tissue, at the expense of a small area of ulceration next to the infected MTP joint. The mobile flail joint created quickly fills up with granulation and fibrous tissue, allowing a reasonable range of movement. This operation is useful in neuropathic diabetic foot ulcers over the MTP joint, where the blood supply is good, or when peripheral arterial disease has been rectified.

References

1. Grimm A, Kastenbauer T, Sauseng S, Sokol G, Irsigler K. Progression and distribution of plantar pressure in Type 2 diabetic patients. *Diabetes Nutr Metab* 2004; **17**: 108–13.
2. Boulton AJ. Pressure and the diabetic foot: clinical science and off-loading techniques. *Am J Surg* 2004; **187** (Suppl): 17S–24S.
3. Ferrari J, Higgins JP, Prior TD. Interventions for treating hallux valgus (abducto-valgus) and bunions. *Cochrane Database System Rev.* 2004; (1): CD000964.

A new technique for reduction of paediatric femoral fractures using elastic stable intramedullary nails

AM WAINWRIGHT, UG NARAYANAN

Division of Orthopaedic Surgery, The Hospital for Sick Children, Toronto, Ontario, Canada

CORRESPONDENCE TO

Unni G Narayanan, Consultant Orthopaedic Surgeon/Assistant Professor, Division of Orthopaedic Surgery, The Hospital for Sick Children, 555 University Avenue, S-107, Toronto, Ontario M5G 1X8, Canada

T: +1 416 813 6432; E: unni.narayanan@sickkids.ca

BACKGROUND

Flexible intramedullary nails have revolutionised paediatric femoral fracture management.^{1–4} Achieving a closed reduction may sometimes be difficult, requiring special measures that can cause additional trauma, including use of a traction table, or an 'F'-shaped radiolucent reduction device (Fig. 1), or even an open reduction. We describe a novel and effective method of achieving reduction whilst the nails are being inserted.

Figure 1 The fracture reduction 'F' tool.

Figure 2 Intra-operative images demonstrate how traction applied to the nails pulls the fracture out to length and aligns the medullary canal to allow the nails to be advanced across the fracture.

TECHNIQUE

With the patient on a radiolucent table, nails are inserted in a standard retrograde manner into the distal fragment. Once both nails reach the fracture, the surgeon applies longitudinal traction on the nails, which pulls the fracture out to length. There is sufficient friction between the correctly sized nails and endosteal surface that the rods do not back out. The nails can be levered antero-posteriorly or side-to-side until the medullary canals of the distal and proximal fragments are aligned. The surgeon maintains the traction on the nails while the assistant strikes the impacting surface of the chuck handles with a mallet to advance the nails across the fracture (Figs 2 and 3). This overcomes the traction and frictional resistance, advancing the nails in a controlled manner across the fracture without displacing the reduction. The operation is completed in a standard manner, but nail ends should be left flush to the bone to avoid irritation.⁵

DISCUSSION

The advantages of this method are that a traction table is not needed, soft tissues are not injured from direct reduction manoeuvres such as an 'F' tool or open reduction.

References

1. Metaizeau JP. *L'osteosynthese chez l'enfant par embrochage centro medullaire elastique stable*. Montpellier: Sauramps Medical, 1988.
2. Ligier JN, Metaizeau JP, Prévot J, Lascombes P. Elastic stable intramedullary nailing of femoral shaft fractures in children. *J Bone Joint Surg Br* 1988; **70**: 74–7.

A

B

Figure 3 While maintaining traction on the nails, the mallet is used to strike the chuck handles to advance the nails across the fracture.

3. Sanders JO, Browne RH, Mooney JF, Raney EM, Horn BD, Anderson DJ *et al.* Treatment of femoral fractures in children by pediatric orthopedists: results of a 1998 survey. *J Pediatr Orthop* 2001; **21**: 436–41.
4. Flynn JM, Hresko T, Reynolds RAK, Blasier RD, Davidson R, Kasser J. Titanium elastic nails for pediatric femur fractures: a multicenter study of early results with analysis of complications. *J Pediatr Orthop* 2001; **21**: 4–8.
5. Narayanan UG, Hyman JE, Wainwright AM, Rang M, Alman BA. Complications of elastic stable intramedullary nail fixation of pediatric femoral fractures, and how to avoid them. *J Pediatr Orthop* 2004; **24**: 363–9.

A low-cost technique for measuring the intra-abdominal pressure in non-industrialised countries

ADHISH BASU

Jawaharlal Institute of Postgraduate Medical Education and Research, Pondicherry, India

CORRESPONDENCE TO

Adhish Basu, 90 Ballygunge Place, Kolkata 700019, India
T: +91 (0)413 3957934; E: adhishbasu@rediffmail.com

BACKGROUND

A low-cost and simple technique for estimating the intra-abdominal pressure (IAP) indirectly by vesical pressure measurement is described.

TECHNIQUE

Using a Foley catheter (16-F or 18-F), an intravenous infusion set, a 50 ml syringe, a measuring scale and a hemostat, provides a low-cost assessment of the IAP. The connector of the intravenous

Figure 1 Intravenous infusion set tubing used as a saline column manometer