3. Schütze M, Kääb MJ, Haas N. Stabilization of proximal tibial fractures with the LISS-system: early clinical experience in Berlin. *Injury* 2003; **34** (Suppl. 1): A30–5.
4. Stannard JP, Wilson TC, Volgas DA, Alonso JE. Fracture stabilization of proximal tibial fractures with the proximal tibial LISS: early experience in Birmingham, Alabama (USA). *Injury* 2003; **34** (Suppl. 1): A36–42.

A simple surgical technique for removal of radio-opaque foreign objects from the plantar surface of the foot

S SHARMA, T AZZOPARDI
Department of Orthopaedics, Southern General Hospital,
Glasgow, UK

CORRESPONDENCE TO

Mr Sunil Sharma, 47 Shuna Place, Newton Mearns, Glasgow
G77 6TN, UK
T: +44 (0)141 639 3820; E: sssharma2@yahoo.co.uk

INTRODUCTION

Removal of foreign objects such as bits of broken glass or a piece of broken needle from the plantar surface of the foot can be challenging. It is common for the surgeon to end up with a larger than planned surgical incision in pursuit of the elusive foreign object. Despite a long surgical incision, the foreign object may not be found; frequently, the surgeon has to use an image intensifier during the procedure to help locate the foreign object. We describe a simple technique, which could be used for removal of radio-opaque foreign objects from the plantar surface of the foot.

TECHNIQUE

The patient is positioned supine with the foot at the end of the operating table. The foot is prepared and draped as usual. Under image intensifier guidance, three 20-gauge needles are inserted, 2 cm apart, angled at 45° to the plantar surface of the foot so that the tips of the needles are in contact with the radio-opaque foreign object. The needles are inserted around the puncture wound on the plantar surface of the foot. The skin is incised in the triangular region between the needles (Fig. 1). Sharp

Figure 1 The surgical incision is through the centre of the triangular area formed by the needle insertion sites (circles)

dissection carried down to the tips of the needles will identify the foreign object and facilitate easy removal.

CONCLUSION

This is a simple technique for identifying and removing radio-opaque foreign objects from the plantar aspect of the sole.

A technique to assist reversal of abdominal stomas

SHIRLEY CHAN, SAMER DOUGHAN, MICHAEL SAUNDERS
Department of General Surgery, Eastbourne District General
Hospital, King's Drive, Eastbourne, East Sussex, UK

CORRESPONDENCE TO

Miss Shirley Chan, SpR in General Surgery, The Royal
Alexander Hospital for Sick Children, Dyke Road, Brighton
BN2 3JN, UK
T: +44 (0)1273 328145; F: +44 (0)1273 736685;
E: shirley.chan@talk21.com

BACKGROUND

The operative procedure of reversal of abdominal stoma is often underestimated in its complexity but its complications can lead to significant morbidity (e.g. anastomotic leak, wound infection). We describe a technique of abdominal stoma reversal that facilitates dissection of tissue planes and good postoperative analgesia.

TECHNIQUE

The patient is cleaned and draped as normal. The area around the stoma is infiltrated with 40 ml of 0.25% bupivacaine with adrenaline diluted with normal saline until the skin blanches (or the maximum dose allowed by body weight is reached). The skin around the stoma is excised elliptically. The stoma is mobilised down to the peritoneal cavity as normal but with the added benefit of the local anaesthetic to dissect and define tissue planes. The anastomosis is performed and the bowel reduced back into the peritoneal cavity. Closure of the rectus and skin is performed.

DISCUSSION

This technique has the advantages of improved stoma definition as well as easier dissection of tissue planes by 'hydrodissection' of the local anaesthetic infiltration. It also benefits the patient with good regional analgesia in the immediate postoperative period.