Figure 5

TECHNIQUE

With the patient supine, the centre of the bandage cradles the tendo-Achilles. The arms of the bandage are wrapped around the ankle crossing over on the dorsum of the foot (Fig. 1). The throws are then brought around the foot sole and made to cross each other again (Fig. 2). The two arms of the bandage are made to pass under the first throw and brought up proximally. Alternate traction to tighten up the construction/sling (Fig. 3). The direction of the traction is now finally reversed so that it is away from the patient (Fig. 4). The axis of traction is then centralised, securing the traction by multiple simple knots inserted between the two free ends of the bandage which is taken over a sturdy bar and 2–7 kg weights applied (Fig. 5).

DISCUSSION

This is an easy and secure method of traction; no skin problems have been encountered to date. The obvious advantages are that the technique is: (i) inexpensive (15 p); (ii) easy to apply using readily available material; and (iii) non-invasive for the patient.

ACKNOWLEDGEMENT

The authors wish to thank Mr David Higgins for the illustrations.

Reference

1. Budworth G. *The Complete Guide to Knots and Knot Tying*. 1999; 210. (ISBN 0-7548-0422-4).

The use of a polyglactin 910 mesh to obtain haemostasis and prevent further splitting in a fractured transplant kidney

R GANDY, S ASTHANA, KV MENON, N AHMAD, JPA LODGE
Department of Transplant Surgery, St James's University
Hospital, Leeds, UK

CORRESPONDENCE TO

JPA Lodge, Department of Transplantation Surgery, St James's
University Hospital, Beckett Street, Leeds LS9 7TF, UK
E: peterlodge@leedsth.nhs.uk

BACKGROUND

Bleeding due to cortical tears in a transplant kidney is a rare, but potentially serious, complication. Friable and oedematous tissue will often not take sutures and bleeding may not be controlled with fibrin glue, thrombin haemostatic matrix (FloSeal®) or oxidised, regenerated cellulose (Surgicel®). These products will also not prevent further fracturing if the kidney becomes engorged.

TECHNIQUE

The kidney is exposed and inspected for bleeding points. Any haematoma must be evacuated. If the kidney is viable but with cortical tearing, direct pressure is applied. Flowseal can be applied to the fracture site. An appropriately sized absorbable mesh is divided in the midline to a distance of roughly two-thirds of its length. The cut surface of the mesh is laid around the hilum of the kidney with care not to obstruct the vessels or ureter (Fig. 1). The outer edges of the mesh are drawn together over the cortex of the kidney and loosely sutured together using a continuous PDS

Figure 1 A kidney has been wrapped in absorbable mesh after the upper pole has fractured in three places (Surgicel applied to upper pole under mesh).

stitch, so there is no cortical hypoperfusion. The kidney is then re-inspected for vascular insufficiency and two suction drains are placed adjacent to the hilum. The patient must be closely observed postoperatively for evidence of re-bleed and post-operative biopsy must be avoided.

DISCUSSION

This technique is a viable option for haemostasis and the prevention of fracturing in a transplanted kidney. It can be used to avoid transplant nephrectomy without compromising vascularity and function.

References

1. Lee SS, Cheng CL, Yu DS, Chang SY, Ma CP. Vicryl mesh repair of severely injured kidneys: an experimental study. *J Trauma* 1993; **34**: 406–10.
2. Heimbach D, Miersch WD, Buszello H, Schoeneich G, Klehr HU. Is the transplant preserving management of renal allograft rupture justified? *Br J Urol* 1995; **75**: 729–32.

A quick and simple technique for irrigation of open fractures

R LANDSMEER, V TEK, V KHANDUJA

Dennis Dunn Orthopaedic Unit, Colchester General Hospital, Colchester, Essex, UK

CORRESPONDENCE TO

Vikas Khanduja, Knee Fellow, The Wellington & Royal Free Hospital, Wellington Place, London NW8 9LE, UK
T: +44 (0)20 7483 5043; F: +44 (0)20 7483 5241; M: +44 (0)7968 098252; E: vikaskhanduja@aol.com

BACKGROUND

Wound irrigation to remove debris and decrease contamination is an essential step in management of open fractures and wounds.¹ Debate still continues over the best irrigation system. Various systems of high- and low-pressure irrigation have been described but all have their pitfalls.² We describe a quick, simple, cheap and effective method for irrigation of open fractures and soft-tissue injuries.

TECHNIQUE

Our technique requires a sterile 1-l bag of normal saline and a sterile Yankauer sucker (Fig. 1A). The outflow port of the bag is cut and the Yankauer sucker is attached (Fig. 1B). This allows the saline to be well directed and, in addition, the holes at the far end act as a sprinkler system. It is perfect for irrigating open fractures and wounds quickly and gives the ideal low-pressure, high-output system. The flow rate measured via this method is 1.5 l/min.

A

B

Figure 1 (A) A 1-l bag of normal saline and the Yankauer sucker. (B) The Yankauer sucker attached to the saline bag for irrigation system.

DISCUSSION

This technique has been used effectively in our theatres and can also be extended to the accident and emergency department setting. Also, as compared to the other irrigation systems described in the literature³ namely, a 60-ml syringe with 19-gauge needle, a giving set and gravity, a giving set with 400 mmHg and the pulsatile lavage system, our technique has the highest flow rate, thereby allowing copious amounts of fluid to be delivered in a very short span of time.

References

1. Gustilo RB, Anderson JT. Prevention of infection in treatment of one and twenty five open fractures of long bones. *J Bone Joint Surg Am* 1976; **58**: 453–8.
2. Dulecki M, Pieper B, Detroit M. Irrigating simple acute traumatic wounds. A review of the current literature. *J Emerg Nurs* 2005; **31**: 156–60.
3. Karupparsamy K, Hutton MJ, Porteous MJL. A novel system for the irrigation of open fractures. *Injury* 2004; **35**: 980–1.